

Children's Mercy Kansas City

SHARE @ Children's Mercy

Research at Children's Mercy Month 2019

Research at Children's Mercy Month

5-2019

Improved Vitamin D Status Over Time in Children Undergoing Hematopoietic Cell Transplantation

Nancy Shreve

Rakesh Goyal

Vincent S. Staggs

Katrina Walters

Jos Domen

See next page for additional authors

Follow this and additional works at: https://scholarlyexchange.childrensmercy.org/research_month2019

Authors

Nancy Shreve, Rakesh Goyal, Vincent S. Staggs, Katrina Walters, Jos Domen, Jean Freundenthal, Uttam Garg, Nancy Green, Rukhsana Rahmetulla, Nancy Terwilliger, Mohamed Radhi, and Jignesh Dalal

Improved Vitamin D Status over Time in Children Undergoing Hematopoietic Cell Transplantation



Nancy Shreve, MS, APRN, Rakesh K Goyal, MD, Vincent S Staggs, PhD, Katrina Walters, CCRC, Jos Domen, PhD, Jean Freudenthal, PharmD, Uttam Garg, PhD, Nancy Green, Rukhsana Rahmetulla, APRN, Nancy Terwilliger, APRN, Mohamed Radhi, MD and Jignesh D. Dalal, MD
 Children's Mercy, Kansas City, MO, Centene Corporation, St. Louis, MO, and Rainbow Babies and Children's Hospital, Cleveland, OH



BACKGROUND

Vitamin D is an essential nutrient that plays an important role in calcium homeostasis and bone health and has important immunomodulatory effects. Vitamin D deficiency is prevalent in children undergoing hematopoietic cell transplantation (HCT), and has been correlated with risk of post-transplant complications and inferior survival.

OBJECTIVE

To determine vitamin D status pre transplant, at 100 days, 180 days, one year, and two years post-transplant

STUDY DESIGN

A prospective non-interventional, IRB approved study of vitamin D status over time in pediatric HCT patients

METHODS

Patients between >1 mo. and ≤21 y were eligible. Blood samples were drawn to measure 25-hydroxy vitamin D ("vitamin D") levels at pre-HCT, and 100 days, 6 mo., 1 y, and 2 y post-HCT time points. Vitamin D levels were classified as sufficient (≥30 ng/ml), insufficient (20–29 ng/ml) or deficient (<20 ng/ml). All patients were followed by registered dietitians. Logistic mixed models were used to examine vitamin D status across time. Chart Review was completed to determine Vitamin D supplementation.

TABLE 1. BASELINE CHARACTERISTICS BY PRE-HCT VITAMIN D GROUPS

	SUFFICIENT (>30 ng/ml)	INSUFFICIENT (20-30 ng/ml)	DEFICIENT (<20 ng/ml)
TOTAL (N = 66)	29 (43%)	22 (33%)	15 (23%)
AGE RANGE			
<4 y (N = 22)	14 (63%)	5 (23%)	3 (14%)
4-12 y (N = 23)	10 (43.5%)	10 (43.5%)	3 (23%)
>12 (N = 21)	5 (24%)	7 (33%)	9 (48%)
GENDER			
Male (N = 45)	20 (45%)	19 (42%)	6 (33%)
Female (N = 21)	5 (24%)	7 (33%)	9 (43%)
RACE / ETHNICITY			
White (N = 47)	19 (40%)	19 (40%)	9 (20%)
Non-white (N = 19)	7 (37%)	6 (31.5%)	6 (31.5%)
TYPE OF HCT			
Allogeneic (N = 48)	21 (44%)	15 (31%)	12 (25%)
Autologous (N = 18)	3 (17%)	7(39%)	8 (44%)

RESULTS

Sixty-six patients were enrolled, median age 8.1 y (0.4 y -20.5 y). 73% of patients received myeloablative conditioning. The median vitamin D level prior to HCT was 27 ng/ml (range: 8 ng/ml – 69 ng/ml). Levels were insufficient and deficient in 33% and 23% patients (Table 1). Only 24% of children >12 y had sufficient vitamin D levels compared with 44% and 64% in ages 4-12 and below 4 y respectively. Only 24% female patients had normal vitamin D levels compared with 45% male patients.

Compared to pre-HCT, estimated odds of "below sufficient" (either insufficient or deficient) were 47% higher at 100 days, 30% higher at 6 mo., 69% lower at 1 y, and 81% lower at 2 y (linear trend p = .023) and estimated odds of sufficiency were 39% lower at 100 days, 21% lower at 6 mo., and 41-42% higher at 1 y and 2 y (linear trend p = .087)(Figure 1). The proportion of patients receiving supplemental vitamin D were 7.5%, 10%, 19%, 30% and 42% at pre-HCT, +100 days, 6 mo., 1 and 2 y post-HCT time-points, respectively.

CONCLUSIONS

We show a high prevalence rate of vitamin D insufficiency and deficiency prior to and early after HCT. We also found that the odds of vitamin D sufficiency improved over 1 – 2 y post-HCT. We suggest that all pediatric patients should be screened for vitamin D status prior to and at day 100 post-transplant and those with low levels (<20 ng/mL) treated. Clinical significance of this approach needs to be examined in a large multicenter trial.

FIGURE 1. VITAMIN D SUFFICIENCY, INSUFFICIENCY AND DEFICIENCY OVER TIME

