

Children's Mercy Kansas City

## SHARE @ Children's Mercy

---

Research Days

GME Research Days 2019

---

May 16th, 11:30 AM - 1:30 PM

### Initial Evaluation of Adolescent Females Hospitalized with Heavy Menstrual Bleeding

Lauren Amos MD

*Children's Mercy Hospital*

Let us know how access to this publication benefits you

Follow this and additional works at: <https://scholarlyexchange.childrensmercy.org/researchdays>



Part of the [Hemic and Lymphatic Diseases Commons](#), and the [Pediatrics Commons](#)

---

Amos, Lauren MD, "Initial Evaluation of Adolescent Females Hospitalized with Heavy Menstrual Bleeding" (2019). *Research Days*. 13.

[https://scholarlyexchange.childrensmercy.org/researchdays/GME\\_Research\\_Days\\_2019/Research\\_Days\\_four/13](https://scholarlyexchange.childrensmercy.org/researchdays/GME_Research_Days_2019/Research_Days_four/13)

This Poster Presentation is brought to you for free and open access by the Conferences and Events at SHARE @ Children's Mercy. It has been accepted for inclusion in Research Days by an authorized administrator of SHARE @ Children's Mercy. For more information, please contact [hlsteel@cmh.edu](mailto:hlsteel@cmh.edu).

## Initial Evaluation of Adolescent Females Hospitalized with Heavy Menstrual Bleeding

Lauren E. Amos, MD<sup>1</sup>, Shannon L. Carpenter, MD, MS<sup>1</sup>

*<sup>1</sup>Division of Hematology-Oncology-BMT, Children's Mercy Hospital, Kansas City, Missouri*

**Background:** Heavy menstrual bleeding (HMB) in adolescents can be severe and life-threatening. Up to 30% of young women who are hospitalized with anemia due to HMB have a bleeding disorder. Guidelines from the American College of Obstetrics and Gynecology (ACOG) and the National Heart, Lung, and Blood Institute (NHLBI) recommend evaluation for bleeding disorders in such patients. ACOG recommendations include testing for von Willebrand disease (VWD) and specify that consultation with a hematologist may help in interpreting results. NHLBI recommends testing for vWD be done in conjunction with a hematologist. ACOG guidelines do not include testing for platelet function disorders (PFD). Early and accurate diagnosis of bleeding disorders is important, yet limited data exists on the diagnostic evaluation for bleeding disorders in adolescent females hospitalized for HMB.

**Objective:** To evaluate the diagnostic evaluation of bleeding disorders in adolescent females hospitalized for HMB.

**Methods:** A retrospective, single center chart review of female patients aged 9-21 years hospitalized for HMB and anemia at a tertiary care children's hospital from January 1, 2000 until December 31, 2017 was done. Patients were identified from our Hemophilia Treatment Center (HTC) registry, review of patients seen at a comprehensive clinic staffed by pediatric hematologists and gynecologists, and by an Electronic Medical Record (EMR) query of admission and discharge diagnoses of HMB and anemia. Data obtained included clinical features, diagnostic evaluation, and laboratory results.

**Results:** 118 patients hospitalized for HMB and anemia were included. Inpatient Hematology consult or outpatient referral occurred in 68 (58%) of the patients; 60/68 (88%) had a bleeding disorder evaluation completed. 34 patients had a hematologic disorder. PFD was the most common (15/34; 44%) followed by VWD (9/34; 26%). 42% (50/118) of the patients did not have a Hematology consult or outpatient referral (Table 1). While hospitalized for HMB and anemia, 29 of the 50 patients had testing for vWD performed and only 4/29 (14%) had testing repeated as an outpatient once hemoglobin normalized. No patients tested for VWD while inpatient had results consistent with the diagnosis. Platelet function testing was performed in 10/50 patients using the platelet function analyzer (PFA-100) in 8 patients and platelet aggregometry in 2 patients.

**Conclusions:** Despite national guidelines and the presence of known risk factors for bleeding disorders, the hematology service was not involved in the diagnostic process for a significant number of adolescent females. In these patients, testing often occurred while patients were hospitalized and was not repeated. Testing for platelet function disorders occurred infrequently and mainly consisted of the PFA-100 which lacks sensitivity and specificity. When patients were evaluated by Hematology and tested for bleeding disorders, a large proportion had a bleeding disorder, of which PFD were most common. This study demonstrates the need for standardization of the evaluation of adolescent females hospitalized

for HMB. Guidelines should be updated to include testing for PFD. Hematologists should be involved when females are hospitalized for HMB and anemia.