

Children's Mercy Kansas City

SHARE @ Children's Mercy

Posters

5-2021

Impact of Early Tracheostomy on Neurodevelopmental Outcome of Infants with Severe Bronchopulmonary Dysplasia Exposed to Postnatal Steroids

Amjad Taha

Children's Mercy Hospital

Gangaram Akangire

Children's Mercy Hospital

Janelle R. Noel-Macdonnell PhD

Children's Mercy Hospital

Tiffany Willis

Children's Mercy Hospital

Winston M. Manimtim

Children's Mercy Hospital

Let us know how access to this publication benefits you

Follow this and additional works at: <https://scholarlyexchange.childrensmercy.org/posters>



Part of the Behavioral Medicine Commons, and the Pediatrics Commons

Recommended Citation

Taha, Amjad; Akangire, Gangaram; Noel-Macdonnell, Janelle R. PhD; Willis, Tiffany; and Manimtim, Winston M., "Impact of Early Tracheostomy on Neurodevelopmental Outcome of Infants with Severe Bronchopulmonary Dysplasia Exposed to Postnatal Steroids" (2021). *Posters*. 190.

<https://scholarlyexchange.childrensmercy.org/posters/190>

This Poster is brought to you for free and open access by SHARE @ Children's Mercy. It has been accepted for inclusion in Posters by an authorized administrator of SHARE @ Children's Mercy. For more information, please contact hlsteel@cmh.edu.

Impact of Early Tracheostomy on Neurodevelopmental Outcome of Infants with Severe Bronchopulmonary Dysplasia Exposed to Postnatal Steroids

Amjad Taha, MD^{1,2}, Gangaram Akangire, MD^{1,2}, Janelle Noel-MacDonnell PhD¹, Tiffany Willis, PsyD^{1,2}, Winston Manimtim, MD^{1,2}

¹Children's Mercy Kansas City; ²University of Kansas City School of Medicine

Introduction

- Infants with severe bronchopulmonary dysplasia (sBPD) are at high risk for neurodevelopmental (ND) impairment
- Prolonged assisted ventilation through a tracheostomy in infants with sBPD may improve ND outcome
- There is no consensus on optimal timing of tracheostomy
- There is evidence that tracheostomy performed at <120 days of age improves ND outcome at 18-22 months of age
- Postnatal steroid (PS) use in BPD had been shown to have negative effect on neurodevelopment
- To date, no data has specifically evaluated the impact of early tracheostomy on ND outcome of infants with severe BPD who are exposed to postnatal steroids

Objective

- To compare the cognitive, language and motor scores among infants with sBPD who received Early (ET) versus Late (LT) versus No tracheostomy (NT)
- Secondly, to evaluate if PS had an additive negative effect on ND outcomes

Design/Methods

- IRB approved retrospective review, 2010-2017
- Infants with sBPD divided into ET (< 121 days of age), LT (≥ 121 days) and NT
- Primary outcome: Cognitive, Language and Motor scores at 2-3 years of age (*Bayley's 3rd edition*)
- Secondary outcome: Cumulative PS use of each group
- Statistics: *Kruskal Wallis test* (nonparametric ANOVA)


Results

Table 1. Patient Demographics with steroids exposure

Variables	Early Tracheostomy n=22	Late Tracheostomy n=19	No Tracheostomy n=27	P Value
Gestational age at birth (weeks) ^a	26(24.00,28.00)	25.00 (24.00, 28.00)	25.00 (24.00, 28.00)	0.967
Birth weight (grams) ^a	840.00 (630.00, 1069.00)	710.00 (600.00, 895.00)	845.00 (710.00, 995.00)	0.138
Gender ^b				< 0.001
Male	12 (54.5%)	8 (42.1%)	10 (37.0%)	
Maternal Race ^c				< 0.001
White	19 (100.0%)	13 (68.4%)	11 (40.7%)	
Black/African American	0 (0.0%)	3 (15.8%)	12 (44.4%)	
Hispanic	0 (0.0%)	0 (0.0%)	3 (11.1%)	
Other	0 (0.0%)	3 (15.7%)	1 (3.7%)	
Unknown	0	0		
Age on admission (days of life) ^a	65.00 (1.00, 92.00)	103.00 (0.00, 129.00)	1.00 (0.00, 32.00)	0.005
Age at tracheostomy (Day of life) ^a	94.00 (82.00, 105.00)	148.00 (134.00, 168.00)		< 0.001
Age at discharge (days of life) ^a	190.00 (159.00, 255.00)	251.00 (210.00, 275.00)	125.00 (110.00, 161.00)	< 0.001
Age at discharge (weeks PMA) ^a	53.50 (49.10, 60.70)	61.50 (56.10, 65.60)	43.86 (41.43, 53.29)	< 0.001
Steroid exposure prior to admission ^b	14 (63.6%)	11 (57.9%)	5 (18.5%)	0.002
Dexamethasone exposure ^b	18 (81.8%)	15 (78.9%)	26 (96.3%)	0.160
Cumulative steroid exposure (hydrocortisone equivalent) ^a	347.20 (132.95, 677.00)	595.05 (67.50, 1213.60)	97.90 (35.60, 237.50)	0.012
PDA ^b	18 (81.8%)	16 (84.2%)	14 (51.9%)	0.022
IVH ^b	6 (27.3%)	8 (42.1%)	10 (37.0%)	0.594

PMA postmenstrual age, PDA patent ductus arteriosus, IVH intraventricular hemorrhage
^aData presented as median (interquartile range)
^bData presented as number (percentage)

- Motor composite scores: ET vs LT (median score 85 vs 73, *P* 0.028)
- Trend for better cognitive scores in ET vs LT vs NT but not significant
- No difference in language scores among 3 groups
- LT had the lowest scores in all 3 domains
- LT had the most PS; NT had the least PS



Conclusion

- Early tracheostomy may improve neurodevelopmental outcome in infants with severe BPD particularly in the **motor domain**
- Delaying tracheostomy in severe BPD may predispose to more postnatal steroids exposure and possible worst neurodevelopmental impairment