

Children's Mercy Kansas City

SHARE @ Children's Mercy

Research Days

GME Research Days 2021

May 12th, 11:30 AM - 1:30 PM

Bar Removal Following Minimally Invasive Pectus Excavatum Repair – Does Removal at 2 Years Affect Recurrence or Satisfaction Rates?

Wendy Jo Svetanoff
Children's Mercy Hospital

Let us know how access to this publication benefits you

Follow this and additional works at: <https://scholarlyexchange.childrensmercy.org/researchdays>



Part of the [Pediatrics Commons](#), [Surgery Commons](#), and the [Surgical Procedures, Operative Commons](#)

Svetanoff, Wendy Jo, "Bar Removal Following Minimally Invasive Pectus Excavatum Repair – Does Removal at 2 Years Affect Recurrence or Satisfaction Rates?" (2021). *Research Days*. 13.
https://scholarlyexchange.childrensmercy.org/researchdays/GME_Research_Days_2021/researchday3/13

This Poster Presentation is brought to you for free and open access by the Conferences and Events at SHARE @ Children's Mercy. It has been accepted for inclusion in Research Days by an authorized administrator of SHARE @ Children's Mercy. For more information, please contact hlsteel@cmh.edu.

Bar Removal Following Minimally Invasive Pectus Excavatum Repair – Does Removal at 2 Years Affect Recurrence or Satisfaction Rates?

Wendy Jo Svetanoff MD MPH; James A Fraser MD; Kayla B Briggs MD; Pablo Aguayo MD; Charles Snyder MD; David Juang MD; Jason D Fraser MD; Tolulope A Oyetunji MD MPh; Shawn D St. Peter MD

Children's Mercy Kansas City; University of Missouri Kansas City School of Medicine

Introduction

- Pectus excavatum is the most common congenital chest wall deformity.
- During minimally invasive pectus excavatum repair (MIRPE), a bar is placed to correct the deformity.
- The bar normally stay in place for 3 years.
- Little is known about recurrence or cosmetic satisfaction when the bar is removed early.

Aim

- Review patients who underwent bar removal less than 3 years after placement:
 - Recurrence Rate
 - Patient Satisfaction


Methods

- Retrospective Review with Prospective Follow-up
- Inclusion Criteria:
 - MIRPE between October 2006 – June 2017
 - Had bar removed < 3 years after placement
- Telephone Follow-up
 - Residual Deformity
 - Cosmetic Satisfaction

Results

44 patients included
 Haller Index: 4.0 (IQR 3.5, 4.5)
 Median time at bar placement: 13.3 years
 Median time to bar removal: 2.1 years

Reasons for Early Bar Removal




- No Reason Identified/Doing Well
- Over-correction
- Leaving Parents' Home
- Other
- Infection
- Loss of Insurance

Post-operative Outcomes	Patients (n=44)
Recurrent Deformity	2
- Required Bracing	1
- Required Redo Repair	1
Bracing for Pectus Carinatum	1
Costal Cartilage Resection	3
Scar Revision	1

Telephone Surveys (n=17):

- *Median follow-up: 9 years
- *9 patients (53%) noted residual deformity
 - *2 thought deformity was >25% of chest
 - *1 sought re-evaluation (bracing)

Patient Satisfaction Rates



Conclusion

- Early bar removal does not increase recurrence rates requiring surgical intervention and maintains good patient cosmetic satisfaction.
 - More long-term follow-up information for all pectus patients is warranted.