

Children's Mercy Kansas City

## SHARE @ Children's Mercy

---

Posters

---

5-2021

### Constrictive Pericarditis After Repair of a Ruptured Sinus of Valsalva

Sarah Studyvin

*Children's Mercy Hospital*

Sanket Shah

*Children's Mercy Hospital*

Let us know how access to this publication benefits you

Follow this and additional works at: <https://scholarlyexchange.childrensmercy.org/posters>



Part of the [Cardiology Commons](#)

---

#### Recommended Citation

Studyvin, Sarah and Shah, Sanket, "Constrictive Pericarditis After Repair of a Ruptured Sinus of Valsalva" (2021). *Posters*. 204.

<https://scholarlyexchange.childrensmercy.org/posters/204>

This Poster is brought to you for free and open access by SHARE @ Children's Mercy. It has been accepted for inclusion in Posters by an authorized administrator of SHARE @ Children's Mercy. For more information, please contact [hlsteel@cmh.edu](mailto:hlsteel@cmh.edu).

# Constrictive Pericarditis After Repair of a Ruptured Sinus of Valsalva

Sarah Studyvin, DO, MPH; Sanket S. Shah, MD  
Children's Mercy Hospital, Kansas City, MO

## BACKGROUND

Constrictive pericarditis is an uncommon complication of cardiac surgery. We report a patient who developed constrictive pericarditis after ruptured sinus of Valsalva (RSOV) repair.

## CASE

A 23-year-old male presented with exertional dyspnea one year after RSOV repair. TTE showed a small, circumferential effusion with thickened pericardium, ventricular septal bounce (Fig 1), left atrial enlargement, diastolic hepatic flow reversal (Fig 2), and trivial mitral regurgitation without stenosis. He underwent cardiac catheterization, which revealed elevated filling pressures (RVEDP 16 mmHg, LVEDP 18 mmHg), RVEDP/RVSP ratio < 0.5, and a low cardiac index (1.65 L/min/m<sup>2</sup>). Cardiac MRI confirmed pericardial thickening (Fig 3) with paradoxical septal motion, dilated pulmonary veins (Fig 4) and retrograde flow in the SVC.

## DECISION-MAKING

Pericardiectomy of thickened and adherent pericardium was performed. The central venous pressure decreased from 23 to 7 mmHg and TEE showed normal systolic function with less septal bounce postoperatively. Pathology specimens of the pericardium exhibited fibrosis and mild chronic inflammation. He continued to do well at one-month follow-up.

## CONCLUSION

Constrictive pericarditis is a rare complication of cardiac surgery. In patients presenting with right sided heart failure and deterioration of cardiac function not explained by other mechanisms, evaluation with TTE and cMRI should be considered; cardiac catheterization with or without fluid challenge can be performed to confirm the diagnosis.


Consider the diagnosis of **constrictive pericarditis** in patients with unexplained **right heart failure and decreased cardiac function.**

Evaluate for constrictive pericarditis with **echocardiography and cardiac MRI.**

For more information email  
[scstudyvin@cmh.edu](mailto:scstudyvin@cmh.edu)


ACC.21

FIGURE 1




Thickened pericardium and septal bounce with deep inspiration on TTE.

FIGURE 2




Diastolic hepatic flow reversal on TTE.

FIGURE 3



Pericardial thickening on cMRI.

FIGURE 4



Dilated pulmonary veins on cMRI.

## DISCLOSURE INFORMATION

Nothing to disclose.