

Children's Mercy Kansas City

SHARE @ Children's Mercy

Posters

9-2021

Improving Skin and Soft Tissue Antibiotic Duration Concordance with National Guidelines in Pediatric Urgent Care Clinics

Megan Hamner

Children's Mercy Hospital

Amanda Nedved

Children's Mercy Hospital

Holly Austin

Children's Mercy Hospital

Donna Wyly

Children's Mercy Hospital

Alaina N. Burns

Children's Mercy Hospital

Let us know how access to this publication benefits you
~~See next page for additional authors~~

Follow this and additional works at: <https://scholarlyexchange.childrensmercy.org/posters>



Part of the [Emergency Medicine Commons](#), [Infectious Disease Commons](#), [Pediatrics Commons](#), and the [Pharmacy and Pharmaceutical Sciences Commons](#)

Recommended Citation

Hamner, Megan; Nedved, Amanda; Austin, Holly; Wyly, Donna; Burns, Alaina N.; King, Diana; Lee, Brian; and El Feghaly, Rana, "Improving Skin and Soft Tissue Antibiotic Duration Concordance with National Guidelines in Pediatric Urgent Care Clinics" (2021). *Posters*. 248.

<https://scholarlyexchange.childrensmercy.org/posters/248>

This Poster is brought to you for free and open access by SHARE @ Children's Mercy. It has been accepted for inclusion in Posters by an authorized administrator of SHARE @ Children's Mercy. For more information, please contact hlsteel@cmh.edu.

Authors

Megan Hamner, Amanda Nedved, Holly Austin, Donna Wyly, Alaina N. Burns, Diana King, Brian Lee, and Rana El Feghaly

Improving Skin and Soft Tissue Antibiotic Duration Concordance with National Guidelines in Pediatric Urgent Care Clinics

Megan Hamner¹ MD; Amanda Nedved¹ MD; Holly Austin¹ MD; Donna Wyly¹ RN, MSN, CPNP-AC, PPCNP-BC; Alaina Burns¹ PharmD, BCPPS; Diana King¹ RN, MSN, CPNP-PC; Brian Lee² PhD, MPH, and Rana E. El Feghaly¹ MD, MSCI

¹ Pediatrics, Children's Mercy Hospitals and Clinics, Kansas City, MO; ² Health Services and Outcomes Research, Children's Mercy Hospitals and Clinics, Kansas City, MO

BACKGROUND

- Skin and soft tissue infections (SSTIs) are the second most common diagnoses leading to antibiotic prescriptions
- Guidelines recommend 5-7 days of antibiotics for most SSTIs
- 58% of patients with SSTIs at CMH urgent care clinics (UCCs) received 5-7 days of oral antibiotics

METHODS

- Multidisciplinary team: UCC and ID providers, pharmacist, data analyst
- Reviewed charts of all patients diagnosed with SSTIs in our UCCs
- Cause-and-effect analyses and driver diagram (Fig 1)
- PICK chart helped identify interventions

AIM

Increase the percentage of patients receiving 5-7 days of oral antibiotics for SSTIs from 58% to 75% by December 31st, 2021

METHODS

- **PDSA cycles:**
 - #1: Educational session
 - #2: Updated prescription sentences in electronic health record
 - #3: Project update to UCC providers
- **Measures:**
 - Outcome: % of patients receiving 5-7 days of oral antibiotics for SSTIs
 - Process: # of updated prescriptions used
 - Balancing: # of patients returning for SSTIs within 14 days of initial visit

Figure 1: Driver Diagram





Figure 2: Annotated Control Chart



RESULTS

- Outcome measure:
 - After project initiation (04/2020)-up to 68%
 - PDSA #1 (10/2020-12/2020)-up to 75%
 - PDSA #2 (02/2021)-up to 80%
 - PDSA #3 (04/2021)-up to 90%
- Balancing measure stable

CONCLUSION

- Using QI methodology, we increased the percentage of patients receiving appropriate antibiotic durations for SSTIs from 58% to over 80%
- We are planning additional PDSA cycles and expansion to other departments
- Planning expansion of stewardship efforts to other diagnoses