

Children's Mercy Kansas City

SHARE @ Children's Mercy

Posters

9-2022

Impact of Remote Monitoring During the Interstage Period on Outcomes in Single Ventricle Patients Across Socioeconomic Groups

Bianca Cherestal

Lori Erickson

Janelle R. Noel-Macdonnell

Girish S. Shirali

Hayley S. Hancock

See next page for additional authors

Follow this and additional works at: <https://scholarlyexchange.childrensmercy.org/posters>



Part of the [Cardiology Commons](#), and the [Pediatrics Commons](#)

Authors

Bianca Cherestal, Lori Erickson, Janelle R. Noel-Macdonnell, Girish S. Shirali, Hayley S. Hancock, Doaa Aly, and Natalie Jayaram

Impact of Remote Monitoring During the Interstage Period on Outcomes in Single Ventricle Patients Across Socioeconomic Groups

Bianca Cherestal, MD; Lori A. Erickson, PhD, RN, MSN, CPNP-PC; Janelle R. Noel-MacDonnell, PhD; Girish Shirali, MBBS; Hayley Hancock, MD, FAAP; Doaa Aly, MBBCH; Natalie Jayaram, MD


Children's Mercy Kansas City

Background

- Despite improving outcomes, morbidity and mortality for single ventricle (SV) infants remains high.
- Infants of low socioeconomic status (SES) are known to be particularly vulnerable following stage 1 palliation.
- **Aim:** To investigate whether use of a novel remote monitoring program, CHAMP® (Cardiac High Acuity Monitoring Program), would mitigate the known disparate outcomes for lower SES SV infants during the interstage period (ISP).
- **Hypothesis:** Interstage outcomes for SV infants are the same across differing SES tertiles.

Methods

- **Data Source:** CHAMP® Database
 - 607 SV interstage infants, across 11 institutions (2014-2021) were included in the analysis.
 - Enrollees download CHAMP app to their own device or were provided an iPad or tablet (with built in cellular and video capability) for instantaneous transfer of input data to the care team.



- **Outcome:** Death or transplant listing during the interstage period.
- Patients were divided into SES tertiles based on a neighborhood summary score (Table 1) which is derived from six unique variables relating to SES.
- **Statistical Analysis:** Baseline characteristics between tertiles were compared using Kruskal-Wallis tests for continuous variables and chi-square or Fisher's exact tests for categorical variables (Table 2).
 - Hierarchical logistic regression adjusted for potential confounding characteristics

Table 1

Neighborhood Summary Score	
Median household income	
Median value of housing units	
Households with interest, dividend, or rental income	
Adult residents who completed high school	
Adult residents who completed college	
Employed residents with executive, managerial, or professional occupations	

Table 2

	Tertiles			P-value
	Lowest N = 198	Middle N = 213	Upper N = 198	
Demographic Characteristics				
Female, n (%)	74 (37.6)	76 (35.7)	70 (35.7)	0.905
Non-White race, n (%)	40 (20.2)	39 (18.3)	41 (20.7)	0.811
Hispanic/Latino, n (%)	41 (21.2)	27 (13)	24 (12.4)	0.027
Private Insurance, n (%)	53 (27.6)	96 (47.1)	110 (57.3)	<0.001
Neighborhood Summary Score (range)	-10.84 to -1.56	-1.55 to 1.23	1.25 to 13.53	
Birth Characteristics				
Prenatal Diagnosis, n (%)	162 (81.8)	174 (82.5)	170 (86.7)	0.356
Gestational Age (mean, weeks), n	38.13	38.18	38.09	0.379
Birth Weight (mean, kg), n	3.19	3.13	3.17	0.530
Clinical Characteristics				
Anatomy – HLHS, n (%)	63 (32)	81 (38.2)	77 (39.3)	0.265
Genetic Syndrome, n (%)	160 (80.8)	172 (80.8)	158 (80.6)	0.999
Other Anomalies, n (%)	172 (86.9)	186 (87.3)	172 (87.8)	0.966
Predischarge AVVR*, n (%)	86 (43.6)	91 (42.9)	97 (50)	0.723
Predischarge Function – normal, n (%)	183 (93.4)	198 (93.8)	179 (91.8)	0.942
Interstage Period (mean, days), n	165.46	155.15	146.66	0.104
Outcome				
Glenn, n (%)	187 (94.4)	192 (90.1)	184 (93.9)	0.298
Death, n (%)	8 (4)	15 (7)	6 (3.1)	
Transplant Listing, n (%)	3 (1.5)	6 (2.8)	6 (3.1)	

*AVVR = Atrioventricular valve regurgitation that was mild or greater on predischarge echocardiogram.


Table 3

	Outcome N = 44	Glenn N = 563	P value
Non-Hispanic/Non-Latino, n (%)	42 (95.5)	460 (83.6)	0.037
Renal failure following stage 1 palliation, n (%)	2 (4.5)	2 (0.4)	0.028
Ventricular dysfunction predischarge, n (%)	7 (16.2)	32 (5.8)	0.034
Predischarge AVVR*, n (%)	30 (69.8)	244 (43.5)	<0.001
Lowest tertile, n (%)	11 (25)	187 (33.2)	
Middle tertile, n(%)	21 (47.7)	192 (34.1)	0.185
Upper tertile, n (%)	12 (27.3)	184 (32.7)	

Results

- Of the 607 SV infants included, 44 (7.2%) met the primary outcome.
- **Univariate Analysis:** Non-Hispanic/Non-Latino patients, patients with pre-discharge ventricular dysfunction, post-op renal failure, or post-op AVVR were more likely to experience the primary outcome (Table 3). Rate of reaching outcome did not correlate with SES tertile (Table 3).
- **Multivariable Analysis:** Even after multivariable adjustment for potentially confounding factors, SES was not associated with death/needing transplant.
- The odds of reaching the outcome were no different for those in the middle or upper tertile when compared to the lowest (Figure 1).

Figure 1



Conclusion

- In this large cohort of SV infants enrolled in a digital remote monitoring program during the ISP, we found no difference in outcomes based upon SES.
- These findings differ from prior studies showing worse outcomes for SV patients of lower SES.
- Our study suggests this novel technology could help mitigate differences in outcomes for this fragile population of patients.