

Children's Mercy Kansas City

SHARE @ Children's Mercy

Posters

10-2022

Alterations in the Metabolic Comorbidities of Youth with Overweight and Obesity during the COVID-19 Pandemic

Safa Farrukh

Erica Wee

Children's Mercy Hospital

Ashley Sherman

Children's Mercy Hospital

Yun Yan

Children's Mercy Hospital

Follow this and additional works at: <https://scholarlyexchange.childrensmercy.org/posters>



Part of the [Endocrinology, Diabetes, and Metabolism Commons](#), and the [Pediatrics Commons](#)

Recommended Citation

Farrukh, Safa; Wee, Erica; Sherman, Ashley; and Yan, Yun, "Alterations in the Metabolic Comorbidities of Youth with Overweight and Obesity during the COVID-19 Pandemic" (2022). *Posters*. 294.

<https://scholarlyexchange.childrensmercy.org/posters/294>

This Poster is brought to you for free and open access by SHARE @ Children's Mercy. It has been accepted for inclusion in Posters by an authorized administrator of SHARE @ Children's Mercy. For more information, please contact hlsteel@cmh.edu.

Alterations in the Metabolic Comorbidities of Youth with Overweight and Obesity during the COVID-19 Pandemic

Safa Farrukh¹, Erica Wee, MD², Ashley K. Sherman², Yun Yan, MD^{1,2}

¹University of Missouri-Kansas City School of Medicine

²Children's Mercy Kansas City

Background


- During the COVID-19 pandemic, the time spent in sedentary activities increased.
- The sedentary lifestyle may exacerbate the metabolic comorbidities in youth with overweight and obesity.
- The progression of metabolic comorbidities in youth with overweight and obesity during the COVID-19 pandemic has not been well investigated.
- **Objective:** To evaluate the metabolic comorbidities of youth with overweight and obesity during the COVID-19 pandemic.

Methodology

- A retrospective chart review was conducted of youths seen in a pediatric endocrinology clinic for type 2 diabetes prevention.
- Age, sex at birth, race, type of insurance, BMI, and laboratory results (HbA1c, LDL, HDL, ALT) were collected pre-pandemic and post-pandemic onset.
- Fisher's exact or Chi-squared test and t-test were used to evaluate metabolic comorbidities and demographics traits.

Results

- A total of 74 patients were assessed with mean pre-pandemic age of 12.5 years and was 56.8% female, 39.2% white and 59.2% publicly insured.
- The overall mean BMI increased from 2.5 (37.1 kg/m²) pre-pandemic to 2.6 (39.8 kg/m²) post-pandemic, although the increase was not statistically different (p=0.45).



- HbA1c progression was significantly seen in older youths with mean age of 14.1 years as compared to those without HbA1c progression with mean age of 12.3 years (p=0.04).
- LDL progression was more common in Hispanic race (83.3%) as compared to other races (p=0.01)
- Those with HbA1c progression were more likely to have LDL progression (p=0.03).

Summary

- Progression of metabolic comorbidities among youth with overweight and obesity were common during the COVID-19 pandemic despite no significant increase in BMI.
- HbA1c progression was more common in older youths with overweight and obesity during the COVID-19 pandemic.
- LDL progression was more common among Hispanic youths with overweight and obesity during the COVID-19 pandemic.
- There was a strong correlation between HbA1c progression and LDL progression.

Acknowledgements

We want to acknowledge Brent Lockee and Mitchell S. Barnes who assisted with data collection.

Author's Disclosure

Authors have no financial relationships to disclose.