

Children's Mercy Kansas City

SHARE @ Children's Mercy

Manuscripts, Articles, Book Chapters and Other Papers

8-1-2009

Functional state following the Fontan procedure.

Ismee A. Williams

Lynn A. Sleeper

Steven D. Colan

Minmin Lu

Elizabeth A. Stephenson

See next page for additional authors

Let us know how access to this publication benefits you

Follow this and additional works at: <https://scholarlyexchange.childrensmercy.org/papers>



Part of the [Cardiology Commons](#), [Cardiovascular System Commons](#), [Congenital, Hereditary, and Neonatal Diseases and Abnormalities Commons](#), [Pediatrics Commons](#), [Surgery Commons](#), and the [Surgical Procedures, Operative Commons](#)

Recommended Citation

Williams, I. A., Sleeper, L. A., Colan, S. D., Lu, M., Stephenson, E. A., Newburger, J. W., Gersony, W. M., Cohen, M. S., Cnota, J. F., Atz, A. M., Williams, R. V., Margossian, R., Powell, A. J., Stylianou, M. P., Hsu, D. T., Shirali, G. S. Functional state following the Fontan procedure. *Cardiology in the young* 19, 320-330 (2009).

This Article is brought to you for free and open access by SHARE @ Children's Mercy. It has been accepted for inclusion in Manuscripts, Articles, Book Chapters and Other Papers by an authorized administrator of SHARE @ Children's Mercy. For more information, please contact hlsteel@cmh.edu.

Creator(s)

Ismee A. Williams, Lynn A. Sleeper, Steven D. Colan, Minmin Lu, Elizabeth A. Stephenson, Jane W. Newburger, Welton M. Gersony, Meryl S. Cohen, James F. Cnota, Andrew M. Atz, Richard V. Williams, Renee Margossian, Andrew J. Powell, Mario P. Stylianou, Daphne T. Hsu, Pediatric Heart Network Investigators, and Girish S. Shirali

Published in final edited form as:

Cardiol Young. 2009 August ; 19(4): 320–330. doi:10.1017/S1047951109990382.

Functional state following the Fontan procedure

Ismee A. Williams¹, Lynn A. Sleeper², Steven D. Colan³, Minmin Lu², Elizabeth A. Stephenson⁴, Jane W. Newburger³, Welton M. Gersony¹, Meryl S. Cohen⁵, James F. Cnota⁶, Andrew M. Atz⁷, Richard V. Williams⁸, Renee Margossian³, Andrew J. Powell³, Mario P. Stylianou⁹, and Daphne T. Hsu¹ For the Pediatric Heart Network Investigators (see Appendix)

¹ Department of Pediatrics, Division of Cardiology, Columbia University Medical Centre, New York, New York, United States of America

² New England Research Institutes, Watertown, Massachusetts, United States of America

³ Department of Cardiology, Children's Hospital Boston, Boston, Massachusetts, United States of America

⁴ Department of Pediatrics, Division of Cardiology, University of Toronto, The Hospital for Sick Children, Toronto, Canada

⁵ Department of Pediatrics, Division of Cardiology, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, United States of America

⁶ Department of Pediatric Cardiology, Wake Forest University School of Medicine, Winston-Salem, North Carolina, United States of America

⁷ Department of Pediatrics, Division of Cardiology, Medical University of South Carolina, Charleston, South Carolina, United States of America

⁸ Department of Pediatrics, Division of Cardiology, University of Utah, Salt Lake City, Utah, United States of America

⁹ National Heart, Lung, and Blood Institute, Bethesda, MD, United States of America

Abstract

Background—Despite improvements in outcomes after completion of the Fontan circulation, long-term functional state varies. We sought to identify pre- and postoperative characteristics associated with overall function.

Methods and Results—We analyzed data from 476 survivors with the Fontan circulation enrolled in the Pediatric Heart Network Fontan Cross-sectional Study. Mean age at creation of the Fontan circulation was 3.4 plus or minus 2.1 years, with a range from 0.7 to 17.5 years, and time since completion was 8.7 plus or minus 3.4 years, the range being from 1.1 to 17.3 years. We calculated a functional score for the survivors by averaging the percentile ranks of ventricular ejection fraction, maximal consumption of oxygen, the physical summary score for the Child Health Questionnaire, and a function of brain natriuretic peptide. The mean calculated score was 49.5 plus or minus 17.3, with a range from 3 to 87. After adjustment for time since completion of the circulation, we found that a lower score, and hence worse functional state, was associated with: right ventricular morphology (p less than 0.001), higher ventricular end-diastolic pressure (p equals 0.003) and lower saturations of oxygen (p equals 0.047) prior to completion of the Fontan

circulation, lower income for the caregiver (p equals 0.003), and, in subjects without a prior superior cavopulmonary anastomosis, arrhythmias after completion of the circulation (p equals 0.003). The model explained almost one-fifth (18%) of the variation in the calculated scores. The score was not associated with surgical centre, sex, age, weight, fenestration, or the period of stay in hospital after completion of the Fontan circuit. A validation model, using 71 subjects randomly excluded from initial analysis, weakly correlated (R equals 0.17, p equals 0.16) with the score calculated from the dataset.

Conclusions—Right ventricular morphology, higher ventricular end-diastolic pressure and lower saturations of oxygen prior to completion of the Fontan circuit, lower income for the provider of care, and arrhythmias after creation of the circuit, are all associated with a worse functional state. Unmeasured factors also influence outcomes.

Keywords

Clinical outcomes; single ventricle; univentricular heart

Advances in medical and surgical techniques have increased the survival of patients with functionally univentricular heart into adolescence and young adulthood following the final stage of palliation, namely creation of the Fontan circulation. While patients with this circulation may lead relatively normal lives, their exercise tolerance and functional state often decline over time. Patients with a failing circuit may manifest signs and symptoms of exercise intolerance, protein-losing enteropathy, intractable arrhythmias, poor growth, cyanosis, or recurrent effusions. Although ventricular dysfunction may underlie these symptoms, patients with normal ventricular function may also be affected. As the population with the functionally univentricular circulation ages, an increasing number are being referred for cardiac transplantation due to their poor functional state.¹ Unfortunately, poor nutritional state, or end-organ dysfunction, increases the risk of death while waiting for, or following, transplantation. Identification of factors associated with a worse functional state following creation of the Fontan circuit may allow the early recognition of patients at risk for poor outcome, permitting appropriate intervention and earlier referral for evaluation for transplantation. The aim of our analysis, therefore, was to identify characteristics of the patients before and after conversion to the functionally univentricular circulation associated with a prospectively collected measure of overall functional state, analysing a well characterized cohort of patients who have undergone the Fontan operation.

Methods

Study Design

The National Heart, Lung, and Blood Institute Pediatric Heart Network Fontan Cross-sectional Study, a multi-centric assessment completed in 2004 by the 7 network centres, enrolled 546 survivors with the Fontan circulation, aged from 6 to 18 years. Details of the overall study have been published previously.^{2–4} In brief, 4 clinically useful and independent measures of functional state, specifically ventricular function, exercise performance, health-related measures for quality of life, and evidence of neurohormonal activation, were measured in subjects at a mean of 8.7 plus or minus 3.4 years after completion of the circuit, with the goal of determining the relationship among the variables collected. At the time of enrollment, we recorded the medical history related to the underlying cardiac anatomy, clinical, and echocardiographic state prior to completion of the circuit, the surgical outcomes, and the course after completion of the functionally univentricular circulation. Each centre obtained approval from the local ethics review committee, and informed consent was obtained for each subject.

Assessment of ventricular function

Ventricular ejection fraction as assessed by echocardiography provided a quantitative measure of ventricular function. In order to justify the use of echocardiographic calculated ejection fractions in the population, we compared the ejection fractions in a 125 subjects who underwent both magnetic resonance imaging and echocardiography, and found the absolute mean difference between modalities to be less than 2%. Imaging was performed at each participating centre using a standard protocol, and all measurements were conducted in a standardized fashion by the Core Laboratory of the Fontan Cross-Sectional Study. Cross-sectional echocardiograms and Doppler evaluations of standard short- and long-axis views of the ventricle or ventricles were interpreted by one of two readers. Ejection fraction was calculated using end-diastolic and end-systolic volumes obtained using the biplane-modified Simpson's method.

Assessment of exercise capacity

A maximal bicycle ergometry exercise test was performed using a standard ramp protocol across all centres. We recorded the percentage predicted maximal consumption of oxygen normalized for age and sex.^{5,6}

Assessment of perceived functional state of health

The Child Health Questionnaire Parent Report Form⁷ is a validated self-administered instrument for parents to report on the general health and functioning of their child. It yields two summary scores, one physical and the other psychosocial. We used the physical score, which includes questions about functional limitations at school and at home, physical discomfort, and overall health, in our analysis.

Assessment of neurohormonal activation

We measured centrally the resting concentration of brain natriuretic peptide in the plasma using the Shionogi brain natriuretic peptide-32 human assay.⁸

Statistical methods

Functional Score—In an effort to characterize the overall functional state at the time of participation in the study, we created a functional score. This score is novel, and was developed over the course of multiple discussions by 15 representatives, including both physicians and statisticians, of the 7 participating centres, the New England Research Institute, and the National Heart, Lung, and Blood Institute. The components were determined by consensus from first principles prior to analysis of study data, and are considered to represent clinically useful and independent measures of the functional state at any given time in the period of follow-up. The composite score was calculated for each subject by averaging the percentile rank, determined relative to the other participants in the study, of the four main assessments collected during the study, namely ventricular ejection fraction measured by echocardiography, percentage predicted maximal consumption of oxygen on exercise testing, the physical summary score on the Child Health Questionnaire, and the negative of the residual of age- and gender-adjusted concentration of brain natriuretic peptide in the serum. Equal weight was attributed to each component of the score. The scale ranged from 0 to 100, with a higher score indicative of better functional state.

As an example, the severity score for an individual subject would be calculated as follows:

Percentage predicted maximal consumption of oxygen	57th percentile
Physical score for the Child Health Questionnaire	49th percentile
Negative brain natriuretic peptide	66th percentile

Total ejection fraction

33th percentile

The overall score equals 57 plus 49 plus 66 plus 33 divided by 4, in other words, 51.3.

Due to known effects of gender on the levels of brain natriuretic peptide, we adjusted the raw values prior to calculating the score. Since an adequate external reference does not exist for children, we normalized the values to take account of age and sex, and we used the percentile rank of the residuals to calculate the score. The equation used for adjustment is:

Normalized brain natriuretic peptide equals

$$\log(\text{brain natriuretic peptide}) + 0.0471 \\ * (11.86 - x_1) - 0.21127 * (0.603 - x_2)$$

where x_1 is age at enrollment, 11.86 is mean age, x_2 is 0 if female and 1 if male, and 0.603 is the prevalence of males in the study group.

Candidate Predictors of the Functional Score—A list of potential correlates of the functional score was generated by discussion after review of available medical history data. In Table 1, we show the complete list of potential variables. Due to their low prevalence in our sample, we did not use an anatomic diagnosis of totally anomalous pulmonary venous connection, found in only 7 subjects, procedures performed on the pulmonary arteries after completion of the circuit, undertaken in only 4 subjects, and repair of the aortic arch during completion of the circuit, undertaken in only 1 subject. Medications were not considered as correlates due to wide variation in institutional practices.

Variable reduction—Cluster analysis was used to explore collinearity among variables, and to limit the number of variables considered for building our model.

Imputation of potential predictors—Some variables had missing values that were replaced prior to building the model using mean imputation from the modeling cohort. These were ventricular end-diastolic pressure, in 21 subjects, saturations of oxygen measured at preoperative catheterization, in 8 subjects, the income of the caregiver, in 49 subjects, and age at unloading surgery, in 5 subjects. Missing values in length of hospital stay, discovered in 12 subjects, were replaced by the median number of hospital days. Of note, although data relating to income was missing for 12% of the subjects, when a complete-case analysis was conducted, excluding the cases with unknown income, the multivariate model contained the same variables.

Development of the model—Linear regression was used to explore the relationship between the calculated functional score and potential correlates, and generalized additive modeling and graphical displays were used to determine whether any associations were non-linear. The number of years since completion of the circulation demonstrated a non-linear relationship with the score and was therefore categorized as less than 6 years, from 6 to 11 years, and greater than 11 years since completion, based on lower and upper quartiles. The number of operative procedures performed after completion of the circuit also demonstrated a non-linear relationship, and was categorized as up to 2, or more than 2, procedures.

Multivariate linear regression was used to identify associations between the calculated score and the potential correlates. The criterion for entry to the model was set at p equals 0.2. The criterion to remain in the model was p equals 0.05. Stepwise regression was first used to select a multivariate model that contained main effects only. Time since completion of the

Fontan circuit was retained in all multivariate modelling, as the value of some potential correlates depended on follow-up time. Bootstrap bagging was used to indicate reliability of variable selection for inclusion in the final multivariable model. In order to assess whether the predictors identified are important across clinical subgroups, we explored the effects of interactions. We examined 3 different interactions with each candidate predictor in separate models. One interaction term was for the presence of a prior superior cavopulmonary anastomosis, noting yes or no, one interaction term was for the age at Fontan completion of less than 3 years as opposed to equal to or more than 3 years, and the final interaction term was for left versus right versus mixed morphology for the dominant ventricle.

Validation of the model—We selected randomly 71 subjects, accounting for 15% of the overall study sample, to serve as a validation cohort. The data of these 71 subjects were not used in the construction of the primary model. The objective was to compare the predicted score from the regression equation with the score that was calculated from the database.

Results

Calculation of the Functional Score

Data for all components of the score were available for 285 subjects. Analyses indicated that the calculated score was robust when using an average of 3 variables if 4 were not available. Using 3 variables for definition resulted in scores that were within one standard deviation of the score based on complete data for 98% of subjects. On this basis, we included 476 subjects, in whom we had at least 3 components of the score. Similar results were achieved using a score restricted to those with all four components.

Among the 476 subjects, 396 (83%) had an ejection fraction available for analysis, 392 (82%) had percentage predicted maximal consumption of oxygen, 463 (97%) had a physical summary score, and all had a measured level of brain natriuretic peptide available. In Figure 1, we show the distribution of each component of the score, and the overall calculated functional score.

Spearman correlation analysis demonstrated that the four components of the score correlated weakly with each other, confirming that their contributions to a composite measure of functioning are largely independent. Only the physical summary score and the percentage predicted maximal consumption of oxygen had a correlation coefficient exceeding 0.15 (R equals 0.24). The calculated functional score itself had a symmetric distribution, with a mean of 49.5 plus or minus 17.3, and a range from 3 to 87.

Results of the Functional Score

In Table 1, we show the characteristics for the entire cohort of 476 subjects, the model cohort of 405 subjects, and the validation cohort of 71 subjects. For the modelling cohort, mean plus or minus standard deviation age at the Fontan procedure was 3.4 plus or minus 2.0 years and, at enrollment, the time since the procedure was 8.8 plus or minus 3.4 years. Of the subjects, 60% were male, and 79% were white. The dominant ventricle was morphologically left in 52% of the subjects, morphologically right in 33%, and was deemed to be of mixed morphology in the remaining 15%. The cohorts used for modeling and validation are similar with respect to all but one variable. The percentage of subjects with a surgical fenestration is lower in the cohort used for analysis than in the one used for validation, at 64.7% versus 80.3% (p equals 0.009).

Results of modelling

Reduction of variables—Cluster analysis revealed expected correlations such as right ventricular morphology and an anatomic diagnosis of hypoplastic left heart syndrome, cardiac anatomic diagnosis and age at Fontan, and pulmonary arterial pressure and end diastolic pressure. Of these, we retained ventricular morphology, age at Fontan, and end diastolic pressure. Postoperative complications were not used in modeling, as this variable was highly collinear with length of hospital stay. Similarly, since annual household income and highest level of education for the caregiver were collinear, we excluded education.

The results of univariate analysis of the candidate predictors using the modelling subset of 405 subjects are listed in Table 2. We identified 6 independent predictors of the functional score as determined by multivariate regression modelling (Table 3). We regarded the time since the procedure as an adjustment variable. The functional score was associated with ventricular morphology (p less than 0.001), with a lower score for those with a dominant right as compared to dominant left (p less than 0.001) or mixed ventricular types (p equals 0.002). The score did not differ between those having dominant left or mixed ventricular morphology (p equals 0.55). A lower score was also associated with higher ventricular end-diastolic pressure (p equals 0.003), and lower saturations of oxygen (p equals 0.047) at the time of the preoperative catheterization, as well as with lower annual family income (p equals 0.003). The multivariate model explained 18 percent of the variation in the calculated score. The multivariate model also included an interaction between performance of a prior superior cavopulmonary anastomosis and the presence of out of hospital arrhythmias after the procedure (p equals 0.003). Among subjects without a superior cavopulmonary anastomosis, the score was significantly lower, with a mean difference of 16.7 points, for those with compared to those without arrhythmias (p less than 0.001). Among subjects with a superior cavopulmonary anastomosis, the mean scores were similar, with a difference of 3.4 points, when those with were compared to those without arrhythmias (p equals 0.153; see Table 3). Thus, the presence of a postoperative arrhythmia was associated with a lower score if a prior superior cavopulmonary anastomosis had not been performed.

No association with the score was found for the centre where the Fontan circulation was created, gender, race, age at the time of the Fontan procedure, weight for age z-score at the time of the procedure, atrioventricular valvar regurgitation observed prior to the procedure, fenestration of the Fontan baffle, time of cardiopulmonary bypass, length of stay in hospital at the time of the Fontan procedure, and the number of operative surgeries or interventions undertaken subsequent to creation of the functionally univentricular circulation.

Validation of the model

The mean score for the cohort of 71 subjects used for validation was 50.0 plus or minus 14.9, with an interquartile range of 40.5 to 59.8, and a range from 12.0 to 81.3. There was no significant difference between the score predicted from the regression equation and the calculated score, the mean difference being 0.9 plus or minus 15.4 (p equals 0.625). In Figure 2, we show a scatter plot for the predicted score versus the original calculated score, revealing a weak correlation (R equals 0.17, p equals 0.16). The scatter plot demonstrates that the derived model predicts the outcome of 'middle' patients reasonably well, but for those with a higher functional state in particular, the model provides an underestimate. There are two extreme observations where the absolute difference between the predicted and the original calculated scores is greater than 40. After removing the two outliers (open circles), the correlation between the predicted and the original score in the remaining 69 subjects increases to 0.29 (p equals 0.017).

Discussion

Despite improvements in surgical outcomes following completion of the Fontan circulation, functional state varies considerably among survivors. Through creation of a functional score, averaging the percentile ranks of four main evaluations collected during our Fontan Cross-Sectional Study, we aimed to identify pre- and postoperative characteristics associated with overall function for patients in the years following their Fontan operation. We chose our 4 specific components because each was felt to represent a unique and important measure of overall functional state of the individual with Fontan physiology. After adjustment for time since the Fontan procedure, multivariate linear regression demonstrated that a lower functional score, indicating a worse functional state, was associated with: right ventricular morphology, higher ventricular end-diastolic pressure, lower saturations of oxygen, lower annual family income, and, in subjects without a superior cavopulmonary anastomosis, occurrence of arrhythmias after the Fontan procedure. The model explained almost one-fifth of the variation in the calculated score.

Previous series have evaluated risk factors for poor outcomes following palliation by creation of the Fontan circulation. In a recent report of 406 patients at a mean follow up of 6.1 plus or minus 5.7 years, factors associated with poor late outcome included qualitatively decreased ventricular function and elevated pulmonary arterial pressures prior to completion of the Fontan circuit.⁹ Cluster analysis of our data demonstrated a strong correlation between pulmonary arterial and ventricular end-diastolic pressures prior to the Fontan procedure, and therefore we removed pulmonary arterial pressure as a variable prior to building our model. In a cohort of 220 patients studied 10.2 plus or minus 0.6 years after a Fontan procedure, early failure of the circulation was associated with right ventricular morphology, preoperative elevated pulmonary vascular resistance and pulmonary arterial distortion, and postoperative elevated right atrial pressures.¹⁰ Late death was associated with a history of repair of coarctation and right ventricular morphology. Our data also suggests that worse functional state is associated with right ventricular morphology. Patients with a dominant right ventricle may fare worse due to an increased likelihood of poor coronary arterial perfusion during the initial stage of surgery, as in patients with hypoplastic left heart syndrome and aortic atresia. Differences intrinsic to the molecular biology of right ventricular myocardium or to anatomic geometry may also contribute to this finding. Another group reported that atrioventricular valvar regurgitation and heterotaxy were associated with late failure among 121 patients at a mean follow up time of 10.9 plus or minus 5.2 years, albeit that only heterotaxy remained significant in a multivariate equation.¹¹ In our study, the preoperative echocardiographic finding of moderate to severe atrioventricular valvar regurgitation in 20 subjects was not significantly associated using univariate analysis with the functional score (p equals 0.360). We had 33 subjects with heterotaxy syndrome in our cohort, albeit that cluster analysis revealed high collinearity with age at the time of the Fontan procedure and the presence of heterotaxy syndrome. Because of this we eliminated heterotaxy syndrome from the list of potential predictors.

Increased ventricular end-diastolic pressure is presumed to be detrimental to patients with the Fontan circulation due to the impeding of pulmonary venous inflow, the increased pulmonary pressures, and hence compromise of the passive flow dynamics of the Fontan circuit. In addition, elevated ventricular end-diastolic pressure can lead to atrial enlargement, which increases the risk for tachyarrhythmias, another reported risk factor for failure of the Fontan circuit. Reasons for the worse functional state of those with lower saturations of oxygen prior to creating the circuit are unclear. This was the weakest of the independent predictors identified. Chronic hypoxaemia may be a marker for decreased pulmonary arterial flow, and poorer pulmonary arterial growth, leading to less favourable outcomes following completion of the Fontan circuit.

The negative impact of postoperative arrhythmias on the patient with a Fontan circuit, both from a functional and a psychosocial perspective, has been well described. The decreased ventricular filling time associated with tachyarrhythmias rapidly leads to impaired ventricular relaxation, increased ventricular end-diastolic pressures, and potentially decreased flow through the Fontan circuit. Interventions such as cardioversion, and medications to control the arrhythmias, may negatively impact on the quality of life, and increase the risk of adverse effects. Our finding that arrhythmias were predictive of the functional score only in the absence of a superior cavopulmonary anastomosis is curious. The effect could not be explained by any single component of the score. Of note, most patients who did not undergo a prior superior cavopulmonary anastomosis had a history of tachyarrhythmia rather than bradyarrhythmia. Within the group of those with a superior cavopulmonary anastomosis, the prevalences of tachy-versus bradyarrhythmia were similar, consistent with the reported association between staged surgery and sinus nodal dysfunction.¹² Although highly statistically significant, we cannot exclude the possibility that the effect of this interaction is spurious. Independent replication of this interaction is of interest, as earlier studies may not have examined interactions by clinical subgroups.

Family income, another independent predictor of the functional score, was associated with the level of education of the parents. This association between the level of education and scores for quality of life has been reported previously.¹³

Limitations

Requirements for entry into the cross sectional study, including survival to the time of the study, ability to attempt an exercise test, and surgery performed at one of the 7 participating centres, necessarily resulted in bias of selection towards higher functioning patients. Brain natriuretic peptide was normalized using least-square residuals derived from the study sample, limiting the ability to apply this equation to all patients with the Fontan circuit. In general, as the score is derived from the percentile ranks of the population studied, caution should be used if seeking to extrapolate the score to other patients with the Fontan circuit. Assessment of cardiac function using calculations of ejection fraction derived from those with normal left ventricles may also introduce error. Despite utilization of all available data in this well characterized cohort, we were able to explain 18% of the variation noted in functional state following the completion of the Fontan circulation. Hence, over four-fifths of the variation remains unexplained. Unmeasured factors, such as subclinical myocarditis or subtle intraoperative events, may contribute significantly to outcomes. Alternatively, this cohort may be too highly functioning and homogeneous with respect to clinical state to allow us to identify predictors of good and poor function following completion of the Fontan circuit. Future studies including patients with failed Fontan physiology, such as those who have died or have undergone cardiac transplantation, are needed to investigate predictors of long-term outcomes following completion of the Fontan circulation.

Acknowledgments

Supported by U01 grants from the National Heart, Lung, and Blood Institute (HL068269, HL068270, HL068279, HL068281, HL068285, HL068292, HL068290, HL068288). Dr. Williams reports support from Grant Number KL2 RR024157 from the National Centre for Research Resources, a component of the National Institutes of Health, and National Institutes of Health Roadmap for Medical Research. Its contents are solely the responsibility of the authors and do not necessarily represent the official view of National Centre for Research Resources or National Institutes of Health. Information on National Centre for Research Resources is available at <http://www.ncrr.nih.gov/>. Information on Re-engineering the Clinical Research Enterprise can be obtained from <http://nihroadmap.nih.gov/clinicalresearch/overview-translational.asp>.

References

1. Jayakumar KA, Addonizio LJ, Kichuk-Chrisant MR, et al. Cardiac transplantation after the Fontan or Glenn procedure. *J Am Coll Cardiol* 2004;44:2065–2072. [PubMed: 15542293]
2. McCrindle BW, Williams RV, Mitchell PD, et al. Relationship of patient and medical characteristics to health status in children and adolescents after the Fontan procedure. *Circulation* 2006;113:1123–1129. [PubMed: 16490823]
3. Sleeper LA, Anderson P, Hsu DT, et al. Design of a large cross-sectional study to facilitate future clinical trials in children with the Fontan palliation. *Am Heart J* 2006;152:427–433. [PubMed: 16923408]
4. Anderson PA, Sleeper LA, Mahony L, et al. Contemporary Outcomes after the Fontan Procedure: A Pediatric Heart Network Multicentre Study. *J Am Coll Cardiol* 2008;52:85–98. [PubMed: 18598886]
5. Cooper DM, Weiler-Ravell D. Gas exchange response to exercise in children. *Am Rev Respir Dis* 1984;129:S47–S48. [PubMed: 6696341]
6. Paridon SM, Mitchell PD, Colan SD, et al. A cross-sectional study of exercise performance during the first 2 decades of life after the Fontan operation. *J Am Coll Cardiol* 2008;52:99–107. [PubMed: 18598887]
7. Landgraf JM. Measuring health-related quality of life in pediatric oncology patients: a brief commentary on the state of the art of measurement and application. *Int J Cancer Suppl* 1999;12:147–150. [PubMed: 10679887]
8. Shimizu H, Aono K, Masuta K, Asada H, Misaki A, Teraoka H. Stability of brain natriuretic peptide (BNP) in human blood samples. *Clin Chim Acta* 1999;285:169–172. [PubMed: 10481933]
9. Hosein RB, Clarke AJ, McGuirk SP, et al. Factors influencing early and late outcome following the Fontan procedure in the current era. The ‘Two Commandments’? *Eur J Cardiothorac Surg* 2007;31:344–352. [PubMed: 17236782]
10. Stamm C, Friehs I, Mayer JE Jr, et al. Long-term results of the lateral tunnel Fontan operation. *J Thorac Cardiovasc Surg* 2001;121:28–41. [PubMed: 11135157]
11. Ono M, Boethig D, Goerler H, Lange M, Westhoff-Bleck M, Breyman T. Clinical outcome of patients 20 years after Fontan operation – effect of fenestration on late morbidity. *Eur J Cardiothorac Surg* 2006;30:923–929. [PubMed: 17074498]
12. Manning PB, Mayer JE Jr, Wernovsky G, Fishberger SB, Walsh EP. Staged operation to Fontan increases the incidence of sinoatrial node dysfunction. *J Thorac Cardiovasc Surg* 1996;111:833–839. [PubMed: 8614144]
13. Saliba Z, Butera G, Bonnet D, et al. Quality of life and perceived health status in surviving adults with univentricular heart. *Heart* 2001;86:69–73. [PubMed: 11410565]

Appendix

National Heart, Lung, and Blood Institute: Gail Pearson, Mario Stylianou, Judith Massicot-Fisher, Marsha Mathis, Victoria Pemberton

Data Coordinating Centre: *New England Research Institutes*, Lynn Sleeper, Steven Colan, Paul Mitchell, Dianne Gallagher, Patti Nash, Gloria Klein, Minmin Lu

Network Chair: Lynn Mahony, *University of Texas Southwestern Medical Centre*

Clinical Site Investigators: *Children’s Hospital Boston*, Jane Newburger (PI), Stephen Roth, Roger Breitbart, Jonathan Rhodes, Jodi Elder, Ellen McGrath; *Children’s Hospital of New York*, Welton M. Gersony (PI), Seema Mital, Beth Printz, Ashwin Prakash, Darlene Servedio; *Children’s Hospital of Philadelphia*, Victoria Vetter (PI), Bernard J. Clark, Mark Fogel, Steven Paridon, Jack Rychik, Margaret Harkins, Jamie Koh; *Duke University*, the late Page A. W. Anderson (PI), Rene Herlong, Lynne Hurwitz, Jennifer S. Li, Ann Marie Nawrocki; *Medical University of South Carolina*, J. Philip Saul (PI), Andrew M. Atz,

Andrew D. Blafox, Girish Shirali, Jon Lucas, Amy Blevins; *Primary Children's Medical Centre, Salt Lake City, Utah*, LuAnn Minich (PI), Richard Williams, Linda Lambert, Michael Puchalski; *Hospital for Sick Children, Toronto*, Brian McCrindle (PI), Timothy Bradley, Kevin Roman, Jennifer Russell, Shi-Joon Yoo, Elizabeth Radojewski, Nancy Slater

Core Laboratories:

Cardiac MRI, Children's Hospital Boston: Tal Geva (Director); Andrew J. Powell

Echocardiography, Children's Hospital Boston: Steven Colan (Director), Marcy Schwartz, Renee Margossian

Protocol Review Committee: Michael Artman, Chair; Dana Connolly, Timothy Feltes, Julie Johnson, Jeffrey Krischer, G. Paul Matherne

Data and Safety Monitoring Board: John Kugler, Chair; Kathryn Davis, David J. Driscoll, Mark Galantowicz, Sally A. Hunsberger, Thomas J. Knight, Catherine L. Webb, Lawrence Wissow




Figure 1. Distribution of the functional score and its components. The middle line in each box indicates median, asterisk indicates the mean, and lower and upper edges of boxes represent the 25th and 75th percentiles, respectively. Three extreme values of Brain Natriuretic Peptide (377, 500, 652) are not shown.




Figure 2. Validation cohort predicted versus empirical functional score. Solid line is line of identity. Pearson correlation is 0.17, p equals 0.16. Outliers are depicted by open circles.

Table 1

Demographic, anatomic, and surgical characteristics of the cohorts. All variables listed were considered in modelling as potential correlates of Fontan Functional Score, unless otherwise noted in Methods.

Variables	Study Cohort		Modelling Cohort		Validation Cohort		p value*
	Number missing	Mean \pm SD, Median (Interquartile Range) or %	Mean \pm SD, Median (Interquartile Range) or %	Mean \pm SD, Median (Interquartile Range) or %	Mean \pm SD, Median (Interquartile Range) or %		
Number of subjects		476	405	71			
Fontan Functional Score		49.5 \pm 17.3	49.4 \pm 17.7	50.0 \pm 14.9			0.803
Age at enrollment, year		12.0 \pm 3.4	12.0 \pm 3.4	12.1 \pm 3.3			0.825
Race	1						0.133
White		382 (80.4%)	319 (79.0%)	63 (88.7%)			
Black		47 (9.9%)	44 (10.9%)	3 (4.2%)			
Other		46 (9.7%)	41 (10.1%)	5 (7.0%)			
Male		287 (60.3%)	244 (60.2%)	43 (60.6%)			1.000
Parental highest grade of school	6						0.391
Some high school or less		29 (6.2%)	22 (5.5%)	7 (10.3%)			
High school graduate or GED		117 (24.9%)	105 (26.1%)	12 (17.6%)			
Vocational school, some college		159 (33.8%)	136 (33.8%)	23 (33.8%)			
4 year college graduate		110 (23.4%)	93 (23.1%)	17 (25.0%)			
Graduate degree		55 (11.7%)	46 (11.4%)	9 (13.2%)			
Household income	61						0.174
<\$20 000		52 (12.5%)	46 (12.9%)	6 (10.2%)			
\$20 000–39 000		76 (18.3%)	70 (19.7%)	6 (10.2%)			
\$40 000–59 000		69 (16.6%)	53 (14.9%)	16 (27.1%)			
\$60 000–79 000		63 (15.2%)	54 (15.2%)	9 (15.3%)			
\$80 000–99 000		58 (14.0%)	51 (14.3%)	7 (11.9%)			
>\$100 000		97 (23.4%)	82 (23.0%)	15 (25.4%)			
Pulmonary atresia		29 (6.1%)	25 (6.2%)	4 (5.6%)			1.000
Hypoplastic left heart syndrome		100 (21.0%)	83 (20.5%)	17 (23.9%)			0.529
Common atrioventricular valve/Heterotaxy/Unbalanced atrioventricular septal defect		52 (10.9%)	42 (10.4%)	10 (14.1%)			0.355
Tricuspid atresia		114 (24.0%)	97 (24.0%)	17 (23.9%)			1.000

Variables	Study Cohort		Modelling Cohort		Validation Cohort		p value*
	Number missing	Mean ± SD, Median (Interquartile Range) or %	Mean ± SD, Median (Interquartile Range) or %	Mean ± SD, Median (Interquartile Range) or %	Mean ± SD, Median (Interquartile Range) or %		
Totally anomalous pulmonary venous return		8 (1.7%)	5 (1.2%)	3 (4.2%)	0.103		
Ventricular type					0.671		
Left		244 (51.3%)	211 (52.1%)	33 (46.5%)			
Right		162 (34.0%)	135 (33.3%)	27 (38.0%)			
Mixed		70 (14.7%)	59 (14.6%)	11 (15.5%)			
Pre-Fontan mean pulmonary arterial pressure, mmHg	43	11.4 ± 3.3	11.4 ± 3.3	11.9 ± 3.3	0.200		
Pre-Fontan end-diastolic pressure, mmHg	22	7.8 ± 3.3	7.8 ± 3.4	8.0 ± 2.6	0.562		
Pre-Fontan oxygen saturation, percent	13	84 ± 5.2	84.0 ± 5.0	84.3 ± 6.4	0.694		
Age at volume unloading surgery (either superior cavopulmonary anastomosis or Fontan), year	6	1.0 (0.6, 2.3)	1.0 (0.6, 2.3)	1.1 (0.6, 2.2)	0.949		
Superior cavopulmonary anastomosis		351 (73.7%)	297 (73.3%)	54 (76.1%)	0.770		
Number of pulmonary arterial surgeries		0.4 ± 0.7	0.4 ± 0.7	0.5 ± 0.7	0.819		
Moderate/Severe atrioventricular valvar regurgitation	48	20 (4.7%)	17 (4.7%)	3 (4.4%)	1.000		
Moderate/Severe ventricular dysfunction	31	6 (1.4%)	5 (1.4%)	1 (1.5%)	1.000		
Age at Fontan, year		2.9 (2.1, 4.0)	2.9 (2.1, 3.9)	3.0 (1.9, 5.0)	0.657		
Weight for age z score at Fontan	8	-1.0 ± 1.2	-1.0 ± 1.2	-0.9 ± 1.2	0.457		
Fenestration		319 (67.0%)	262 (64.7%)	57 (80.3%)	0.009		
Cardiopulmonary bypass time, minutes	37	113.9 ± 47.4	114.0 ± 48.1	113.0 ± 43.4	0.865		
Type of Fontan					0.346		
Atriopulmonary connection		67 (14.1%)	61 (15.1%)	6 (8.5%)			
Total cavopulmonary connection intracardiac lateral tunnel		287 (60.3%)	244 (60.2%)	43 (60.6%)			
Total cavopulmonary connection extracardiac lateral tunnel		53 (11.1%)	41 (10.1%)	12 (16.9%)			
Total cavopulmonary connection extracardiac conduit		60 (12.6%)	51 (12.6%)	9 (12.7%)			
Other		9 (1.9%)	8 (2.0%)	1 (1.4%)			
Post-operation complication: Prolonged pleural/pericardial effusions/chylothorax		177 (37.6%)	149 (37.3%)	28 (39.4%)	0.791		
Length of hospital stay, days	9	11 (9, 18)	12 (9, 18)	11 (8, 18)	0.454		
Post-Fontan arrhythmia		101 (21.2%)	86 (21.2%)	15 (21.1%)	1.000		
Post-Fontan surgeries ≥ 2		13 (2.7%)	12 (3.0%)	1 (1.4%)	0.702		
Number of post-Fontan catheterization interventions		0.8 ± 1.1	0.7 ± 1.1	1.0 ± 1.2	0.107		

Variables	Study Cohort		Modelling Cohort		Validation Cohort		p value*
	Number missing	Mean ± SD, Median (Interquartile Range) or %	Mean ± SD, Median (Interquartile Range) or %	Mean ± SD, Median (Interquartile Range) or %	Mean ± SD, Median (Interquartile Range) or %		
Aortic arch repair		1 (0.2%)	0 (0.0%)	1 (1.4%)		0.149	
Pulmonary arterial surgeries		4 (0.8%)	4 (0.9%)	0 (0.0%)		1.000	
Number of years since Fontan		8.7 ± 3.4	8.8 ± 3.4	8.4 ± 3.7		0.398	
Currently on pacemaker		64 (13.5%)	51 (12.6%)	13 (18.3%)		0.191	

* Test of significance for differences between model sample (405 subjects) and validation sample (71 subjects).

Table 2

Unadjusted linear regression of Fontan Functional Score on each independent variable (405 subjects).

Variables	R ²	Estimate	Mean	p
Demographics				
Race	0.008			0.216
White		1.04	50.1	
Black		-3.93	45.2	
Other		-	49.1	
Male	0.001	1.03		0.566
Parental highest grade of school				
Some high school or less	0.019	-4.25	47.1	0.109
High school graduate/GED		1.04	52.4	
Vocational school, some college		-4.77	46.6	
4 year college graduate		-1.08	50.3	
Graduate degree		-	51.4	
Annual household income, \$20K increments	0.027	1.76		0.001
Ventricular type				
Right	0.022	-	45.9	0.011
Left		4.79	50.7	
Mixed		7.19	53.1	
Pre-Fontan end-diastolic pressure, mmHg	0.041	-1.08		< 0.001
Pre-Fontan oxygen saturation, %	0.019	0.50		0.005
Superior cavopulmonary anastomosis:				
Yes	0.020	5.68	50.9	0.004
No			45.2	
Age at unloading surgery	0.022	-1.68		0.003
Moderate/Severe atrioventricular valvar regurgitation:				
Yes	0.003	-1.85	45.8	0.502
No		2.22	49.8	
Unknown		-	47.6	
Number of pulmonary arterial surgeries	0.001	-0.74		0.546
Age at Fontan, year	0.005	-0.62		0.157
Weight for age z score at Fontan	0.001	0.55		0.446
Surgical fenestration:				
Yes	0.004	2.35	50.2	0.202
No		-	47.9	
Type of Fontan				
Atriopulmonary connection	0.032	-	46.6	0.011
Total cavopulmonary connection intracardiac lateral tunnel		2.25	48.9	
Total cavopulmonary connection extracardiac lateral tunnel		6.59	53.2	
Total cavopulmonary connection extracardiac conduit		7.86	54.5	
Other		-11.3	35.3	

Variables	R ²	Estimate	Mean	p
Cardiopulmonary bypass time, minutes	0.002	-0.02		0.347
Length of hospital stay, days	0.009	-0.09		0.062
Log transformed length of stay	0.018	-3.62		0.007
Arrhythmias:				
Yes	0.034	-7.93	43.2	< 0.001
No		-	51.1	
Currently on pacemaker:				
Yes	0.021	-7.73	42.7	0.004
No		-	50.4	
Post-Fontan surgeries ≥2				0.004
Yes	0.020	-14.9	34.9	
No		-	49.9	
Number of post-Fontan catheterization interventions	0.011	-1.68		0.033
Number of years since Fontan	0.013	-0.59		0.023
Number of years since Fontan categorized	0.038			< 0.001
Less than 6 years		4.91	49.3	
6 to 11 years		8.16	52.5	
Greater than or equal to 11 years		-	44.4	

Table 3Independent predictors of Fontan Functional Score (Model $R^2 = 0.18$).

Variable	Estimate \pm Standard Error	Least-Squares Mean	p-value
Years since Fontan			0.042
Less than 6	2.15 \pm 2.54	45.3	
6 to 11	5.19 \pm 2.19	48.3	
Greater than or equal to 11	–	43.1	
Ventricular type			<0.001
Left	–1.45 \pm 2.42	47.2	
Right	–7.88 \pm 2.57	40.8	
Mixed	–	48.7	
Annual household income, \$20 K increments	1.53 \pm 0.50		0.003
Pre-Fontan end diastolic pressure, mmHg	–0.75 \pm 0.25		0.003
Pre-Fontan oxygen saturation	0.34 \pm 0.17		0.047
Post Fontan arrhythmia*			<0.001
Superior Cavopulmonary anastomosis			0.004
Superior cavopulmonary anastomosis \times post-Fontan arrhythmia interaction			0.003
No superior cavopulmonary anastomosis			
No arrhythmia		50.3	<0.001
Arrhythmia		33.6	
Superior cavopulmonary anastomosis performed			
No arrhythmia		50.9	0.153
Arrhythmia		47.5	

* Main effects estimates for post-Fontan arrhythmia and superior cavopulmonary anastomosis are not interpretable and not shown due to the presence of a significant interaction.