

Children's Mercy Kansas City

SHARE @ Children's Mercy

Manuscripts, Articles, Book Chapters and Other Papers

11-1-2016

The Influence of Age on the Diagnostic Performance of White Blood Cell Count and Absolute Neutrophil Count in Suspected Pediatric Appendicitis.

Richard G. Bachur

Peter S. Dayan

Nanette C. Dudley

Lalit Bajaj

Michelle D. Stevenson

See next page for additional authors

Follow this and additional works at: <https://scholarlyexchange.childrensmercy.org/papers>



Part of the [Diagnosis Commons](#), [Emergency Medicine Commons](#), [Medical Cell Biology Commons](#), and the [Pediatrics Commons](#)

Recommended Citation

Bachur RG, Dayan PS, Dudley NC, et al. The Influence of Age on the Diagnostic Performance of White Blood Cell Count and Absolute Neutrophil Count in Suspected Pediatric Appendicitis. *Acad Emerg Med.* 2016;23(11):1235-1242. doi:10.1111/acem.13018

This Article is brought to you for free and open access by SHARE @ Children's Mercy. It has been accepted for inclusion in Manuscripts, Articles, Book Chapters and Other Papers by an authorized administrator of SHARE @ Children's Mercy. For more information, please contact hlsteel@cmh.edu.

Creator(s)

Richard G. Bachur, Peter S. Dayan, Nanette C. Dudley, Lalit Bajaj, Michelle D. Stevenson, Charles G. Macias, Manoj K. Mittal, Jonathan Bennett, Kelly A. Sinclair, Michael C. Monuteaux, and Anupam B. Kharbanda

ORIGINAL CONTRIBUTION

The Influence of Age on the Diagnostic Performance of White Blood Cell Count and Absolute Neutrophil Count in Suspected Pediatric Appendicitis

Richard G. Bachur, MD, Peter S. Dayan, MD, MSc, Nanette C. Dudley, MD, Lalit Bajaj, MD, MPH, Michelle D. Stevenson, MD, MS, Charles G. Macias, MD, MPH, CGM, Manoj K. Mittal, MD, Jonathan Bennett, MD, Kelly Sinclair, MD, Michael C. Monuteaux, ScD, and Anupam B. Kharbanda, MD, MSc

Abstract

Objective: White blood cell (WBC) count and absolute neutrophil count (ANC) are a standard part of the evaluation of suspected appendicitis. Specific threshold values are utilized in clinical pathways, but the discriminatory value of WBC count and ANC may vary by age. The objective of this study was to investigate whether the diagnostic value of WBC count and ANC varies across age groups and whether diagnostic thresholds should be age-adjusted.

Methods: This is a multicenter prospective observational study of patients aged 3–18 years who were evaluated for appendicitis. Receiver operator characteristic curves were developed to assess overall discriminative power of WBC count and ANC across three age groups: <5, 5–11, and 12–18 years of age. Diagnostic performance of WBC count and ANC was then assessed at specific cut-points.

Results: A total of 2,133 patients with a median age of 10.9 years (interquartile range = 8.0–13.9 years) were studied. Forty-one percent had appendicitis. The area under the curve (AUC) for WBC count was 0.69 (95% confidence interval [CI] = 0.61 to 0.77) for patients < 5 years of age, 0.76 (95% CI = 0.73 to 0.79) for 5–11 years of age, and 0.83 (95% CI = 0.81 to 0.86) for 12–18 years of age. The AUCs for ANC across age groups mirrored WBC performance. At a commonly utilized WBC cut-point of 10,000/mm³, the sensitivity decreased with increasing age: 95% (<5 years), 91% (5–11 years), and 89% (12–18 years) whereas specificity increased by age: 36% (<5 years), 49% (5–12 years), and 64% (12–18 years).

Conclusion: WBC count and ANC had better diagnostic performance with increasing age. Age-adjusted values of WBC count or ANC should be considered in diagnostic strategies for suspected pediatric appendicitis.

From the Division of Emergency Medicine, Children's Hospital Boston and Harvard Medical School (RGB, MCM), Boston, MA; the Department of Pediatrics, Columbia University College of Physicians and Surgeons (PSD), New York, NY; the Department of Pediatrics, University of Utah School of Medicine (NCD), Salt Lake City, UT; the Department of Pediatrics, University of Colorado School of Medicine (LB), Denver, CO; the Department of Pediatrics, University of Louisville (MDS), Louisville, KY; the Department of Pediatrics, Baylor College of Medicine (CGM), Houston, TX; the Department of Pediatrics, The Children's Hospital of Philadelphia and Perelman School of Medicine, University of Pennsylvania (MKM), Philadelphia, PA; the Department of Pediatrics, Alfred I. DuPont Hospital for Children (JB), Wilmington, DE; the Division of Emergency Medicine, Children's Mercy Hospitals and Clinics (KS), Kansas City, MO; and the Department of Pediatric Emergency Medicine, Children's Hospital and Clinics of Minnesota (ABK), Minneapolis, MN.

Received April 17, 2016; revision received May 22, 2016; accepted May 31, 2016.

The authors have no relevant financial information or potential conflicts to disclose.

Author contributions—ABK is the principal investigator for the original multicenter study proposal and had oversight over the parent study protocol, data collection, data security, and multicenter coordination; ABK approved the secondary analysis and provided critical review of this manuscript. PSD is the senior investigator for the original multicenter study and provided critical review of the manuscript. RGB conceived of this secondary study and was responsible for primary data analysis, drafting of manuscript, and final responsibility for the submitted manuscript. LB, JB, NCD, CGM, MKM, KS, and MDS contributed as site-PIs under the parent protocol and provided critical review of this manuscript. MCM provided statistical expertise, partially drafted the manuscript, and provided critical review. All authors approved final version of manuscript prior to submission.

Supervising Editor: Mark R. Zonfrillo, MD, MSCE.

Address for correspondence and reprints: Richard Bachur, MD; e-mail: Richard.bachur@childrens.harvard.edu.

The diagnostic evaluation of children and adolescents for possible appendicitis is common in outpatient settings such as primary care clinics and emergency departments (EDs). The presentation of appendicitis is known to vary by the duration of symptoms, the age of the patient, and specific location of the appendix. Prior to the past 20 years, the preoperative diagnosis was based on history and physical examination alone; recent advances to improve outcomes have included clinical decision algorithms and scores,^{1–9} advanced diagnostic imaging,^{10–15} and the pursuit of new biomarkers^{16–20} to help clinicians discriminate appendicitis from other causes of abdominal pain.

Nearly all clinical scores and diagnostic algorithms utilize the white blood cell (WBC) count and associated WBC differential (as the proportion of neutrophils or absolute neutrophil count [ANC]) to guide care.^{2,3,7,9,21} Previous investigations have shown the diagnostic importance of WBC and neutrophil predominance^{22–29} in the evaluation of possible appendicitis. Furthermore, prior studies have demonstrated that the WBC count and ANC have the best test performance of currently available laboratory tests to predict appendicitis.^{18,20} The two most utilized diagnostic scoring systems, the Alvarado Score³ and Pediatric Appendicitis Score,⁷ include an elevated WBC and neutrophil predominance as predictive factors. Previous investigations focused specifically on WBC have been limited by sample size,²⁸ investigated in a selective population of those undergoing surgery,^{30–34} or included broader abdominal pain populations²⁶ rather than focusing the investigation to those being considered for appendicitis. Although many current algorithms employ the WBC count and ANC (or neutrophil predominance) in decision-making, specific diagnostic thresholds are not adjusted based on age. In this study we aimed to determine whether the discriminative performance of the WBC count and ANC vary by age and if the diagnostic thresholds should be adjusted by age for guiding care in the evaluation of possible pediatric appendicitis.

METHODS

Study Design and Setting

We conducted a secondary analysis of a prospective multicenter observational study whose primary objective was to refine a clinical prediction rule for appendicitis.³⁵ The original study enrolled children with suspected appendicitis at nine pediatric EDs who were members of the Pediatric Emergency Medicine Clinical Research Committee of the American Academy of Pediatrics. Subjects were enrolled from March 2009 through April 2010. The study was approved by each site's institutional review board (IRB). Seven IRBs granted a waiver of written consent/assent and obtained verbal consent. At the two remaining sites, written consent was obtained from the guardians and assent from children 7 years of age and older.

Selection of Participants

ED patients who were 3 to 18 years of age presenting with acute abdominal pain and had suspected appendicitis were enrolled. Suspected appendicitis was defined by the clinical team's evaluation for appendicitis by obtaining blood tests, advanced radiologic studies, or surgical consultation. Patients with any of the following conditions were excluded: pregnancy, prior abdominal surgery, chronic gastrointestinal condition, or severe developmental delay (that might interfere with an accurate clinical assessment). For this secondary analysis, we included only those patients who had acute abdominal pain for less than 72 hours and had a WBC count and WBC differential obtained as part of the evaluation.

Data Collection

Site primary investigators received a manual of operations and were instructed on proper completion of case report forms. A pediatric emergency medicine physician completed a standardized case report form with regard to symptoms and signs prior to any advanced imaging or operative care. Research assistants entered data from the medical record, including all laboratory values, for electronic transfer to a central data center.

Outcome Measures

The primary outcome was the presence or absence of appendicitis. For those patients who had operative care, the presence of appendicitis was determined from the pathologist's final report. Presence or absence of perforation was determined from the attending surgeon's operative report. For patients discharged without surgery, telephone follow-up was performed between 1 and 2 weeks to determine resolution of signs and symptoms, visits to other sites of care, and need for any surgery. If we were unable to contact the guardian, research coordinators reviewed the medical record for the 90 days after the ED visit to determine if the patient underwent an operation at the index facility.

Data Analysis

Descriptive statistics of patients' demographic and clinical characteristics were calculated using frequencies and proportions for categorical variables and medians with interquartile ranges (IQRs) for continuous variables. For our primary analysis, we determined the diagnostic performance (sensitivity, specificity, positive predictive value [PPV], negative predictive value [NPV], and positive likelihood ratio) of WBC count and ANC (using 1000/mm³ increments from 5,000 to 15,000/mm³ as cut-points), stratified across three age groups: pre-school (<5 years), school age (5–11 years), and adolescents (12–18 years). Before conducting data analyses, the WBC count and ANC data were checked for the presence of outliers. To assess overall discriminative value, receiver operator characteristic curves (ROC) and the area under the curve (AUC) were determined and

compared between age groups using a nonparametric test as described by DeLong et al.³⁶ The presence of influential data points was determined using the Delta-Beta influence statistic, a standardized measure of the difference in the coefficient vector (in this case, either WBC count or ANC) that is due to deletion of observations with the identical covariate pattern.³⁷ For illustrative purposes, the test characteristics corresponding to the commonly used WBC cutoff of 10,000/mm³ were highlighted and presented.

We also considered the possibility that WBC count and ANC are affected by the duration of symptoms prior to the ED evaluation. To test if symptom duration contributed any predictive information to appendicitis status beyond that already provided by WBC count, we estimated a logistic regression model for each age subgroup with appendicitis as the outcome and WBC count as the independent variable. We then estimated a second logistic model with both WBC count and symptom duration as the independent variables. We then tested the null hypothesis that the AUCs from both models were equal. An identical assessment was made for ANC. All analyses were conducted using Stata 13.1.

RESULTS

Study Group

A total of 2,624 patients were enrolled in the parent study of which 2,133 (81%) patients had pain less than 72 hours and had a WBC count and WBC differential as part of their ED evaluation. Details of patient characteristics are presented in Table 1. The median age was 10.9 years (IQR = 8.0–13.9 years) and 48% were female. Forty-one percent of patients had appendicitis of which 219 (10%) were perforated. A total of 163 (8%) children were less than 5 years of age, 1,107 (52%) were

5–11 years of age, and 863 (40%) were 12–18 years of age. Of those patients not classified as appendicitis cases ($n = 1,236$), the most common diagnosis was non-specific abdominal pain, followed by gastroenteritis and constipation. This pattern was similar within each age group (Table 2).

Diagnostic Performance of WBC Count and ANC

ROC analysis for WBC count and ANC by age group is shown in Figure 1. The AUC for WBC count was significantly greater in the 12- to 18-year-old group (0.83 [95% CI = 0.81 to 0.86]) compared to both the <5-year-old group (0.69 [95% CI = 0.61 to 0.77]; $p = 0.001$) and the 5- to 11-year-old group (0.76 [95% CI = 0.73 to 0.79]; $p < 0.001$). No difference was found between the WBC count AUC values of the <5- and 5- to 11-year-old groups ($p = 0.11$). Similarly, the AUC for ANC was significantly greater in the 12- to 18-year-old group (0.83 [95% CI = 0.80 to 0.86]) compared to both the <5-year-old group (0.68 [95% CI = 0.60 to 0.77]; $p = 0.001$) and the 5- to 11-year-old group (0.76 [95% CI = 0.73 to 0.78]; $p < 0.001$). No difference was found between the ANC AUC values of the <5- and 5- to 11-year-old groups ($p = 0.11$).

Evaluation of the Delta-Beta statistics revealed five WBC data points as influential (i.e., large Delta-Beta values outside the range of the rest of the distribution). However, repeating the analysis without these observations did not change the magnitude of the AUC values, their CIs, or the pattern of significance for the pairwise comparisons. No influential data points were detected in the analysis of ANC.

The detail of a range of cut-points is shown in Table 3 (WBC count) and Table 4 (ANC). The PPV and NPV are displayed across a range of WBC cut-points in Figures 2 and 3, respectively. The PPV and NPV are similarly graphed across a range of ANC cut-points in Data

Table 1
Clinical Characteristics of Study Sample ($N = 2,133$)

	No Appendicitis, $n = 1,263$	Appendicitis, $n = 870$	Nonperforated Appendicitis, $n = 651$	Perforated Appendicitis, $n = 219$
Age (y)	11.0 (7.7–14.2)	10.9 (8.3–13.3)	11.3 (8.7–13.6)	9.6 (7.1–12.4)
Age subgroups (y)				
<5	125 (10)	38 (4)	21 (3)	17 (8)
5–11	609 (48)	498 (57)	363 (56)	135 (62)
>12	529 (42)	334 (38)	267 (41)	67 (31)
Female (%)	683 (54)	348 (40)	252 (39)	96 (44)
Duration of abdominal pain (h)				
<12	416 (33)	210 (17)	193 (30)	17 (8)
12–24	400 (32)	296 (34)	244 (37)	52 (24)
25–48	312 (25)	265 (30)	176 (27)	89 (41)
49–72	135 (11)	99 (11)	38 (6)	61 (28)
WBC count by age group ($\times 1,000/\text{mm}^3$)				
<5 y	13.5 \pm 6.7	17.1 \pm 4.9	17.4 \pm 3.1	16.7 \pm 6.6
5–11 y	11.5 \pm 5.7	16.3 \pm 5.0	15.6 \pm 4.5	18.1 \pm 5.7
>12 y	9.7 \pm 4.0	15.3 \pm 4.7	14.9 \pm 4.4	16.6 \pm 5.6
ANC by age group ($\times 1,000/\text{mm}^3$)				
<5 y	10.7 \pm 6.2	13.9 \pm 4.7	14.1 \pm 3.3	13.7 \pm 6.2
5–11 y	8.7 \pm 5.7	13.4 \pm 4.9	12.7 \pm 4.5	15.5 \pm 5.3
>12 y	6.8 \pm 4.2	12.5 \pm 4.8	12.1 \pm 4.5	14.1 \pm 5.6

Categorical data are presented as n (%); other data are reported as median (interquartile range) or mean \pm SD.

Table 2
Other diagnoses among patients without appendicitis, stratified by age group

Other Diagnoses	< 5 years (n = 122)	5-11 years (n = 597)	12-18 years (n = 517)	Total (n = 1,236)
Non-specific abdominal pain	42 (34)	256 (43)	222 (43)	520 (42)
Gastroenteritis	30 (25)	103 (17)	51 (10)	184 (15)
Other, not classified	21 (17)	86 (14)	69 (13)	176 (14)
Constipation	16 (13)	73 (12)	62 (12)	151 (12)
Ovarian cyst	0 (0)	2 (<1)	58 (11)	60 (5)
Mesenteric adenitis	3 (2)	24 (4)	15 (3)	42 (3)
Pneumonia	4 (3)	19 (3)	2 (<1)	25 (2)
Urinary tract infection	3 (2)	12 (2)	9 (2)	24 (2)
Pyelonephritis	2 (2)	9 (2)	5 (1)	16 (1)
Renal stone	0 (0)	4 (1)	10 (2)	14 (1)
Gastritis	1 (1)	9 (2)	3 (1)	13 (1)
Pelvic inflammatory disease	0 (0)	0 (0)	11 (2)	11 (1)

Values in table represent frequency (percent)




Figure 1. Receiver operator characteristic curves for WBC count and ANC across three age groups of children with suspected appendicitis. ANC = absolute neutrophil count; AUC = area under the curve; WBC = white blood cell.

Supplements S1 and S2 (available as supporting information in the online version of this paper). Within the three age subgroups, the AUCs of WBC count alone did not significantly differ from the AUCs of WBC count

combined with symptom duration in classifying appendicitis (p-values = 0.79, 0.12, and 0.74 for the <5, 5–11, and 12–18 age groups, respectively). The same pattern of results was found for the AUCs of ANC with and without symptom duration in classifying appendicitis.

Illustrative Example: WBC Count 10,000/mm³ Threshold

At a WBC count cut-point of 10,000/mm³, the sensitivity decreased with increasing age: <5 years, 95% (95% CI = 82% to 99%); 5–11 years, 91% (95% CI = 89% to 94%); and 12–18 years, 89% (95% CI = 85% to 92%). Specificity increased with increasing age (by group): 36% (95% CI = 28% to 45%), 49% (95% CI = 45% to 53%), and 64% (95% CI = 59% to 68%). The PPV increased with increasing age (by group) from 31% (95% CI = 23% to 40%) to 59% (95% CI = 56% to 63%) to 61% (95% CI = 56% to 65%), whereas the NPV was highest for children less than 5 years of age (96% [95% CI = 85% to 99%]) compared to the 5- to 11-year-old patients (87% [95% CI = 83% to 91%]) or those over 12 years of age [90% (95% CI 87 - 93%)].

DISCUSSION

Despite the near-universal use of WBC count and ANC as part of the diagnostic evaluation of suspected appendicitis in children, threshold values in published scores and decision algorithms have not been previously age-adjusted. We demonstrated that the predictive capability of WBC count and ANC varies by age. More importantly, at any specific cut-point, the predictive value varies significantly between age groups thereby arguing for age-adjusted WBC count or ANC thresholds. Furthermore, the diagnostic value of WBC count and ANC did not vary with duration of abdominal pain and is consistent with previous investigations;^{22,23,28,38} specific to our investigation, this lack of association with pain duration remained true among the age subgroups.

Despite the ubiquity of patient presentations with abdominal pain and a concern for appendicitis, the diagnostic evaluation for appendicitis remains challenging especially in children. The diagnostic value of

Table 3
Diagnostic Performance of WBC Count Across Different Cut-points by Age of Patient

WBC classification cut-point (×1000/mm ³)	Age Group (y)														
	<5 (n = 163)					5–11 (n = 1,107)					12–18 (n = 863)				
	Sens	Spec	PPV	NPV	+LR	Sens	Spec	PPV	NPV	+LR	Sens	Spec	PPV	NPV	+LR
≥5	100	5	24	100	1.05	99	7	46	88	1.06	100	5	40	100	1.05
≥6	100	10	25	100	1.12	99	14	48	93	1.15	99	13	42	97	1.15
≥7	100	14	26	100	1.16	98	23	51	92	1.27	98	27	46	95	1.33
≥8	100	20	28	100	1.25	97	33	54	92	1.43	97	41	51	95	1.64
≥9	97	28	29	97	1.35	95	41	57	91	1.6	94	52	55	93	1.96
≥10	95	36	31	96	1.48	91	49	59	87	1.78	89	64	61	90	2.46
≥11	92	42	32	95	1.58	87	58	63	85	2.06	81	71	64	86	2.81
≥12	89	50	35	94	1.78	82	63	64	81	2.19	73	77	67	82	3.15
≥13	84	56	37	92	1.91	76	67	66	77	2.33	62	82	68	77	3.4
≥14	76	60	37	89	1.91	67	72	66	72	2.35	55	87	73	76	4.37
≥15	68	63	36	87	1.86	60	76	67	70	2.5	48	91	77	73	5.25

Values for Sens, Spec, PPV, and NPV represent percentages;
+LR = positive likelihood ratio; NPV = negative predictive value; PPV = positive predictive value; Sens = sensitivity; Spec = specificity; WBC = white blood cell.

Table 4
Diagnostic Performance of ANC Across Different Cut-points by Age of Patient

ANC classification cut-point (×1000/mm ³)	Age Group (y)														
	<5 (n = 163)					5–11 (n = 1,107)					12–18 (n = 863)				
	Sens	Spec	PPV	NPV	+LR	Sens	Spec	PPV	NPV	+LR	Sens	Spec	PPV	NPV	+LR
≥5	100	20	28	100	1.25	97	33	54	94	1.45	96	43	52	95	1.7
≥6	97	27	29	97	1.34	95	40	57	91	1.6	93	56	57	93	2.11
≥7	95	30	29	95	1.36	92	47	59	88	1.75	89	64	61	90	2.45
≥8	92	38	31	94	1.48	88	55	61	85	1.94	83	70	63	87	2.72
≥9	89	45	33	93	1.62	82	60	63	80	2.04	75	76	66	83	3.12
≥10	87	54	36	93	1.87	77	66	65	78	2.27	64	81	68	78	3.41
≥11	74	59	35	88	1.81	70	70	66	74	2.33	57	85	70	76	3.72
≥12	63	62	34	85	1.68	61	75	67	70	2.44	51	89	74	74	4.49
≥13	55	70	36	84	1.87	52	79	67	67	2.49	45	91	76	72	5.02
≥14	50	73	36	83	1.84	41	83	66	63	2.36	37	93	77	70	5.37
≥15	34	75	30	79	1.38	32	87	66	61	2.42	29	95	79	68	5.85

Values for Sens, Spec, PPV and NPV represent percentages.
ANC = absolute neutrophil count; +LR = positive likelihood ratio; NPV = negative predictive value; PPV = positive predictive value; Sens = sensitivity; Spec = specificity.

WBC count and ANC has been previously studied in adults^{22,30,34,39,40} and children.^{16,23,26,28,31,33,34,38,40,41} Even with limitations in the sample size or biases in the study population, the markers have consistently shown discriminatory value for appendicitis and complicated appendicitis. High-quality cohort studies inclusive of pediatric patients have reported overall diagnostic value of WBC count and ANC to have AUCs of 0.78–0.92 and 0.78–0.89 respectively.^{28,39}

WBC count and ANC (or neutrophil predominance) are incorporated into nearly all diagnostic pathways for appendicitis. The two most referenced clinical scores are the Alvarado score³ and the Pediatric Appendicitis Score.⁷ Both scoring systems considered an elevated WBC count (>10,000/mm³) and neutrophil predominance (Alvarado score > 75% PMN; Pediatric Appendicitis Score includes “neutrophilia” without a definition) as predictors for appendicitis. Other

guidelines for suspected appendicitis have assigned similar cutoffs^{2,8,9} or performed an independent analysis of threshold values.^{21,29} The exact cut-points incorporated into guidelines depended on the balance between sensitivity and specificity and whether the goal of the guideline was to identify high-risk or low-risk patients. None of the prior published guidelines or scoring systems has provided age-adjusted values for WBC count or ANC.

Outside of clinical score and decision pathways, we found two prior relevant investigations that considered age-adjusted values of WBCs. In a study of 200 children undergoing surgery for appendicitis, the AUC using ROC analysis was 0.46 for those < 6 years of age compared to 0.79 for those 6 to 19 years of age.³² Unfortunately, the study population only included those with operative care and therefore does not reflect a standard sample of children with suspected appendicitis. Another




Figure 2. Positive predictive value of WBC count for appendicitis over a range of cut-points and stratified by age group. WBC = white blood cell.




Figure 3. Negative predictive value of WBC count for appendicitis over a range of cut-points and stratified by age group. WBC = white blood cell.

study of nontraumatic acute abdominal pain used the laboratory-derived range of normal values for WBC that were age-adjusted; although this study was not focused on patients with suspected appendicitis (only 10.2% of sample had appendicitis), the age adjustment was integrated into the definition of an abnormal result as a predictor of appendicitis.²⁶

LIMITATIONS

Our study has some strengths and limitations that deserve discussion. The prospective enrollment of a study sample across multiple sites allows for greater generalizability to a population of patients with suspected appendicitis. However, each site represented an academic pediatric ED that may not mirror the population evaluated in other settings; accordingly, the relatively high prevalence of appendicitis in the study

population influences the reported predictive values. We also acknowledge that we artificially dichotomized the patients into the outcomes of appendicitis and absence of appendicitis when in fact other diagnoses are important to consider and may explain some of the differences in discriminatory power of WBC count and ANC across age groups. Furthermore, we did not investigate the impact of race and sex on total WBC count and ANC, both of which have been previously shown to vary across populations of healthy adults. Finally, even with a cohort of over 2,000 patients, the sample size of children less than 5 years of age was relatively small and therefore estimates of diagnostic performance of the WBC count and ANC have less precision than the two other age groups.

CONCLUSION

The diagnostic performance of white blood cell count and absolute neutrophil count improves with increasing age. Age-adjusted values of white blood cell count or absolute neutrophil count should be considered in diagnostic strategies for suspected pediatric appendicitis. In the short term, clinical decision pathways might include age-adjusted thresholds for age groups. Future applications of these findings might integrate age-based predictive values of white blood cell count and absolute neutrophil count into electronic medical record-based decision support algorithms.

References

1. Russell WS, Schuh AM, Hill JG, et al. Clinical practice guidelines for pediatric appendicitis evaluation can decrease computed tomography utilization while maintaining diagnostic accuracy. *Pediatr Emerg Care* 2013;29:568–73.
2. Santillanes G, Simms S, Gausche-Hill M, et al. Prospective evaluation of a clinical practice guideline for diagnosis of appendicitis in children. *Acad Emerg Med* 2012;19:886–93.
3. Alvarado A. A practical score for the early diagnosis of acute appendicitis. *Ann Emerg Med* 1986;15:557–64.
4. Bachur RG, Callahan MJ, Monuteaux MC, Rangel SJ. Integration of ultrasound findings and a clinical score in the diagnostic evaluation of pediatric appendicitis. *J Pediatr* 2015;166:1134–9.
5. Blitman NM, Anwar M, Brady KB, Taragin BH, Freeman K. Value of focused appendicitis ultrasound and Alvarado score in predicting appendicitis in children: can we reduce the use of CT? *AJR Am J Roentgenol* 2015;204:W707–12.
6. Bond GR, Tully SB, Chan LS, Bradley RL. Use of the MANTRELS score in childhood appendicitis: a prospective study of 187 children with abdominal pain. *Ann Emerg Med* 1990;19:1014–8.
7. Samuel M. Pediatric appendicitis score. *J Pediatr Surg* 2002;37:877–81.
8. Fleischman RJ, Devine MK, Yagapen MA, et al. Evaluation of a novel pediatric appendicitis pathway using high- and low-risk scoring systems. *Pediatr Emerg Care* 2013;29:1060–5.

9. Saucier A, Huang EY, Emeremni CA, Pershad J. Prospective evaluation of a clinical pathway for suspected appendicitis. *Pediatrics* 2014;133:e88–95.
10. Kearl YL, Claudius I, Behar S, et al. Accuracy of magnetic resonance imaging and ultrasound for appendicitis in diagnostic and nondiagnostic studies. *Acad Emerg Med* 2016;23:179–85.
11. Orth RC, Guillerman RP, Zhang W, Masand P, Bisset GS 3rd. Prospective comparison of MR imaging and US for the diagnosis of pediatric appendicitis. *Radiology* 2015;277:927.
12. Frush DP, Frush KS, Oldham KT. Imaging of acute appendicitis in children: EU versus U.S. or US versus CT? A North American perspective. *Pediatr Radiol* 2009;39:500–5.
13. Garcia Pena BM, Cook EF, Mandl KD. Selective imaging strategies for the diagnosis of appendicitis in children. *Pediatrics* 2004; 113:24–8.
14. Doria AS, Moineddin R, Kellenberger CJ, et al. US or CT for diagnosis of appendicitis in children and adults? A meta-analysis. *Radiology* 2006;241:83–94.
15. Schuh S, Chan K, Langer JC, et al. Properties of serial ultrasound clinical diagnostic pathway in suspected appendicitis and related computed tomography use. *Acad Emerg Med* 2015;22:406–14.
16. Gronroos JM, Gronroos P. Leucocyte count and C-reactive protein in the diagnosis of acute appendicitis. *Br J Surg* 1999;86:501–4.
17. Kentsis A, Ahmed K, Kurek K, et al. Detection and diagnostic value of urine leucine-rich alpha-2-glycoprotein in children with suspected acute appendicitis. *Ann Emerg Med* 2012;60(78–83):e1.
18. Kharbanda AB, Rai AJ, Cosme Y, Liu K, Dayan PS. Novel serum and urine markers for pediatric appendicitis. *Acad Emerg Med* 2012;19:56–62.
19. Huckins DS, Simon HK, Copeland K, Spiro DM, Gogain J, Wandell M. A novel biomarker panel to rule out acute appendicitis in pediatric patients with abdominal pain. *Am J Emerg Med* 2013;31:1368–75.
20. Andersson M, Ruber M, Ekerfelt C, Hallgren HB, Olaison G, Andersson RE. Can new inflammatory markers improve the diagnosis of acute appendicitis? *World J Surg* 2014;38:2777–83.
21. Kharbanda AB, Taylor GA, Fishman SJ, Bachur RG. A clinical decision rule to identify children at low risk for appendicitis. *Pediatrics* 2005;116:709–16.
22. Atema JJ, Gans SL, Beenen LF, et al. Accuracy of white blood cell count and C-reactive protein levels related to duration of symptoms in patients suspected of acute appendicitis. *Acad Emerg Med* 2015;22:1015–24.
23. Beltran MA, Almonacid J, Vicencio A, Gutierrez J, Cruces KS, Cumsille MA. Predictive value of white blood cell count and C-reactive protein in children with appendicitis. *J Pediatr Surg* 2007;42:1208–14.
24. Kim HC, Yang DM, Lee CM, et al. Acute appendicitis: relationships between CT-determined severities and serum white blood cell counts and C-reactive protein levels. *Br J Radiol* 2011;84:1115–20.
25. Kohn MA. White blood cell count intervals and likelihood ratios for appendicitis. *Acad Emerg Med* 2005; 12:480; author reply 480–1.
26. Wang LT, Prentiss KA, Simon JZ, Doody DP, Ryan DP. The use of white blood cell count and left shift in the diagnosis of appendicitis in children. *Pediatr Emerg Care* 2007;23:69–76.
27. Yu CW, Juan LI, Wu MH, Shen CJ, Wu JY, Lee CC. Systematic review and meta-analysis of the diagnostic accuracy of procalcitonin, C-reactive protein and white blood cell count for suspected acute appendicitis. *Br J Surg* 2013;100:322–9.
28. Kharbanda AB, Cosme Y, Liu K, Spitalnik SL, Dayan PS. Discriminative accuracy of novel and traditional biomarkers in children with suspected appendicitis adjusted for duration of abdominal pain. *Acad Emerg Med* 2011;18:567–74.
29. Anandalwar SP, Callahan MJ, Bachur RG, et al. Use of white blood cell count and polymorphonuclear leukocyte differential to improve the predictive value of ultrasound for suspected appendicitis in children. *J Am Coll Surg* 2015;220:1010–7.
30. Keskek M, Tez M, Yoldas O, et al. Receiver operating characteristic analysis of leukocyte counts in operations for suspected appendicitis. *Am J Emerg Med* 2008;26:769–72.
31. Okamoto T, Sano K, Ogasahara K. Receiver-operating characteristic analysis of leukocyte counts and serum C-reactive protein levels in children with advanced appendicitis. *Surg Today* 2006;36:515–8.
32. Paajanen H, Mansikka A, Laato M, Kettunen J, Kosttinen S. Are serum inflammatory markers age dependent in acute appendicitis? *J Am Coll Surg* 1997;184:303–8.
33. Stefanutti G, Ghirardo V, Gamba P. Inflammatory markers for acute appendicitis in children: are they helpful? *J Pediatr Surg* 2007;42:773–6.
34. Xharra S, Gashi-Luci L, Xharra K, et al. Correlation of serum C-reactive protein, white blood count and neutrophil percentage with histopathology findings in acute appendicitis. *World J Emerg Surg* 2012;7:27.
35. Kharbanda AB, Dudley NC, Bajaj L, et al. Validation and refinement of a prediction rule to identify children at low risk for acute appendicitis. *Arch Pediatr Adolesc Med* 2012;166:738–44.
36. DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. *Biometrics* 1988;44:837–45.
37. Pregibon D. Logistic regression diagnostics. *Ann Stat* 1981;9:705–24.
38. Wu HP, Yang WC, Wu KH, Chen CY, Fu YC. Diagnosing appendicitis at different time points in children with right lower quadrant pain: comparison between Pediatric Appendicitis Score and the Alvarado score: reply. *World J Surg* 2012; 36:216–21.
39. Schellekens DH, Hulsewe KW, van Acker BA, et al. Evaluation of the diagnostic accuracy of plasma markers for early diagnosis in patients suspected for acute appendicitis. *Acad Emerg Med* 2013;20:703–10.
40. Sengupta A, Bax G, Paterson-Brown S. White cell count and C-reactive protein measurement in patients with possible appendicitis. *Ann R Coll Surg Engl* 2009;91:113–5.
41. Sack U, Biereder B, Elouahidi T, Bauer K, Keller T, Trobs RB. Diagnostic value of blood inflammatory

markers for detection of acute appendicitis in children. *BMC Surg* 2006;6:15.

Supporting Information

The following supporting information is available in the online version of this paper:

Data Supplement S1. Positive predictive value of absolute neutrophil count (ANC) for appendicitis over a range of cut-points and stratified by age group.

Data Supplement S2. Negative predictive value of absolute neutrophil count (ANC) for appendicitis over a range of cut-points and stratified by age group.

Announcing Usus – A community website on usage

Usus (Latin for usage) is a new, independent, community-run website (<http://www.usus.org.uk/>) for all those interested in the usage of online content. It is designed to support a productive conversation among librarians, publishers, aggregators, and repository managers so that we can all get the best possible usage reports for our electronic resources.

The Usus website provides:

- A source of hints and tips on solving known problems
- A list of vendors with problems that are affecting the credibility and/or usefulness of the COUNTER reports
- A collection point for suggestions for new COUNTER usage reports and metrics

Supervisory Board

The Usus Supervisory Board will ensure that the website is editorially independent and will serve the needs of the community. Chaired by Anne Osterman, Deputy Director of VIVA (the Virtual Library of Virginia), the members of the Supervisory Board are:

Anne Osterman, VIVA, USA (Chair)
Simon Bevan, Cranfield University, UK
Melissa Blaney, ACS Publications, USA
Anna Creech, University of Richmond, USA
Lorraine Estelle, JISC, UK
Oliver Pesch, EBSCO, USA
Kari Schmidt, Montgomery College, USA
Mark Tullos, ProQuest, USA

Travel Award

Thank you to Project COUNTER for the financial support needed to get Usus off the ground. COUNTER has also offered to provide a travel award worth £1,000/\$1,500 to a librarian who contributes the best opinion piece for the News & Opinions section of the Usus site. The award can be used to travel to the Charleston Conference, UKSG Conference, or Electronic Resources & Libraries Conference. Please send your submissions of 1,000 words or less to usus.stats@gmail.com by December 31, 2014.

For more information, please contact Usus at: usus.stats@gmail.com