

Children's Mercy Kansas City

SHARE @ Children's Mercy

Manuscripts, Articles, Book Chapters and Other Papers

4-1-2017

Impact of CYP2D6 genotype on amitriptyline efficacy for the treatment of diabetic peripheral neuropathy: a pilot study.

Mamoonah Chaudhry

Marco Alessandrini

Jacobus Rademan

Tyren M. Dodgen

Francois E. Steffens

See next page for additional authors

Follow this and additional works at: <https://scholarlyexchange.childrensmercy.org/papers>



Part of the [Endocrine System Diseases Commons](#), [Medical Genetics Commons](#), [Medical Pharmacology Commons](#), and the [Pharmaceutical Preparations Commons](#)

Recommended Citation



Chaudhry, Mamoonah; Alessandrini, Marco; Rademan, Jacobus; Dodgen, Tyren M.; Steffens, Francois E.; van Zyl, Danie G.; Gaedigk, Andrea; and Pepper, Michael S., "Impact of CYP2D6 genotype on amitriptyline efficacy for the treatment of diabetic peripheral neuropathy: a pilot study." (2017). *Manuscripts, Articles, Book Chapters and Other Papers*. 1107.

<https://scholarlyexchange.childrensmercy.org/papers/1107>

This Article is brought to you for free and open access by SHARE @ Children's Mercy. It has been accepted for inclusion in Manuscripts, Articles, Book Chapters and Other Papers by an authorized administrator of SHARE @ Children's Mercy. For more information, please contact bpfannenstiel@cmh.edu.

Creator(s)

Mamoonah Chaudhry, Marco Alessandrini, Jacobus Rademan, Tyren M. Dodgen, Francois E. Steffens, Danie G. van Zyl, Andrea Gaedigk, and Michael S. Pepper



Preliminary Communication

For reprint orders, please contact: reprints@futuremedicine.com

Impact of *CYP2D6* genotype on amitriptyline efficacy for the treatment of diabetic peripheral neuropathy: a pilot study

Aim: Therapy with low-dose amitriptyline is commonly used to treat painful diabetic peripheral neuropathy. There is a knowledge gap, however, regarding the role of variable *CYP2D6*-mediated drug metabolism and side effects (SEs). We aimed to generate pilot data to demonstrate that SEs are more frequent in patients with variant *CYP2D6* alleles. **Method:** To that end, 31 randomly recruited participants were treated with low-dose amitriptyline for painful diabetic peripheral neuropathy and their *CYP2D6* gene sequenced. **Results:** Patients with predicted normal or ultra-rapid metabolizer phenotypes presented with less SEs compared with individuals with decreased *CYP2D6* activity. **Conclusion:** Hence, *CYP2D6* genotype contributes to treatment outcome and may be useful for guiding drug therapy. Future investigations in a larger patient population are planned to support these preliminary findings.

First draft submitted: 26 November 2016; Accepted for publication: 13 January 2017; Published online: 28 March 2017

Keywords: activity score • adverse drug reactions • amitriptyline • *CYP2D6* • diabetic neuropathy

Diabetes is a public health problem worldwide and will be the seventh leading cause of death by 2030 (WHO). Its occurrence is greatest in low- and middle-income countries. In the sub-Saharan African region, almost 7 million people live with diabetes and this number is likely to double by 2025 [1]. Diabetic Peripheral Neuropathy (DPN) is the most common complication of diabetes and occurs in up to 50% of diabetic patients [2]. DPN is defined as 'the presence of symptoms and/or signs of peripheral nerve dysfunction in people with diabetes, after exclusion of other causes' [3]. The common clinical symptoms are allodynia, hyperalgesia and spontaneous pain [4]. About 26% of all diabetic patients suffer from neuropathic pain which is characterized as burning and shooting pain in the lower limbs and feet [5]. This discomfort often impacts directly on productivity, as reported by Gore *et al.*, who observed that

65% of working patients with DPN either missed work or experienced decreased work productivity due to pain [6].

There are no treatments that can restore nerve function, and hence drug therapy is mainly aimed at controlling pain [5]. However, up to two thirds of patients do not obtain adequate pain relief from their current drug regimens [7]. Drug therapy most often includes antidepressants (tricyclic antidepressants [TCAs], selective serotonin reuptake inhibitors and serotonin-noradrenaline reuptake inhibitors), sodium channel blockers and anticonvulsants [8]. Amitriptyline is the drug of choice for treatment of painful peripheral neuropathy in the South African public health sector. Given its cost-effectiveness, it is also widely used in other developing countries [9]. It is a TCA and is administered at doses of 150–250 mg/day for depression. However, when prescribed

Mamoonah Chaudhry¹, Marco Alessandrini¹, Jacobus Rademan^{1,2}, Tyren M Dodgen^{1,2}, Francois E Steffens³, Danie G van Zyl⁴, Andrea Gaedigk⁵ & Michael S Pepper^{*1}

¹Department of Immunology & the Institute for Cellular & Molecular Medicine, South African Medical Research Council Extramural Unit for Stem Cell Research & Therapy, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa

²Department of Pharmacology, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa

³Department of Consumer Science, Faculty of Natural & Agricultural Sciences, University of Pretoria, Pretoria, South Africa

⁴Department of Internal Medicine, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa

⁵Division of Clinical Pharmacology, Toxicology & Therapeutic Innovation, Children's Mercy Kansas City & School of Medicine, University of Missouri-Kansas City, Kansas City, MO 64108, USA

*Author for correspondence:

Tel.: +27 0 12 319 2190

Fax: +27 0 12 319 2946

michael.pepper@up.ac.za

for pain management, it is used at significantly lower starting doses of 10–25 mg/day [10]. Amitriptyline is mainly metabolized by CYP2C19 and CYP2D6, both of which are members of the CYP450 family of Phase I drug-metabolizing enzymes. The parent drug is rapidly demethylated by CYP2C19 to an active secondary amine, nortriptyline, which has both analgesic and antidepressant effects. Nortriptyline is then hydroxylated by CYP2D6 to form 10-hydroxynortriptyline, an inactive metabolite that is cleared by the kidneys. It has been shown that nortriptyline is predominantly responsible for the adverse drug reactions (ADRs) experienced in amitriptyline therapy [9]. If CYP2D6 clearance of nortriptyline is compromised due to absent or decreased enzyme activity, enhanced receptor occupation could potentially lead to ADRs.

The *CYP2D6* gene is located on chromosome 22q13.1 and has nine exons. It is highly polymorphic with over 100 allelic variants and subvariants having been designated by the Human Cytochrome P450 (*CYP*) Allele Nomenclature Database [11] (many additional variants have been observed, but have not been systematically captured). CYP2D6 represents only 2–4% of hepatic P450 protein, but is responsible for the metabolism of approximately 25% of drugs [12]. *CYP2C19* is also highly polymorphic. To date, 35 allelic variants have been defined not counting subvariants. The gene is located on chromosome 10q24 and contributes to the metabolism of about 10% of drugs [13]. The frequencies of *CYP2D6* and *CYP2C19* genetic variants vary across world populations and ethnic groups [14–16]. Phenotypically, people can be broadly classified according to their CYP2D6 and CYP2C19 activity: poor metabolizers (PMs, no activity), intermediate metabolizers (IMs, decreased activity), normal metabolizers (NMs, wild-type/normal activity) and ultra-rapid metabolizers (UMs, increased activity) [17]. PMs are at a higher risk of exhibiting side effects (SEs) or ADRs, while UMs may have a poor or absent response owing to subtherapeutic drug levels. Some of the SEs for amitriptyline include anticholinergic effects such as constipation, dizziness, dry mouth, blurred vision and urinary retention. The ADRs that are more severe and include cardiotoxicity and CNS toxicity [18].

The Clinical Pharmacogenetics Implementation Consortium (CPIC) has published guidelines providing dose recommendations for TCAs. This includes amitriptyline dosage and drug choice based on a patient's genotype that is used to optimize drug therapy and thereby minimize ADRs for TCAs according to *CYP2D6* and *CYP2C19* genotypes [18]. It needs to be noted though, that the recommendations are for the treatment of depressive disorders, and hence are in ref-

erence to the higher TCA dose ranges. Since amitriptyline is typically used at lower dosages for the treatment of neuropathic pain, CYP2D6 or CYP2C19 PMs or IMs are less likely to experience ADRs, but rather SEs. The recommendations by CPIC, in other words, that patients on low-dose TCAs should be closely monitored for SEs, have not changed in the recently published guideline update due to lack of information regarding low-dose amitriptyline therapy for neuropathic pain [19]. In our experience with low-dose amitriptyline, we often encounter SEs such as blurred vision, drowsiness, constipation, urinary retention and dryness of mouth/eyes in patients who take amitriptyline at lower dosages, which thus warrants further investigation.

Given the extensive and broad usage of amitriptyline for the management of chronic neuropathic pain in South Africa, and the well-described metabolic disposition of the drug, we hypothesized that pharmacogenetic testing might be useful for tailoring treatment and thereby improving amitriptyline efficacy in these patients. To test this hypothesis, and prior to embarking on a large-scale definitive study, we conducted a pilot investigation with the objectives of demonstrating proof of concept, and generating data for power calculations in order to assist in the planning of a larger scale study. For the purpose of this pilot study, the patient population was restricted to those with painful DPN, and pharmacogenetic testing was limited to *CYP2D6*, given that CYP2C19 contributes to the ratio of tertiary to secondary amine plasma concentrations; CYP2D6, however, is believed to have a larger impact on overall drug clearance than CYP2C19 [20]. Predicted phenotype status was correlated with SE scores that were derived from a daily questionnaire completed by the study participants.

Methods

Study population & questionnaire

Approval for this observational study was obtained from the University of Pretoria Research Ethics Committee (Ethics approval number 102/2005). Study participants over the age of 18 were recruited at random over a period of 6 months from the Diabetic Clinic at Kalafong Hospital (Atteridgeville, South Africa), and were included in the study after providing written informed consent. Participants were included if they had DPN and were on amitriptyline treatment for the management of chronic pain. Diagnosis of DPN was confirmed using monofilament and vibration sensory tests [21]. A monofilament test is commonly employed to evaluate peripheral nerve sensation in diabetic patients (usually done on the feet). The patient is asked to score the sensations caused by a thin fiber that is

used to touch the patient's foot at various positions. In the vibration sensory test, a vibration fork (128 Hz) is placed on the distal interphalangeal joint where the vibration sense should normally be felt. If it is not felt the vibration is sequentially tested on more proximal bony prominences. The lowest prominence felt is used to describe the level at which vibration can be felt.

Participants received a diary which they completed each day for 7 days following their clinic visit. The diary contained information regarding efficacy and SEs related to amitriptyline. Scoring was based on a rating scale from 1 to 10, where 1 indicated that the participant was completely satisfied and 10 indicated that the participant was completely dissatisfied. The diary comprised six questions that addressed the following for each day: the intensity of pain/burning in feet, blurred vision, constipation, drowsiness, dryness of the mouth and/or eyes, and difficulty urinating. Patients were asked to return the diary to the recruiter or study coordinator on the day of their next clinic visit or another suitable date. For interpretation, average scores of more than 5.0 were regarded as the threshold confirming the presence of pain or one of the recorded SEs.

Information regarding concomitant drugs (including doses) received by the patients, amitriptyline dose and ethnicity was recorded at the time of recruitment.

Sample processing & *CYP2D6* characterization

Blood was collected in ethylenediaminetetraacetic acid (EDTA) vacutainers and high-quality genomic DNA isolated from the leukocyte layer using the Maxwell®16 system (Promega, WI, USA). DNA was stored at -20°C prior to analysis.

Amplification of *CYP2D6*, detection of the *CYP2D6*5* gene deletion and gene duplication alleles was achieved using a modified tetramer multiplex assay [22]. Prior to sequencing, amplified products (5.1 kb) of *CYP2D6* were treated with FastAP Thermosensitive Alkaline Phosphatase (Fermentas Life Sciences, Vilnius, Lithuania) according to the supplied protocol. Sequencing of the entire gene (introns and exons) including 5'- and 3'-flanking regions (start and end positions for M33388, AY545216 and hg19 coordinates were: -1543–4180, 2658–8381 and 42,528,341–42,522,613, respectively) was performed by Inqaba Biotechnological Industries (Pty) Ltd. (Pretoria, South Africa), using an ABI Big Dye Terminator Cycle Sequencing kit version 3.1 and 3130 XL and 3500 XL DNA analyzer instruments (Applied Biosystems, Inc., CA, USA). Electropherograms were analyzed using CLC DNA Workbench version 5.5 (CLC bio, Aarhus, Denmark) and aligned to the National Center for Biotechnology Information (NCBI) AY545216

reference sequence for the detection and annotation of sequence variations. The assignment of alleles was done according to The Human Cytochrome P450 Allele Nomenclature Database for *CYP2D6* [23].

CYP2D6 genotype calls

Alleles were assigned using a set of 'key' SNPs defining star (*) alleles as shown in [Supplementary Table 1](#). These assignments were used to determine diplotypes and phenotype prediction as well as to calculate allele frequencies.

The majority of subjects contained additional SNPs that could not or only in part, be reconciled with allele definitions described by the nomenclature database. Novel variant haplotypes were manually inferred by taking into account allele definitions (per the nomenclature database), variants described previously [24] and novel variants found in this study that could unequivocally be determined to be due to the presence of the *CYP2D6*5* gene deletion ([Supplementary Table 1](#)). A subset of individuals carried novel SNP(s) that could not be assigned to either allele.

CYP2D6 phenotype prediction

CYP2D6 genotypes were translated into phenotype as previously described [25], using the activity score system ([Supplementary Table 2](#)). Briefly, the sum of values assigned to each allele provides the activity score that is used to group subjects into the following phenotype groups: 0 indicates PM status; a score of 0.5 indicates IM status; scores between 1 and 2 indicate NM status and a score more than 2 indicates UM status.

Statistical evaluation

The GraphPad Prism version 5 for Windows (GraphPad Software, CA, USA) was used to generate scatter plots.

For power calculations, and given that six measurables were reported in the participant questionnaires, multiple testing was corrected for by applying the Bonferroni inequality principle [26]. Therefore, each variable was measured against a significance level of 0.00833 (i.e., 0.05/6). Based on the data gathered during the preliminary study, the noncentrality parameters of the Analysis of Variance (ANOVA) F-tests [27] were calculated to compare the sample means. Based on the noncentral F-distribution, the power was calculated for overall sample sizes at 30, 60, 90, 120, 150 and 180.

Results

The sample cohort consisted of 21 Black African, nine Caucasian and one Indian participant. [Supplementary Table 1](#) provides a summary of the sequence variations and alleles identified. We observed 42 SNPs, a dele-

tion (del>C), two insertions (ins>G), a four-nucleotide insertion (ins>CAAC) and a series of SNPs defining the intron 1 conversion. We identified seven novel haplotypes: *1 V1, *1 V2, *1 V3, *1 V2xN, *2 V4 and V6 and *2 V1xN, found variants described previously [24] that are not cataloged by the P450 Nomenclature website, but also discovered a number of known and novel SNPs for which haplotype could not be determined. Of the novel SNPs, one was exonic (exon 5, 2509G>T, [K245N]). **Supplementary Table 1** provides the genotype assigned to each case considering all SNPs found (column B) and the genotype using only 'key' allele-defining SNPs (column D).

CYP2D6 allele frequencies

Allele frequencies (based on genotype assignments shown in **Supplementary Table 1** (column D) are summarized in **Table 1**. Considerable differences in frequencies were determined when the data were stratified according to ethnicity. In our Black African patients, the *CYP2D6**1 (33%) and *2 (19%) alleles were the most frequent, while the so-called African-specific alleles *CYP2D6**17 and *29 were less abundant at frequencies of 12 and 7.0%, respectively. The latter were absent in the Caucasian and Indian patients. The most common alleles among the nine Caucasian patients were *CYP2D6**2 and *4 (both 28%) followed by *CYP2D6**1 (22%). Gene duplications were only

observed in Black African patients. Also notable is the relatively high frequency of the *CYP2D6**5 gene deletion (14%) among the Black African patients. The alleles we have observed in our study population are consistent with those previously described for a South African population [28].

Phenotypic prediction of CYP2D6 & correlation with SE profiles

Genotypes were translated into predicted phenotype using the CYP2D6 activity score system [25]. As expected, the majority of study participants (n = 28; 90.32%) were classified as CYP2D6 NMs. Two patients (6.45%) were IM (one Caucasian and one Black African), and one Black African patient had a *CYP2D6* genotype predictive of UM status. We did not observe any genotypes predictive of a PM phenotype. For data analysis, subjects were grouped by activity score. As shown in **Figure 1B–F**, there is a trend for study participants with lower activity scores to report more severe SEs. These observations are in agreement with the expected higher amitriptyline and nortriptyline plasma levels in participants with decreased CYP2D6 activity.

By assigning average threshold scores of more than 5.0 as confirmation of the presence of a SE, it was possible to qualitatively report on the occurrence of SEs relative to predicted CYP2D6 phenotype (**Table 2**).

Table 1. *CYP2D6* allele frequencies in the South African diabetic peripheral neuropathy cohort sampled.

<i>CYP2D6</i> allele	Caucasians (n = 9) [†]	Black Africans (n = 21) [†]	Indians (n = 1) [†]	Total (n = 31) [†]	Dodgen et al. [‡] (n = 100) [†]
*1	4 (22)	14 (33)	1 (50)	19 (31)	57 (29.1)
*2	5 (28)	8 (19)	1 (50)	14 (22)	27 (13.8)
*4	5 (28)	0	0	5 (8)	6 (3.1)
*5	0	6 (14)	0	6 (10)	17 (8.7)
*17	0	5 (12)	0	5 (8)	38 (19.4)
*29	0	3 (7)	0	3 (4)	7 (3.6)
*35	2 (11)	0	0	2 (3)	1 (0.5)
*41	2 (11)	0	0	2 (3)	7 (3.6)
*43	0	1 (2)	0	1 (2)	3 (1.5)
*45B	0	1 (2)	0	1 (2)	9 (4.6)
*84	0	1 (2)	0	1 (2)	1 (0.5)
*1xn	0	1 (2)	0	1 (2)	–
*2xn	0	1 (2)	0	1 (2)	1 (0.5)
*4xn	0	1 (2)	0	1 (2)	4 (2.0)

[†]Indicates the number of study participants; in columns the number represents the allele count, and frequency (in percent) is provided in brackets.

[‡]Indicates *CYP2D6* allele frequencies in a representative South African cohort consisting of 70 black, ten Caucasian (white), ten colored (mixed-ancestry) and ten Indian (Asian) subjects [28].




Figure 1. Comparison of side effects experienced by study participants grouped according to their CYP2D6 activity score. Side effects (SEs) were scored on a scale of 0 to 10 (0 = no SEs and effective treatment; 10 = high number of SEs and ineffective treatment). CYP2D6 activity is represented as activity scores and the number of individuals per category is shown in brackets. Gray-shaded triangles and diamonds are for patients on CYP2C19 inhibitor drugs. The open square denotes a patient on a CYP2D6 inhibitor. The large solid triangle represents patient # 17 carrying the 2509G>T SNP in exon 5 that causes a K245N amino acid change. (B-F) Represent study participants with lower activity scores and severe SEs.

Table 2. Description of study participants based on predicted CYP2D6 phenotype and side effect profile.

ID	Ethnicity	Genotype	AS	Predicted phenotype	Pain intensity/burning feet	Blurred vision	Constipation	Drowsiness	Dryness of mouth/eyes	Difficulty in urinating	Number of SEs present >5	AT dose (mg)	Number of other drugs	Inhibitor drugs	Comments
1	Black	*1/*1xN	3	UM	1.3	1.3	1	1.3	1.9	1	0	25	11	Indo-metacin	Category I (1)
2	Black	*1/*45	2	NM	3.1	1.4	1	1.3	1.7	1.1	0	25	10		Category II (11)
3	Black	*1/*1	2	NM	1.9	2.6	3.6	3	2.7	3.3	0	25	12	Indo-metacin	
4	Black	*2/*2	2	NM	4.7	1.7	3.7	2.7	3.6	1.7	0	25	6		
5	Cauc	*2M/*35	2	NM	2.9	3	0.1	2.9	3.4	0	0	50	8		
6	Cauc	*1/*2	2	NM	1.7	3	1	2.1	2	1	0	25	ND		
7	Black	*2/*2	2	NM	10.0 [†]	1.1	1.3	1	1.1	1.1	0	25	5		
8	Black	*1/*1	2	NM	0.0	0	0	0	0.1	0	0	25	12		
9	Cauc	*35/*41	1.5	NM	1.3	3.1	2.9	4	3	4.6	0	ND	ND		
10	Black	*1/*17	1.5	NM	3.3	3.9	1.7	3.3	1.7	3.4	0	50	14	Indo-metacin	
11	Black	*2/*17	1.5	NM	1.3	1.4	1.7	1.9	1.3	1.9	0	10	7		
12	Cauc	*2/*4	1	NM	4.3	0.7	0	0	0	0	0	50	13		
13	Indian	*1/*2	2	NM	7.1 [†]	8 [†]	9.1 [†]	7.7 [†]	8.6 [†]	3.9	4	75	11		Category III (17)
14	Cauc	*1/*1	2	NM	8.0 [†]	6.1 [†]	5	5.1 [†]	5.7 [†]	4.9	3	25	12		
15	Black	*1/*2	2	NM	3.3	4.6	4.3	5.4 [†]	5	4.1	1	25	7		
16	Black	*2xN/*5	2	NM	5.9 [†]	0	2.1	5.4 [†]	5.9 [†]	1	2	ND	ND		
17	Black	*2/*29+1SNP (2509G>T)	1.5	NM	4.3	5.1 [†]	5.1 [†]	6.7 [†]	1.3	6.4 [†]	4	10	12		
18	Black	*1/*17	1.5	NM	7.4 [†]	3.4	5.4 [†]	4.1	6.3 [†]	3.1	2	25	10		
19	Black	*1/*17	1.5	NM	8.0 [†]	5	1.3	2.3	6.7 [†]	1.9	1	25	8	Indo-metacin	
20	Black	*1/*29	1.5	NM	6.4 [†]	0	3.7	0	0	5.1 [†]	1	25	9		
21	Black	*17/*84	1	NM	8.4 [†]	7.1 [†]	5.3 [†]	3.1	4.4	1	2	25	8		

[†]Individuals experiencing pain/burning feet with scores >5.0.

[‡]Average questionnaire scores of >5.0.

Genotype calls using only 'key' SNPs to determine the 'major' alleles; present in each subject are represented (for details see Supplementary Table 1, column D).

AS: Activity score; AT: Amitriptyline; Category I: No SEs + increased CYP2D6 activity; Category II: No SEs + normal CYP2D6 activity; Category III: SEs + normal CYP2D6 activity; Category IV: SEs + decreased CYP2D6 activity; Cauc: Caucasian; ID: Subject number; IM: Intermediate metabolizer; ND: No data; NM: Normal metabolizer; SE: Side effect; UM: Ultra-rapid metabolizer.

Table 2. Description of study participants based on predicted CYP2D6 phenotype and side effect profile (cont.).

ID	Ethnicity	Genotype	AS	Predicted phenotype	Pain intensity/burning feet	Blurred vision	Constipation	Drowsiness	Dryness of mouth/eyes	Difficulty in urinating	Number of SEs present >5	AT dose (mg)	Number of other drugs	Inhibitor drugs	Comments
22	Cauc	*2/*4	1	NM	6.9 [†]	6 [†]	9.1 [†]	7.3 [†]	6.6 [†]	4	4	75	8	Cimetidine	
23	Cauc	*2/*4	1	NM	8.0 [†]	7.3 [†]	8.1 [†]	6.9 [†]	7.3 [†]	3	4	10	9		
24	Black	*1/*5	1	NM	10.0 [†]	1	10 [†]	10 [†]	10 [†]	1.4	3	25	10		
25	Black	*1/*5	1	NM	7.0 [†]	6.7 [†]	7.3 [†]	6.7 [†]	6.3 [†]	5	4	25	15		
26	Black	*5/*43	1	NM	6.4 [†]	6.3 [†]	6.3 [†]	6.4 [†]	6.7 [†]	5.4 [†]	5	25	10		
27	Cauc	*1/*4	1	NM	6.4 [†]	4	8.3 [†]	4.3	6.6 [†]	3.3	2	100	ND		
28	Black	*2/*5	1	NM	2.9	7.9 [†]	1	9.9 [†]	4.1	1.1	2	25	10		
29	Black	*1/*5	1	NM	8.4 [†]	4.7	4	4.1	4.4	5.7 [†]	1	10	12		
30	Black	*4XN/*29	0.5	IM	0.7	3.1	9.6 [†]	0	4.3	2.1	1	25	5		Category IV (2)
31	Cauc	*4/*41	0.5	IM	5.1 [†]	0	2.4	0.3	5.1 [†]	7.1 [†]	2	ND	ND		

[†]Individuals experiencing pain/burning feet with scores >5.0.

[†]Average questionnaire scores of >5.0.

Genotype calls using only 'key' SNPs to determine the 'major' alleles present in each subject are represented (for details see Supplementary Table 1, column D).

AS: Activity score; AT: Amitriptyline; Category I: No SEs + increased CYP2D6 activity; Category II: No SEs + normal CYP2D6 activity; Category III: SEs + normal CYP2D6 activity; Category IV: SEs + decreased CYP2D6 activity; Cauc: Caucasian; ID: Subject number; IM: Intermediate metabolizer; ND: No data; NM: Normal metabolizer; SE: Side effect; UM: Ultra-rapid metabolizer.

Using this approach, we observed that IM individuals experienced up to two SEs during the period of evaluation. In contrast, 61% (17/28) of NMs experienced SEs and 39% (11/28) did not experience any SEs. Of those NMs who experienced 1–5 SEs, 53% had an activity score of 1 (less CYP2D6 activity compared with the patients with activity scores of 1.5 or 2). Of note, the UM individual did not report any SEs. The majority of the study participants were on 25 mg of amitriptyline (i.e., 64%; 18/28) at the time of recruitment. Participants were also taking a number of other drugs during the study (5–15 drugs including CYP2D6 or CYP2C19 inhibitors). In order to compare the predicted CYP2D6 phenotypes with participant scores for the severity of SEs, data were grouped into four categories. These include: category 1, no SEs and increased CYP2D6 activity; category 2, no SEs and normal CYP2D6 activity; category 3, SEs and normal CYP2D6 activity; and category 4, SEs and decreased CYP2D6 activity.

Power calculations

Given the distribution of data generated in this pilot study, power calculations were performed to assist in the planning of a larger scale study (Supplementary Table 3). The calculations indicate that for $n = 180$ subjects, each measurement will have a probability of at least 0.9 of being significant. With a choice of $N = 150$, one measurement will have a power of slightly below 0.9.

Discussion

Genetic variability of CYP2D6 & phenotype prediction

The most notable novel SNP is 2509G>T that was found in exon 5 in a Black African patient. This nonsynonymous SNP causes an amino acid change (K245N) having the potential to alter enzymatic activity. 2509G>T was observed in a *CYP2D6**2/*29 genotype (Table 2) and was classified as NM for this analysis. This patient experienced all SEs except dryness of eyes and mouth (subject 17 in Table 2) which suggests that K245N may cause decreased CYP2D6 function toward amitriptyline. The variant effect predictor available through Ensemble predicted that K245N has only a moderate impact. Further functional investigations are warranted, however, to fully elucidate the consequences of this amino acid change on catalytic activity.

Although sample size was a limitation in this investigation, allele frequencies (Table 1) were similar to those reported in the literature. Based on the literature for South African populations, 1–10% of the population are PM (>2% for Black Africans and 5–10% of Caucasians) [22,29,30]. Therefore, not having any PM

patients in our cohort falls within the expected norm. Albeit that we did not find a PM subject, one UM individual was identified that is consistent with previous reports [22,28].

Correlation of predicted CYP2D6 activity & reported SEs

The data were grouped into four categories to compare the predicted CYP2D6 phenotypes with participant scores for the severity of SEs. In the first category, the individual is UM and did not report any SEs. Pain was also adequately controlled. Individuals with UM status are likely to experience subtherapeutic levels of drug due to ultra-rapid metabolism. Of note, the individual was also taking indometacin; however, O-demethylation by CYP2D6 appears only to be a minor pathway [31,32] and therefore may not have interfered with amitriptyline metabolism in this UM patient. Taken together, low initial doses of amitriptyline, in other words, 25 mg, are not only effective but also safe and are in line with the CPIC dosing recommendations [18]. Individuals with NM status (second category) did not experience adverse effects due to normal enzyme activity. The majority (64%) of those patients had an activity score of 2, which predicts normal CYP2D6 activity. Most of the NMs were receiving 25 mg of amitriptyline while three were on 50 mg and one was on 10-mg amitriptyline. These individuals were also taking a range of other medications from a minimum of 5 to a maximum of 14 including CYP2C19 inhibitors (indomethacin). There was only one NM individual whose pain was not controlled and who was not experiencing other SEs. This individual was on 25 mg of amitriptyline and was using five other medications. Since none of the other drugs interfere with CYP2D6 activity, the dose of amitriptyline could be increased in this patient for more effective pain control. The third category includes individuals with normal CYP2D6 activity but who experience SEs. Out of 17 individuals, nine have an activity score of 1, four have an activity score of 1.5 and four have an activity score of 2. Similar to category 2, the majority of these individuals were receiving 25 mg of amitriptyline; three were receiving 10 mg; two were receiving 75 mg and one individual was on 100 mg. It should also be noted that these individuals were on average taking ten other medications including CYP2D6 inhibitors (cimetidine) and CYP2C19 inhibitors (indometacin), increasing the potential for drug–drug interactions. Individuals with an activity score of 1 reported more SEs compared with those with activity scores of 1.5 and 2. This observation is consistent considering that individuals with an activity score of 1 or 1.5 are predicted to have less metabolic activity

compared with individuals having an activity score of 2 [18]. Among the category 3 patients, amitriptyline treatment was only effective for three individuals and the remaining patients were still experiencing pain. Therefore, amitriptyline dose should not be increased in those patients to relieve the pain, as adverse effects might worsen. Treatment with an alternative should be considered for this group of patients. The fourth category comprised two IM individuals each possessing only one decreased function allele; both subjects experienced SEs.

The data generated in this pilot investigation suggest a relationship between risk of experiencing SE(s) and decreased *CYP2D6* activity. However, a number of limitations need to be considered. First, the sample size for this study was very small. As a result, there was only one UM and two IMs in our patient cohort. Second, the study cohort was genetically heterogeneous, and included individuals that were of Black African, Caucasian and Indian descent. Diverse genetic backgrounds may constitute an additional layer of complexity adding to the variability of results. Third, amitriptyline metabolism is not only dependent on *CYP2D6* but also *CYP2C19*. Other enzymes with minor contributions may also contribute to variability especially in patients with extreme *CYP2D6* and/or *CYP2C19* phenotypes, in other words, UMs and PMs. Fourth, *in vivo* *CYP2D6* and *CYP2C19* metabolic capacity (i.e., actual phenotype status) should be determined to refine the relationship between genotype and phenotype and the impact of concomitant medications. The metabolizer status of patients was predicted by genotype, which may not accurately reflect the actual phenotype status of a patient. Furthermore, for some alleles activity is unknown or uncertain. Fifth, plasma concentrations of amitriptyline and its metabolites should be measured to better understand the relationship between drug level, genotype, *in vivo* phenotype and adverse effects to facilitate the development of an algorithm for genotype-based amitriptyline dosing for neuropathic pain. Sixth, ideally the same low-dose should be given to each patient when therapy is initiated to monitor for SEs and aid in the evaluation of whether SEs are dose dependent, impacted by the patient's genetic makeup and/or are also influenced by other factors such as age, other health conditions or drug interactions. Finally, since the phenotypic component of the study was derived from daily questionnaires, participant subjectivity and bias are factors that may impact data interpretation. Although care was taken to ensure that the questions were phrased for easy interpretation and scoring, a certain level of subjectivity will inevitably result from self-scoring studies such as this.

Planning of large-scale definitive study

Both study objectives were met and a larger study is being planned. Based on the data generated in this pilot, a minimum cohort size of $n = 180$ will need to be recruited to be sufficiently powered. Our future investigation will be considerably more extensive and include, for example, phenotyping of both *CYP2D6* and *CYP2C19* using dextromethorphan and omeprazole as test drugs, measurement of amitriptyline plasma levels, genotyping of *CYP2D6* and *CYP2C19* and other pharmacogenes that are involved in the metabolism and disposition of amitriptyline and other commonly prescribed concomitant medications. It will likely be impossible, however, to completely rule out polypharmacy in a larger prospective study. When polypharmacy was included in power calculations, it was found that the use of chronic medications, in particular *CYP2D6* and/or *CYP2C19* inhibitors, should not affect the sample size requirements. Rather, the number of drugs taken by a patient can be used to explain some of the observed variability in pain response to amitriptyline. Furthermore, results obtained in non-polypharmacy patients would not reflect actual 'real-life' patient populations and therefore, would be of limited value. In this larger study, SEs will also be captured by a self-rating questionnaire completed by the patients, which will be carefully assessed as subjective measurements may impact SE scoring.

Conclusion

In conclusion, this study has found a genotype-phenotype correlation that may potentially be clinically relevant. Subjects with predicted normal or ultrarapid metabolizer phenotypes who are likely to have lower levels of amitriptyline, presented with fewer side effects compared to individuals with decreased or no *CYP2D6* activity who are likely to have higher amitriptyline levels.

Future perspective

The findings of our pilot study support undertaking a larger scale prospective study to address the limitations discussed above. Our preliminary findings also support the use of pharmacogenetic testing in the context of amitriptyline therapy for the management of diabetic pain. Given that *CYP2D6* also contributes to the metabolism of several other drugs used to treat DPN, such as duloxetine, various opioids and dextromethorphan [8,33], we argue that pharmacogenetic screening of *CYP2D6* together with *CYP2C19* will be valuable to guide drug choice and dosage and thereby improve treatment outcomes in patients with DPN. For instance, for *CYP2D6* UMs an alternative drug such as gabapentin may be a better choice to avoid risk

of therapeutic failure with amitriptyline. For PMs, lowering the drug dose may reduce SEs; however, if this is not sufficient, this may result in the prescription of an alternative drug.

Supplementary data

To view the supplementary data that accompany this paper, please visit the journal website at: www.futuremedicine.com/doi/full/10.2217/pgs-2016-0185

Financial & competing interests disclosure

This study was supported by funds provided by the South African Medical Research Council, the National Research Foundation of South Africa, the National Health Laboratory Services and the Institute for Cellular and Molecular Medicine, University of Pretoria. The authors have no other relevant affiliations

or financial involvement with any organization or entity on the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript.

Ethical conduct of research

The authors state that they have obtained appropriate institutional ethics approval for this study. For investigations involving human subjects, informed consent has been obtained from the participants involved.

Open access

This work is licensed under the Attribution-NonCommercial-NoDerivatives 4.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>

Executive summary

- An association between reported side effects to amitriptyline therapy and CYP2D6 metabolizer status was observed.
- This study provides preliminary data supporting the need for pharmacogenetic testing of CYP2D6 in order to individualize amitriptyline therapy for the management of pain.
- CYP2D6 gene sequencing revealed additional variation including one nonsynonymous SNP that could potentially impact enzyme function.

References

Papers of special note have been highlighted as:

• of interest; •• of considerable interest

- 1 International Diabetes Foundation. www.idf.org
- 2 Boulton AJM. Management of diabetic peripheral neuropathy. *Clin. Diabetes* 23, 9–15 (2005).
- 3 Boulton AJ, Gries FA, Jervell JA. Guidelines for the diagnosis and outpatient management of diabetic peripheral neuropathy. *Diabet. Med.* 15, 508–514 (1998).
- 4 Ulugol A, Karadag HC, Tamer M, Firat Z, Aslantas A, Dokmeci I. Involvement of adenosine in the anti-allodynic effect of amitriptyline in streptozotocin-induced diabetic rats. *Neurosci. Lett.* 328, 129–132 (2002).
- 5 Devi P, Madhu K, Ganapathy B, Sarma G, John L, Kulkarni C. Evaluation of efficacy and safety of gabapentin, duloxetine, and pregabalin in patients with painful diabetic peripheral neuropathy. *Indian J. Pharmacol.* 44, 51–56 (2012).
- 6 Gore M, Brandenburg NA, Hoffman DL, Tai K-S, Stacey B. Burden of illness in painful diabetic peripheral neuropathy: the patients' perspectives. *J. Pain* 7, 892–900 (2006).
- 7 Attal N, Cruccu G, Haanpää M *et al.* EFNS guidelines on pharmacological treatment of neuropathic pain. *Eur. J. Neurol.* 13, 1153–1169 (2006).
- 8 Callaghan BC, Cheng HT, Stables CL, Smith AL, Feldman EL. Diabetic neuropathy: clinical manifestations and current treatments. *Lancet Neurol.* 11, 521–534 (2012).
- 9 Steimer W, Zöpf K, Von Amelunxen S *et al.* Amitriptyline or not, that is the question: pharmacogenetic testing of CYP2D6 and CYP2C19 identifies patients with low or high risk for side effects in amitriptyline therapy. *Clin. Chem.* 51, 376–385 (2005).
- 10 Chetty S, Baalbergen E, Bhigjee AI *et al.* Clinical practice guidelines for management of neuropathic pain: expert panel recommendations for South Africa. *SAMJ.* 102, 312–325 (2012).
- 11 The Human Cytochrome P450 (CYP) Allele Nomenclature Database. www.cypalleles.ki.se/
- 12 Teh LK, Bertilsson L. Pharmacogenomics of CYP2D6: molecular genetics, interethnic differences and clinical importance. *Drug Metab. Pharmacokinet.* 27, 55–67 (2012).
- 13 Zhou S-F, Liu J-P, Chowbay B. Polymorphism of human cytochrome P450 enzymes and its clinical impact. *Drug Metab. Rev.* 41, 89–295 (2009).
- 14 Gaedigk A, Sangkuhl K, Whirl-Carrillo M, Klein T, Leeder JS. Prediction of CYP2D6 phenotype from genotype across world populations. *Genet. Med.* 19, 69–76 (2017).
- **This is the first comprehensive study that summarizes allele frequencies, diplotypes and predicted phenotype across major populations, providing a rich data resource for clinicians and researchers.**
- 15 Llerena A, Dorado P, Peñas-Lledó EM. Pharmacogenetics of debrisoquine and its use as a marker for CYP2D6 hydroxylation capacity. *Drug Metab. Pharmacokinet.* 10(1), 17–28 (2009).
- 16 Llerena A, Naranjo MEG, Rodrigues-Soares F, Penas-Lledó EM, Fariñas H, Tarazona-Santos E. Interethnic variability of CYP2D6 alleles and of predicted and measured metabolic phenotypes across world populations. *Expert Opin. Drug Metab. Toxicol.* 10(11), 1569–1583 (2014).

- 17 Caudle KE, Dunnenberger HM, Freimuth RR *et al.* Standardizing terms for clinical pharmacogenetic test results: consensus terms from the Clinical Pharmacogenetics Implementation Consortium (CPIC). *Genet. Med.* 19(2), 215–223 (2017).
- It is of considerable interest because it provides standardized pharmacogenetic terms that will facilitate reporting and sharing of pharmacogenetic tests.
- 18 Hicks JK, Swen JJ, Thorn CF *et al.* Clinical Pharmacogenetics Implementation Consortium guideline for CYP2D6 and CYP2C19 genotypes and dosing of tricyclic antidepressants. *Clin. Pharmacol. Ther.* 93, 402–408 (2013).
- It is of considerable interest because it provides dose recommendations for tricyclic antidepressants.
- 19 Hicks JK, Sangkuhl K, Swen JJ *et al.* Clinical Pharmacogenetics Implementation Consortium Guideline (CPIC®) for CYP2D6 and CYP2C19 genotypes and dosing of tricyclic antidepressants: 2016 update. *Clin. Pharmacol. Ther.* doi:10.1002/cpt.597 (2016) (Epub ahead of print).
- 20 Jiang Z-P, Shu Y, Chen X-P *et al.* The role of CYP2C19 in amitriptyline N-demethylation in Chinese subjects. *Eur. J. Clin. Pharmacol.* 58, 109–113 (2002).
- 21 Mayfield JA, Sugarman JR. The use of the Semmes–Weinstein monofilament and other threshold tests for preventing foot ulceration and amputation in persons with diabetes. *J. Fam. Pract.* 49(11), S17–S17 (2000).
- 22 Dodgen TM, Hochfeld WE, Fickl H *et al.* Introduction of the AmpliChip CYP450 test to a South African cohort: a platform comparative prospective cohort study. *BMC Med. Genet.* 14, 20 (2013).
- Provides useful information on genetic diversity of South African population in context to CYP2D6.
- 23 The Human Cytochrome P450 (CYP) Allele Nomenclature Database
www.cypalleles.ki.se/cyp2d6.htm
- 24 Twist GP, Gaedigk A, Miller NA *et al.* Constellation: a tool for rapid, automated phenotype assignment of a highly polymorphic pharmacogene, CYP2D6, from whole-genome sequences. *NPJ Genom. Med.* 1, 15007 (2016).
- 25 Gaedigk A, Simon SD, Pearce RE, Bradford LD, Kennedy MJ, Leeder JS. The CYP2D6 activity score: translating genotype information into a qualitative measure of phenotype. *Clin. Pharmacol. Ther.* 83, 234–242 (2008).
- The key paper on CYP2D6 activity score to determine individual phenotype from genotype.
- 26 Dunn OJ. Estimation of the means of dependent variables. *Ann. Math. Stat.* 29(4), 1095–1111 (1958).
- 27 Weisstein EW. Noncentral F-Distribution. (2002).
<http://mathworld.wolfram.com/>
- 28 Dodgen TM, Labuschagne CDJ, Van Schalkwyk A *et al.* Pharmacogenetic comparison of CYP2D6 predictive and measured phenotypes in a South African cohort. *Pharmacogenomics J.* 16, 566–572 (2015).
- 29 Gaedigk A, Coetsee C. The CYP2D6 gene locus in South African coloureds: unique allele distributions, novel alleles and gene arrangements. *Eur. J. Clin. Pharmacol.* 64(5), 465–475 (2008).
- 30 Wright GE, Niehaus DJ, Drögemöller BI, Koen L, Gaedigk A, Warnich L. Elucidation of CYP2D6 genetic diversity in a unique African population: implications for the future application of pharmacogenetics in the Xhosa population. *Ann. Hum. Genet.* 74(4), 340–350 (2010).
- 31 Nakajima M, Inoue T, Shimada N, Tokudome S, Yamamoto T, Kuroiwa Y. Cytochrome P450 2C9 catalyzes indomethacin O-demethylation in human liver microsomes. *Drug Metab. Dispos.* 26(3), 261–266 (1998).
- 32 Xie G, Zhou D, Cheng K-W, Wong CC, Rigas B. Comparative *in vitro* metabolism of phospho-tyrosol-indomethacin by mice, rats and humans. *Biochem. Pharmacol.* 85(8), 1195–1202 (2013).
- 33 Ziegler D, Fonseca V. From guideline to patient: a review of recent recommendations for pharmacotherapy of painful diabetic neuropathy. *J. Diabetes Complications* 29(1), 146–156 (2015).