

Children's Mercy Kansas City

SHARE @ Children's Mercy

Manuscripts, Articles, Book Chapters and Other Papers

5-1-2017

Role of biomechanical forces in hyperfiltration-mediated glomerular injury in congenital anomalies of the kidney and urinary tract.

Tarak Srivastava
Children's Mercy Hospital

Ganesh Thiagarajan

Uri S. Alon
Children's Mercy Hospital

Ram Sharma

Ashraf El-Meanawy

See next page for additional authors

Follow this and additional works at: <https://scholarlyexchange.childrensmercy.org/papers>



Part of the [Congenital, Hereditary, and Neonatal Diseases and Abnormalities Commons](#), [Nephrology Commons](#), [Pediatrics Commons](#), and the [Urogenital System Commons](#)

Recommended Citation

Srivastava, Tarak; Thiagarajan, Ganesh; Alon, Uri S.; Sharma, Ram; El-Meanawy, Ashraf; McCarthy, Ellen T.; Savin, Virginia J.; and Sharma, Mukut, "Role of biomechanical forces in hyperfiltration-mediated glomerular injury in congenital anomalies of the kidney and urinary tract." (2017). *Manuscripts, Articles, Book Chapters and Other Papers*. 1178.

<https://scholarlyexchange.childrensmercy.org/papers/1178>

This Article is brought to you for free and open access by SHARE @ Children's Mercy. It has been accepted for inclusion in Manuscripts, Articles, Book Chapters and Other Papers by an authorized administrator of SHARE @ Children's Mercy. For more information, please contact bpfannenstiel@cmh.edu.

Creator(s)

Tarak Srivastava, Ganesh Thiagarajan, Uri S. Alon, Ram Sharma, Ashraf El-Meanawy, Ellen T. McCarthy, Virginia J. Savin, and Mukut Sharma

Full Reviews

Role of biomechanical forces in hyperfiltration-mediated glomerular injury in congenital anomalies of the kidney and urinary tract

Tarak Srivastava^{1,2}, Ganesh Thiagarajan³, Uri S. Alon¹, Ram Sharma², Ashraf El-Meanawy⁴, Ellen T. McCarthy⁵, Virginia J. Savin² and Mukut Sharma²

¹Section of Nephrology, Children's Mercy Hospital and University of Missouri at Kansas City, Kansas City, MO, USA, ²Renal Research Laboratory, Research and Development, Kansas City VA Medical Center, Kansas City, MO, USA, ³School of Computing and Engineering, University of Missouri at Kansas City, MO, USA, ⁴Division of Nephrology, Medical College of Wisconsin, Milwaukee, WI, USA and ⁵Kidney Institute, Kansas University Medical Center, Kansas City, KS, USA

Correspondence and offprint requests to: Tarak Srivastava; E-mail: tsrivastava@cmh.edu

ABSTRACT

Congenital anomalies of the kidney and urinary tract (CAKUT) including solitary kidney constitute the main cause of progressive chronic kidney disease (CKD) in children. Children born with CAKUT develop signs of CKD only during adolescence and do not respond to renin-angiotensin-aldosterone system blockers. Early cellular changes underlying CKD progression to end-stage renal disease by early adulthood are not well understood. The mechanism of maladaptive hyperfiltration that occurs from loss of functional nephrons, including solitary kidney, is not clear. We re-examine the phenomenon of hyperfiltration in the context of biomechanical forces with special reference to glomerular podocytes. Capillary stretch exerts tensile stress on podocytes through the glomerular basement membrane. The flow of ultrafiltrate over the cell surface directly causes fluid flow shear stress (FFSS) on podocytes. FFSS on the podocyte surface increases 1.5- to 2-fold in animal models of solitary kidney and its effect on podocytes is a subject of ongoing research. Podocytes (i) are mechanosensitive to tensile and shear forces, (ii) use prostaglandin E₂, angiotensin-II or nitric oxide for mechanoperception and (iii) use specific signaling pathways for mechanotransduction. We discuss (i) the nature of and differences in cellular responses to biomechanical forces, (ii) methods to study biomechanical forces and (iii) effects of biomechanical forces on podocytes and glomeruli. Future studies on FFSS will likely identify novel targets for strategies for

early intervention to complement and strengthen the current regimen for treating children with CAKUT.

Keywords: CAKUT, fluid flow shear stress, hyperfiltration, podocytes, tensile stress

INTRODUCTION

Congenital anomalies of the kidney and urinary tract (CAKUT) including solitary kidney constitute the most common cause of chronic kidney disease (CKD) in children. CAKUT accounts for nearly 57% cases of CKD in children, according to the North American Pediatric Renal Trials and Collaborative Studies registry [1]. The NIH-CKiD (Chronic Kidney Disease in Children Study), USRDS (United States Renal Data System), and several European and Asian registries report the incidence of CAKUT as 49–62% in children with CKD and as 34–43% in those with end-stage renal disease (ESRD) [2–7]. The rate of decline in kidney function is related to modifiable factors such as proteinuria, hyperfiltration, blood pressure and diabetes, and to non-modifiable factors such as age, gender and ethnicity. Among the modifiable factors, hyperfiltration takes temporal precedence over others in modulating the progression of CKD.

In children with CAKUT, lack of postnatal nephrogenesis and hyperfiltration-mediated injury set up a vicious cycle of nephron loss with progressive CKD. Hyperfiltration not only results from but also contributes to ongoing and repetitive glomerular injury [8, 9]. With regard to the prevalence of risk

factors in 586 children (aged 1–16 years) in cohort I [glomerular filtration rate (GFR) 30–75 mL/min/1.73 m²] of the NIH-CKiD study none had diabetes, 84% were of normal weight, 46% had normal blood pressure and 24% showed normal urinary protein [7, 10]. These findings suggest that despite moderate CKD at baseline and ongoing progression on follow-up, hyperfiltration was the main driver of CKD progression during early years in children with CAKUT. Renin-angiotensin-aldosterone system (RAAS) blockers effectively delay the progression of the disease in some proteinuric CKD patients, but not all. Specifically, RAAS blockers have not been effective in delaying CKD progression in children with CAKUT [11].

Hyperfiltration is considered a long-known and frequently studied phenomenon, and early work has explored the physiological aspects of glomerular filtration. These studies and principles of fluid engineering suggest that glomerular capillary pressure (P_{GC}) maintains an outward flow of the plasma fluid while causing tensile stress on the capillary wall that is transmitted to podocytes. In parallel, continuous flow of the ultrafiltrate into Bowman's space generates shear stress on podocytes. Reconsidering hyperfiltration in terms of biomechanical forces provides an opportunity for identifying new facets of CKD progression in children with CAKUT. We discuss (i) the nature of and differences in the response of podocytes to tensile stress and shear stress, (ii) methods available to study biomechanical forces and (iii) effects of biomechanical forces on podocytes and glomeruli.

FLUID MECHANIC PERSPECTIVE OF STRESS AND STRAIN

The passage of blood through the capillary and the flow of the ultrafiltrate along the podocyte can be explored using principles of fluid mechanics. Capillary loops in the glomerulus appear as a coiled structure. Being hollow, the tubular structure of capillary may be visualized from within as a porous, thin-walled cylinder with holes and bends, with blood flowing under pressure. This structure results in one column of fluid (blood) that moves within the capillary and another (ultrafiltrate) that moves outside of the capillary between podocytes and over cell surface.

Stresses (force/area) are of two types, namely axial and shear. An object experiences strain when subjected to stress defined as the intensity of deformation resulting from the applied stress. Axial stress (σ), tensile or compressive in nature, exerts force perpendicular to the area to which it is applied and is expressed as $= \Delta l / l$, where Δl represents the change in length and l is the original length. Axial stress could be uniaxial, where force is applied along one of the axial directions or biaxial when force is applied along two perpendicular axes and the object stretches/compresses in two perpendicular directions.

Fluid passing through a tube (e.g. capillary) indicates fluid pressure within a hollow cylinder with closed ends. The closed end is an idealization for the bends in a tube, e.g. looped capillary. Radially directed internal pressure perpendicular to the wall of the tube (cylinder) and the axial thrust from the closed ends (a bend in the tube) stretch the walls of the cylinder in two axial directions. A tube is considered thin when r/t , the ratio of

the radius (r) and the thickness of the wall (t), is less than 10. Circumferential stretch results in a hoop stress in the cylinder wall given by $\sigma_{\theta} = p.r/t$. The axial stress, which is one half of the hoop stress, is given as $\sigma_a = p.r/2t$, where p is the pressure inside the capillary. The axial/hoop stress causes axial and hoop strains, due to the elastic modulus of components of the capillary wall. Thus, endothelial cells along the lumen, and the glomerular basement membrane (GBM) and podocytes covering the capillary will experience uniaxial tensile stress, and possibly biaxial tensile stress at the capillary bends (Figure 1).

The second type of stress is shear stress (τ) caused by a force parallel to the surface of the object and also defined as force divided by the area. However, shear strain (γ) is defined as change in angle (in radians) of an object exposed to shear stress. It should be noted that shear stress/strain on one surface automatically causes complementary shear stress on all surfaces, thereby changing its shape, e.g. change of a rectangular shape to parallelogram following deformation. Under small deformation and static behavior, the stresses and strains are related as follows: $\sigma = E\epsilon$ and $= G\gamma$, where E and G are the elastic modulus and shear modulus, respectively. The two moduli are related by the expression $= E/2(1+\nu)$, where ν is the Poisson's ratio of the membrane. The Poisson's ratio of a material represents the reduction in lateral strain when a membrane is stretched in the longitudinal direction.

Blood flow in capillaries over endothelium and ultrafiltrate flow between podocytes would be analogous to a fluid flowing over surface(s) of an object causing shear stress. Shear stress caused by a fluid column in a parallel plate configuration is given by the equation: $\tau = \frac{6\eta q}{wh}$, where τ is fluid flow shear stress (FFSS) in dynes/cm², w and h are the width and height of the fluid




FIGURE 1: Stresses, defined as force divided by a unit area, are of two types, namely axial and shear. Axial stress (σ) results from a force perpendicular to the area to which it is applied, and shear stress (τ) is caused by a force parallel to the surface of the area. Blood flow in glomerular capillary exerts force perpendicular to the capillary wall (two solid arrows) creating an axial (tensile) stress on podocytes localized along the outer aspect of capillary wall. The 3D architecture of podocyte attached to the GBM is highly complex. Each slit junction regulates the flow of a fraction of the total ultrafiltrate that flows into Bowman's space. Ultrafiltrate emerging through a large number of slit junctions between podocyte processes coalesces to form a fluid column. Dashed arrows represent fluid column from a number of filtration sites that are too numerous to depict in a simple cartoon.

column in cm, η is viscosity in centipoises, and q is flow rate in mL/s. Similarly, shear stress of a fluid column in a cylinder is given by the equation: $\tau = \frac{4\eta v}{r}$, where η is viscosity in centipoise, v is the velocity in mL/s and r is the radius in cm. In the glomerulus, the flowing fluid column (blood) within the capillary will create shear stress on endothelial cells. Similarly, the flowing fluid column (ultrafiltrate) creates shear stress on podocytes (Figure 1).

METHODS TO STUDY THE EFFECTS OF BIOMECHANICAL FORCES ON PODOCYTES AND GLOMERULI

Published research mentions Flexcell International Corporation (Burlington, NC, USA) and Strex Incorporation (Osaka, Japan) as major suppliers of equipment to study the effect of biomechanical forces. We have described a convenient flow chamber to study the effect of low FFSS on cells grown on glass slides or isolated rat glomeruli [12].

Tensile stress

Methods to study tensile or compressive stress (σ) are based on techniques to induce elongation or compression of cells or organelle (e.g. glomerulus). Podocytes are grown on a flexible membrane that constitutes the bottom of specially constructed cell culture dishes. Negative pressure (vacuum) causes a computer-controlled cyclical or sustained stretch in the adherent cell layer. Stretch can be applied in 1D or 2D, resulting in uniaxial or biaxial stress. Other variables include the (i) frequency (cycles/s, Hertz), (ii) magnitude (percentage change), (iii) pattern of application (square wave or sine wave) and (iv) the duration of stretch. Studies of podocytes have used 0.5–1 Hz, 2.5–20% stretch for 30 min to 3 days of continuous tensile stress. Additional methods to study axial stress include application of hydrostatic force, hyposmotic stretch or magnetic pull [13–15].

FFSS

Methods to study shear stress (τ) use technique(s) for controlled flow of a fluid column over podocytes or glomeruli. Fluid flow over the cells grown on glass slides illustrates a parallel plate geometry, where FFSS $\tau = \frac{6\eta q}{wh^2}$. In these instruments, height, width and viscosity of the fluid column (e.g. cell culture medium) are maintained constant. Therefore, the magnitude of FFSS is proportional to the rate of flow. Fluid column can be patterned as laminar, pulsatile or oscillating. Rotational flow of fluid is generated by placing the cell culture dish on an orbital shaker [16]. FFSS at the bottom of the dish is calculated using the equation $\tau_{\max} = \alpha[\rho\eta(2\pi f)^3]^{1/2}$, where α is the radius of orbit rotation, ρ is the density of the fluid medium, η is the viscosity of the fluid medium and f is the frequency of rotations per second. We have developed a flow chamber to study the effect of low FFSS on podocytes (0.2 dynes/cm²) and glomeruli (0.3 dynes/cm²) [17]. Others have described application of low FFSS between 0.015 and 1.75 dynes/cm², as well as a high range between 8 and 649 dynes/cm² [16, 18, 19].

Elastic modulus and shear modulus

Biophysical properties of podocytes and glomeruli [20, 21] include the elastic and shear moduli. Atomic force microscopy (AFM) and microindentation equipment are employed to measure the elastic modulus using information on the indentation depth, unloading slope and the type of indenter. The Oliver–Pharr methodology is generally used to determine the elastic indentation modulus. AFM revealed that the stiffness of HIV-infected podocytes is considerably lower than that of a normal podocyte [20]. Wyss *et al.* [21] measured the elastic properties (compressive modulus and shear modulus) of isolated mouse glomeruli using ‘capillary micromechanics’ and detected much lower Young’s modulus in glomeruli from mouse models of Alport syndrome or HIV-associated nephropathy compared with normal mouse glomeruli.

HYPERFILTRATION AND PODOCYTES

Human glomerular capillaries maintain transmural hydrostatic pressure gradient (ΔP) at ~ 60 mmHg [19] and filter plasma through a barrier to generate approximately 180 L ultrafiltrate/day. The filtration barrier is composed of capillary endothelial cells covered by the GBM, which, in turn, is supported from outside by podocyte foot processes that interdigitate to form adherens slitpore junctions. These structural features and spatial organization expose podocytes to biomechanical forces resulting from glomerular capillary stretch and plasma ultrafiltrate flow [22]. Capillary stretch exerts tensile stress on the capillary wall that transmits through the GBM to the podocyte, causing cellular deformation. In a conceptual model, tensile stress is sensed at the podocyte foot process, where the force generated by the fluid (blood) column will be perpendicular to the foot process. Kriz and Lemley [23] have suggested that the GBM, an elastic structure, counteracts tensile stress as podocytes are not true pericytes to develop an effective counterforce to tensile stress. Thus, tensile stress on the capillary wall is dampened by the elastic property of the GBM. Endlich and Endlich [22] calculated the uniaxial tensile stress on podocyte foot processes to be ~ 50 kPa based on ΔP of 40 mmHg and capillary radius and thickness of 5 and 0.5 μm , respectively. The biaxial tensile stress at a 90° capillary bend was calculated to be small at ~ 0.3 kPa.

The flow of ~ 180 L ultrafiltrate/day along the surface of podocytes would result in cellular deformation. In contrast to tensile stress, FFSS is exerted directly over the slit diaphragms, outer aspect of major processes and the soma of podocytes. Shear stress occurs when there is fluid flow along the surface of an object, albeit at different magnitudes based on the nature of the surface, viscosity, rate of the fluid flow, dimensions of the fluid column, etc. The architecture of podocytes attached to the GBM is complex. Each podocyte participates in the formation of a number of slit pore junctions through its foot processes that interdigitate with foot processes from adjacent podocytes. Each slit pore junction regulates the flow of a fraction of the total ultrafiltrate that flows into Bowman’s space. Thus, mathematical modeling for FFSS is convenient to understand by treating the ultrafiltrate as a single fluid column over podocytes. FFSS is

calculated by the equation $\tau = \frac{6\eta q}{wh^2}$ for rectangular flow or $\tau = \frac{4\eta v}{r}$ for cylindrical flow, indicating that FFSS is determined by the viscosity of the ultrafiltrate, flow rate [i.e. single nephron glomerular filtration rate (SNGFR)], and the dimensions of the fluid column. Endlich and Endlich [21] calculated the FFSS at the site of filtration slit diaphragm and the apical surface to be 8 Pa and 0.05 Pa, respectively. Friedrich *et al.* [18] calculated FFSS over the surface of the podocyte to be 0.3 dynes/cm² (or 0.03 Pa). Thus, under normal conditions, podocytes are exposed to tensile stress from P_{GC} and FFSS from ultrafiltration.

Using the equation $\tau = \frac{3\eta \cdot f \cdot \text{SNGFR}}{\pi \cdot s^2 \cdot (s + 2R_T)} \cdot \frac{z}{\sqrt{z \cdot (1-z)}}$ to estimate FFSS on the surface of podocytes, we showed a 1.5- to 2-fold increase in the calculated FFSS over podocytes in animal models of solitary kidney (Figure 2) [24]. We also found that: (i) increased SNGFR, not filtration fraction, was the basis of increased FFSS over podocytes in solitary kidney and (ii) the glomerular hypertrophy that accompanies adaptation to hyperfiltration did not compensate for the increased FFSS from increased SNGFR. No similar calculations have been made to determine changes in tensile stress with decreased functional nephron mass. Increased FFSS may contribute to detachment and loss of podocytes in animal models of hyperfiltration-mediated glomerular injury [23].

BIOMECHANICAL FORCES AND PODOCYTES

The effect of tensile stress on podocytes is well described. However, reports on the effect of shear stress on podocytes have started appearing only recently. Computational modeling shows that FFSS and tensile stress each may cause cellular deformation to different extent and may involve separate mechanisms [17].

Podocytes are intrinsically mechanosensitive

Mechanical stretch was found to cause narrowing of podocytes cell body, elongation of cytoplasmic processes and




FIGURE 2: Reduction in nephron number by uninephrectomy alters intrarenal hemodynamics to increase SNGFR. The calculated FFSS (τ) over podocytes using the equation $\tau = [(3\eta \cdot f \cdot \text{SNGFR} / \pi \cdot s^2 \cdot (s + 2R_T)) \cdot (z / \sqrt{z \cdot (1-z)})]$, where f is filtration fraction, $2R_T$ is diameter of glomerular tuft, s is width of Bowman's space and z is distance from the vascular to the urinary pole. The FFSS over podocytes in rats is increased 1.5- to 2-fold at 60 days after unilateral nephrectomy (shown in black) on Day 5 of life compared with control (shown in grey) using three separate models for 's'.

hypertrophy without proliferation [25]. Tensile stress (5% linear strain at 0.5 Hz for 24–72 h) decreases transversal stress F-actin filaments but microtubules and intermediate filaments appear unaffected. An increase in radial stress fibers, which converge on actin-rich centers, occurs in almost all cells by Day 3 and reverses to baseline within 1 week post-tensile stress [26, 27]. Actin cytoskeletal changes apparently involve α -actinin-4 [28] as well as Rho kinase activity, since Rho kinase inhibitor (Y-27632) blocks cellular changes. Tensile stress renders the actin cytoskeleton more sensitive to exogenous PGE₂ [29]. Overall, tensile stress alters podocyte morphology gradually through changes in the actin cytoskeleton that revert back to normal when tensile stress is stopped.

Conversely, podocytes are highly sensitive and respond immediately to FFSS [19]. FFSS increases membrane ruffling and macropinocytosis activity, diminishes transversal stress fibers and induces cortical actin within minutes. These changes can be blocked by pre-treating podocytes with indomethacin and recapitulated by treating podocytes with PGE₂ for 24 h without FFSS [19, 30].

Biomechanical forces may also activate pathways for cell death. Thus, while tensile stress up to 10% stretch or FFSS up to 2 dynes/cm² do not induce apoptosis in podocytes [19, 27, 30], tensile stress of 20% stretch [31] or FFSS of 10 dynes/cm² result in apoptotic changes [16].

Mechanoperception in podocytes

Deformation of the surface glycocalyx with the underlying stiff core proteins may mediate the perception of biomechanical forces in podocytes [32]. Among the known sensors of flow, primary cilia have been observed only in cultured podocytes and embryonic rat glomeruli. Slit diaphragm linked to adaptor molecules and actin cytoskeleton, and TRPC6 (ion channel) with BK_{Ca} (stretch channel) may also act as sensors for tensile stress [23, 33].

Mechanotransduction in podocytes

Molecules activated in response to perturbation of the plasma membrane include stretch channels and small effector molecules such as nitric oxide, Ca²⁺, angiotensin-II (Ang-II), Adenosine triphosphate (ATP) and PGE₂ [26, 34–36]. The following sections summarize the molecular changes that contribute to conversion of mechanical force into biochemical signals.

Tensile stress-induced molecular changes in podocytes

Both human and mouse podocytes show an increase in Ang-II secretion and expression of Ang-II Type 1 Receptor (AT1R) with a 50% reduction in nephrin expression when exposed to tensile stress for 24–72 h [26, 35]. The downregulation of nephrin could be blocked by AT1R antagonists or anti-Ang-II antibody. However, Liebau *et al.* [36] did not observe an increase in Ang-II associated with tensile stress in an earlier study.

Calcium mobilization is an important mechanism for transmitting rapid changes. Podocytes do not express voltage-gated Ca²⁺ channels. Endlich *et al.* [27] showed that tensile stress-induced changes in actin cytoskeleton were not blocked by gadolinium, an inhibitor of stretch-activated channel, but could be blocked by nickel, a non-specific calcium blocker. These




FIGURE 3: PGE₂ is increased in supernatant media at 2 h and 24 h following application of FFSS at 2 dynes/cm² for 2 h to podocytes compared with control (left). No such change in PGE₂ was seen in supernatant media at 2 h and 24 h following application of tensile stress of biaxial stretch of 5% at 1 Hz for 2 h (right).

observations suggest the significance of calcium influx and indicate the role of an unidentified channel.

Tensile stress increases the expression/release of secreted protein acidic and rich in cysteine (SPARC, osteonectin or BM-40) and osteopontin, but results to-date appear conflicting. SPARC is a matricellular protein that alters actin cytoskeleton through focal adhesion complexes and plays a role in wound healing, tissue remodeling and fibrosis. Tensile stress induces the expression and release of SPARC [37]. SPARC^{-/-} podocytes resist detachment by tensile stress compared with SPARC^{+/+} cells [38]. Tensile stress upregulates osteopontin expression in cultured podocytes, and in the DOCA salt-sensitive rat model of glomerular hypertension [39]. Gene deletion in mice showed osteopontin to be mechanoprotective. Osteopontin coating increases podocytes motility, accelerates actin cytoskeleton changes and diminishes cell loss in response to stretch [40]. The protective effects of osteopontin were mediated by increased $\alpha_v\beta_1$, $\alpha_v\beta_3$ and $\alpha_v\beta_5$ integrins. However, others observed that tensile stress increases cell detachment mediated by downregulation of $\alpha_3\beta_1$ integrin [31].

Intracellular signaling induced by stretch is not well understood. Some researchers have observed that *in vitro* application of stretch resulted in upregulation of p38 MAPK, ERK1/2 and JNK, but not Akt and GSK3 β . Other researchers, however, did not observe such changes [29, 37, 40–42].

Tensile stress activates arachidonic metabolism by cyclooxygenase-2 (COX-2) enzyme. Martineau *et al.* [29] found that tensile stress (1–8% stretch) resulted in increased expression of COX-2 but not COX-1, and EP₄ but not EP₁. A small increase in cAMP following tensile stress was significantly enhanced by exogenous PGE₂ with a greater disruption of the actin cytoskeleton.

FFSS-induced molecular changes in podocytes

Using laminar flow at 0.25–1.75 dynes/cm², Friedrich *et al.* [19] showed enhanced α -actinin at cell junctions, decreased vinculin, cortactin distributing to cell margin, and unchanged ZO-1 expression. While broadband tyrosine kinase inhibitors (TKI) such as genistein and AG 82 (Tyrphostin A25) caused podocyte loss, specific inhibitors such as PP2 (src family TKI) or

PD153035 (EGF receptor TKI) did not. The authors concluded that specific TKs play a protective role in podocytes against FFSS.

Huang *et al.* [16] applied rotational fluid movement with maximum FFSS of 10 dynes/cm². They demonstrated increased phospholipase D (PLD) activity, c-Src phosphorylation and mTOR activation, suggesting the importance of c-src/PLD1/mTOR pathway in podocytes exposed to FFSS.

We applied laminar FFSS (2 dynes/cm²) to podocytes and found increased synthesis and secretion of prostaglandin E₂ (PGE₂) during and after FFSS treatment [12, 30, 43]. Increased PGE₂ synthesis appears to be a specific response of podocytes to FFSS, as neither Martineau *et al.* [29] nor we detected increased PGE₂ in podocytes subjected to tensile stress (Figure 3). PGE₂ interacts with four membrane localized receptors, namely EP₁–EP₄, of which only EP₂ was upregulated by FFSS [43]. Changes in the actin cytoskeleton, COX-2, PGE₂ and EP₂ receptor induced by FFSS were blocked by indomethacin [12].

Hyperfiltration induced by unilateral nephrectomy of sv129 mice also increases the expression of COX-2 and EP₂ in the remnant kidney [12]. A similar increase in COX-2 expression and PGE₂ synthesis in renal-collecting duct cells treated with FFSS was shown by Rohatgi *et al.* [44, 45]. Preliminary results of our recent study suggest that application of FFSS induces an EP₂-mediated Akt-GSK3 β - β catenin and not the cAMP-PKA signaling pathway (unpublished observations) (see Figure 4).

Major differences between tensile stress and FFSS-induced changes in podocytes

Tensile stress (i) causes formation of actin-rich centers and radial stress fibers, (ii) upregulates COX-2 without increasing PGE₂ levels, (iii) upregulates EP₄, but not EP₂ and (iv) activates p38-MAPK and ERK1/2 pathways. On the other hand, FFSS (i) disrupts actin stress fibers with the formation of a cortical actin ring, (ii) upregulates COX-2 with increased PGE₂ levels, (iii) upregulates EP₂, but not EP₄ and (iv) activates Akt-GSK3 β - β catenin and c-src/PLD1/mTOR pathways. Thus, tensile stress and FFSS differ with regard to their effects on the actin cytoskeleton and specificity for PGE₂ receptors.

Effect of FFSS on glomeruli


Prolonged FFSS on podocytes alters glomerular filtration barrier characteristics. We have developed and used an *in vitro* method to study changes in glomerular albumin permeability (P_{alb}) as a surrogate of changes in glomerular function [46]. We exposed isolated decapsulated rat glomeruli to FFSS at 0.3 dynes/cm² for 2 h in a flow chamber designed in our laboratory. FFSS-induced increase in P_{alb} was replicated by incubation with exogenous PGE₂ without FFSS, and blocked by pre-treatment with indomethacin [12]. Thus, the COX2–PGE₂–EP₂ pathway appears to participate in cellular response to FFSS.

SIGNIFICANCE OF BIOMECHANICAL FORCES IN CAKUT

As previously mentioned, CAKUT is the most common cause of CKD, with hyperfiltration as a major driver of glomerular dysfunction in affected children. RAAS blockers do not effectively delay the progression of disease [11]. Approximately 50% of children born with solitary kidney progress to ESRD as young adults [47, 48]. The Italian (ItaKid) study found that most CAKUT children progressed to ESRD post-puberty [11].

Traditional risk factors of CKD including hypertension, diabetes or proteinuria do not arise until late puberty. Children with a solitary kidney manifest hypertension (16–26%), proteinuria (19–21%) or estimated GFR <60 mL/min/1.73 m² (6–10%)

Reduced Nephron Mass (CAKUT/Solitary Kidney)



Progression of Chronic Kidney Disease

FIGURE 4: Conceptual model for hyperfiltration-induced glomerular injury in CAKUT or solitary kidney in progression of CKD. Reduced number of functional nephron leads to an adaptive increase in RBF and P_{GC} . These changes result in increased SNGFR that results in increased FFSS over podocytes. RBF, renal blood flow; COX-2, cyclooxygenase-2 enzyme; PGE₂, prostaglandin E₂; EP₂, prostanoïd receptor; Akt, protein kinase B; GSK3β, glycogen synthase kinase 3 beta; cAMP, cyclic adenosine monophosphate; PKA, protein kinase A; Src, a non-receptor tyrosine kinase; EGFR, epidermal growth factor receptor.

at a median age of ~15 years [49, 50]. We submit that while persistent maladaptive hyperfiltration results in increased FFSS as well as tensile stress, FFSS is mainly responsible for early renal changes leading to progressive CKD in children born with CAKUT. While RAAS inhibitors can address glomerular hypertension, we believe ameliorating the effect of FFSS on podocytes will complement the current treatment approach.

SUMMARY AND FORWARD THOUGHTS

The term hyperfiltration-mediated injury has been used for decades but the mechanism of glomerular injury by hyperfiltration remains unclear. We have developed a paradigm that connects hyperfiltration-mediated podocyte injury to biomechanical forces. Advances in understanding mechanoperception and mechanotransduction of shear stress on podocytes will lead to novel targets to develop treatments for children with CAKUT.

ACKNOWLEDGEMENTS

Authors thank members of the laboratory who contributed to the research cited here. We acknowledge the contribution of a large number of investigators who have contributed to research on hyperfiltration and whose work could not be cited here.

FUNDING

This work was supported by NIDDK R01DK107490 (T.S.), Department of Veterans Affairs VA BX001037 (V.J.S.), DK 1R01 DK064969 (E.T.M.) and funds from the Midwest Biomedical Research Foundation (V.J.S. and M.S.). The views expressed in this article are those of the authors and do not necessarily reflect the position or policy of the Department of Veterans Affairs or the US government.

CONFLICT OF INTEREST

None declared.

REFERENCES

- Seikaly MG, Ho PL, Emmett L *et al*. Chronic renal insufficiency in children: the 2001 Annual Report of the NAPRTCS. *Pediatr Nephrol* 2003; 18: 796–804
- Ardissino G, Daccò V, Testa S *et al*. Epidemiology of chronic renal failure in children: data from the ItaKid project. *Pediatrics* 2003; 111: e382–e387
- Ishikura K, Uemura O, Ito S *et al*. Pre-dialysis chronic kidney disease in children: results of a nationwide survey in Japan. *Nephrol Dial Transplant* 2013; 28: 2345–2355
- McTaggart S, McDonald S, Henning P *et al*. Pediatric report. *ANZDATA Registry Report*. Adelaide, South Australia: Australia and New Zealand Dialysis and Transplant Registry, 2009
- ESPN/ERA-EDTA Registry Annual Report. <http://www.espn-reg.org> (2010) (17 June 2016, date last accessed).

6. USRDS 2013 Annual Data Report. <http://www.usrds.org/adr.aspx> (2013) (17 June 2016, date last accessed).
7. Wong CS, Pierce CB, Cole SR *et al.* Association of proteinuria with race, cause of chronic kidney disease, and glomerular filtration rate in the chronic kidney disease in children study. *Clin J Am Soc Nephrol* 2009; 4: 812–819
8. Brenner BM. Nephron adaptation to renal injury or ablation. *Am J Physiol* 1985; 249: F324–F337
9. Brenner BM, Lawler EV, Mackenzie HS. The hyperfiltration theory: a paradigm shift in nephrology. *Kidney Int* 1996; 49: 1774–1777
10. Furth SL, Abraham AG, Jerry-Fluker J *et al.* Metabolic abnormalities, cardiovascular disease risk factors, and GFR decline in children with chronic kidney disease. *Clin J Am Soc Nephrol* 2011; 6: 2132–2140
11. Ardissino G, Viganò S, Testa S *et al.* No clear evidence of ACEi efficacy on the progression of chronic kidney disease in children with hypodysplastic nephropathy-report from the ItalKid Project database. *Nephrol Dial Transplant* 2007; 22: 2525–2530
12. Srivastava T, Alon US, Cudmore PA *et al.* Cyclooxygenase-2, prostaglandin E2 and prostanoid receptor EP2 in fluid flow shear stress mediated injury in solitary kidney. *Am J Physiol Renal Physiol* 2014; 307: F1323–F1333
13. Coers W, Vos JT, Huitema S *et al.* Biological alterations of rat podocytes cultured under basolateral hydrostatic pressure. *Pathobiology* 1996; 64: 222–232
14. Anderson M, Kim EY, Hagmann H *et al.* Opposing effects of podocin on the gating of podocyte TRPC6 channels evoked by membrane stretch or diacylglycerol. *Am J Physiol Cell Physiol* 2013; 305: C276–C289
15. Glogauer M, Ferrier J. A new method for application of force to cells via ferri-oxide beads. *Pflugers Arch* 1998; 435: 320–327
16. Huang C, Bruggeman LA, Hydo LM *et al.* Shear stress induces cell apoptosis via a c-src phospholipase D-mTOR pathway in cultured podocytes. *Exp Cell Res* 2012; 318: 1075–1085
17. McGarry JG, Klein-Nulend J, Mullender MG *et al.* A comparison of strain and fluid shear stress in stimulating bone cell responses—a computational and experimental study. *FASEB J* 2005; 19: 482–484
18. Dandapani SV, Sugimoto H, Matthews BD *et al.* Alpha-actinin-4 is required for normal podocyte adhesion. *J Biol Chem* 2007; 282: 467–477
19. Friedrich C, Endlich N, Kriz W *et al.* Podocytes are sensitive to fluid shear stress in vitro. *Am J Physiol Renal Physiol* 2006; 291: F856–F865
20. Tandon R, Levental I, Huang C *et al.* HIV infection changes glomerular podocyte cytoskeletal composition and results in distinct cellular mechanical properties. *Am J Physiol Renal Physiol* 2007; 292: F701–F710
21. Wyss HM, Henderson JM, Byfield FJ *et al.* Biophysical properties of normal and diseased renal glomeruli. *Am J Physiol Cell Physiol* 2011; 300: C397–C405
22. Endlich N, Endlich K. The challenge and response of podocytes to glomerular hypertension. *Semin Nephrol* 2012; 32: 327–341
23. Kriz W, Lemley KV. A potential role for mechanical forces in the detachment of podocytes and the progression of CKD. *J Am Soc Nephrol* 2015; 26: 258–269
24. Srivastava T, Celsi GE, Sharma M *et al.* Fluid flow shear stress over podocytes is increased in the solitary kidney. *Nephrol Dial Transplant* 2014; 29: 65–72
25. Petermann AT, Pippin J, Durvasula R *et al.* Mechanical stretch induces podocyte hypertrophy in vitro. *Kidney Int* 2005; 67: 157–166
26. Durvasula RV, Petermann AT, Kiromura K *et al.* Activation of a local tissue angiotensin system in podocytes by mechanical strain. *Kidney Int* 2004; 65: 30–39
27. Endlich N, Kress KR, Reiser J *et al.* Podocytes respond to mechanical stress in vitro. *J Am Soc Nephrol* 2001; 12: 413–422
28. Michaud JL, Hosseini-Abardeh M, Farah K *et al.* Modulating alpha-actinin-4 dynamics in podocytes. *Cell Motil Cytoskeleton* 2009; 66: 166–178
29. Martineau LC, McVeigh LI, Jasmin BJ *et al.* P38 MAP kinase mediates mechanically induced COX-2 and PG EP4 receptor expression in podocytes: implications for the actin cytoskeleton. *Am J Physiol Renal Physiol* 2004; 286: F693–F701
30. Srivastava T, McCarthy ET, Sharma R *et al.* Prostaglandin E2 is crucial in the podocytes response to fluid flow shear stress. *J Cell Commun Signal* 2010; 4: 79–90
31. Dessapt C, Baradez MO, Hayward A *et al.* Mechanical forces and TGFbeta1 reduce podocyte adhesion through alpha3beta1 integrin downregulation. *Nephrol Dial Transplant* 2009; 24: 2645–2655
32. Tarbell JM, Weinbaum S, Kamm RD. Cellular fluid mechanics and mechanotransduction. *Ann Biomed Eng* 2005; 33: 1719–1723
33. Dryer SE, Reiser J. TRPC6 channels and their binding partners in podocytes: role in glomerular filtration and pathophysiology. *Am J Physiol Renal Physiol* 2010; 299: F689–F701
34. Weinbaum S, Duan Y, Thi MM *et al.* An integrative review of mechanotransduction in endothelial, epithelial (renal) and dendritic cells (osteocytes). *Cell Mol Bioeng* 2011; 4: 510–537
35. Miceli I, Burt D, Tarabra E *et al.* Stretch reduces nephrin expression via an angiotensin II-AT(1)-dependent mechanism in human podocytes: effect of rosiglitazone. *Am J Physiol Renal Physiol* 2010; 298: F381–F390
36. Liebau MC, Lang D, Böhm J *et al.* Functional expression of the renin-angiotensin system in human podocytes. *Am J Physiol Renal Physiol* 2006; 290: F710–F719
37. Durvasula RV, Shankland SJ. Mechanical strain increases SPARC levels in podocytes: implications for glomerulosclerosis. *Am J Physiol Renal Physiol* 2005; 289: F577–F584
38. Sussman AN, Sun T, Krofft RM *et al.* SPARC accelerates disease progression in experimental crescentic glomerulonephritis. *Am J Pathol* 2009; 174: 1827–1836
39. Endlich N, Sunohara M, Nietfeld W *et al.* Analysis of differential gene expression in stretched podocytes: osteopontin enhances adaptation of podocytes to mechanical stress. *FASEB J* 2002; 16: 1850–1852
40. Schordan S, Schordan E, Endlich K *et al.* AlphaV-integrins mediate the mechanoprotective action of osteopontin in podocytes. *Am J Physiol Renal Physiol* 2011; 300: F119–F132
41. Eekhoff A, Bonakdar N, Alonso JL *et al.* Glomerular podocytes: a study of mechanical properties and mechano-chemical signaling. *Biochem Biophys Res Commun* 2011; 406: 229–233
42. Faour WH, Thibodeau JF, Kennedy CR. Mechanical stretch and prostaglandin E2 modulate critical signaling pathways in mouse podocytes. *Cell Signal* 2010; 22: 1222–1230
43. Srivastava T, McCarthy ET, Sharma R *et al.* Fluid flow shear stress upregulates prostanoid receptor EP2 but not EP4 in murine podocytes. *Prostaglandins Other Lipid Mediat* 2013; 104–105: 49–57
44. Liu Y, Flores D, Carrisoza-Gaytán R *et al.* Biomechanical regulation of cyclooxygenase-2 in the renal collecting duct. *Am J Physiol Renal Physiol* 2014; 306: F214–F223
45. Flores D, Liu Y, Liu W *et al.* Flow-induced prostaglandin E2 release regulates Na and K transport in the collecting duct. *Am J Physiol Renal Physiol* 2012; 303: F632–F638
46. Savin VJ, Sharma R, Lovell HB *et al.* Measurement of albumin reflection coefficient with isolated rat glomeruli. *J Am Soc Nephrol* 1992; 3: 1260–1269
47. Sanna-Cherchi S, Ravani P, Corbani V *et al.* Renal outcome in patients with congenital anomalies of the kidney and urinary tract. *Kidney Int* 2009; 76: 528–533
48. Westland R, Schreuder MF, Bökenkamp A *et al.* Renal injury in children with a solitary functioning kidney—the KIMONO study. *Nephrol Dial Transplant* 2011; 26: 1533–1541
49. Westland R, Schreuder MF, Ket JC *et al.* Unilateral renal agenesis: a systematic review on associated anomalies and renal injury. *Nephrol Dial Transplant* 2013; 28: 1844–1855
50. Westland R, Kurvers RA, van Wijk JA *et al.* Risk factors for renal injury in children with a solitary functioning kidney. *Pediatrics* 2013; 131: e478–e485

Received: 30.8.2016; Editorial decision: 15.11.2016