

Children's Mercy Kansas City

SHARE @ Children's Mercy

Manuscripts, Articles, Book Chapters and Other Papers

5-7-2015

A patient with polymerase E1 deficiency (POLE1): clinical features and overlap with DNA breakage/instability syndromes.

Isabelle Thiffault

Children's Mercy Hospital

Carol Saunders

Children's Mercy Hospital

Janda Jenkins

Nikita Raje

Kristi Canty

See next page for additional authors

Follow this and additional works at: <https://scholarlyexchange.childrensmc.org/papers>



Part of the [Medical Genetics Commons](#)

Recommended Citation

Thiffault, I., Saunders, C., Jenkins, J., Raje, N., Canty, K., Sharma, M., Grote, L., Welsh, H. I., Farrow, E., Twist, G., Miller, N., Zwick, D., Zellmer, L., Kingsmore, S. F., Safina, N. P. A patient with polymerase E1 deficiency (POLE1): clinical features and overlap with DNA breakage/instability syndromes. *BMC medical genetics [electronic resource]* 16, 31-31 (2015).

This Article is brought to you for free and open access by SHARE @ Children's Mercy. It has been accepted for inclusion in Manuscripts, Articles, Book Chapters and Other Papers by an authorized administrator of SHARE @ Children's Mercy. For more information, please contact library@cmh.edu.

Creator(s)

Isabelle Thiffault, Carol Saunders, Janda Jenkins, Nikita Raje, Kristi Canty, Mukta Sharma, Lauren Grote, Holly I. Welsh, Emily Farrow, Greyson Twist, Neil Miller, David Zwick, Lee Zellmer, Stephen F. Kingsmore, and Nicole P. Safina

CASE REPORT

Open Access

A patient with polymerase E1 deficiency (*POLE1*): clinical features and overlap with DNA breakage/instability syndromes

Isabelle Thiffault^{2,3*}, Carol Saunders^{2,3,8}, Janda Jenkins^{1,7,8}, Nikita Raje⁴, Kristi Canty⁵, Mukta Sharma⁶, Lauren Grote^{1,7,8}, Holly I Welsh^{1,7,8}, Emily Farrow², Greyson Twist², Neil Miller², David Zwick³, Lee Zellmer², Stephen F Kingsmore^{2,3,7,8} and Nicole P Safina^{1,7,8}

Abstract

Background: Chromosome instability syndromes are a group of inherited conditions associated with chromosomal instability and breakage, often leading to immunodeficiency, growth retardation and increased risk of malignancy.

Case presentation: We performed exome sequencing on a girl with a suspected chromosome instability syndrome that manifested as growth retardation, microcephaly, developmental delay, dysmorphic features, poikiloderma, immune deficiency with pancytopenia, and myelodysplasia. She was homozygous for a previously reported splice variant, c.4444 + 3A > G in the *POLE1* gene, which encodes the catalytic subunit of DNA polymerase E.

Conclusion: This is the second family with *POLE1*-deficiency, with the affected individual demonstrating a more severe phenotype than previously described.

Keywords: *POLE1*, FILS syndrome, Immunodeficiency, Dysmorphism, Primordial dwarfism

Background

Chromosome instability syndromes are a group of inherited conditions associated with chromosomal instability and breakage which includes *LIG4* [1,2], Seckel type 1 [3,4], Bloom syndrome [5], Nijmegen breakage syndrome [6-8], and Fanconi anemia. These genetic conditions are characterized by pre and postnatal growth retardation, microcephaly, dysmorphic features and bone marrow failure [9-11]. There are other conditions with overlapping phenotypes including microcephaly, such as *Rad50* deficiency [12,13], Cernunnos-XLF syndrome [14] and Warsaw breakage syndrome [15-18] which can complicate molecular diagnosis. FILS syndrome (facial dysmorphism, immunodeficiency, livedo, and short stature) is a recently described condition caused by variants in *POLE1*, encoding the catalytic subunit of polymerase E. We describe the second family with a homozygous

variant in *POLE1*, and a more severely affected individual, suggesting a broader phenotypic spectrum for this condition.

Case presentation

Patient CMH812 is a female infant born to healthy non consanguineous Palestinian parents, weighing 1745 g and measuring 38.1 cm at birth. The pregnancy was complicated by subchorionic bleeding in the first trimester, fetal abnormalities on ultrasound including intra-uterine growth restriction, short long bones, suspected skull abnormalities and oligohydramnios. TORCH titers were negative. Amniocentesis revealed normal 46,XX karyotype. She was delivered at 37 weeks gestation by elective C-section secondary to breech presentation. Dysmorphic features noted included malar and mandibular hypoplasia (Figure 1A, B). Initial clinical suspicion was for primordial dwarfism such as Seckel type 1 syndrome, however her microcephaly was not as severe. Over several months, lacy reticular pigmentation was noted of the face and extremities. She had recurrent pruritic papular eruptions and skin findings progressed

* Correspondence: ithiffault@cmh.edu

²Center for Pediatric Genomic Medicine, Children's Mercy Hospital, Kansas City, MO 64108, USA

³Department of Pathology and Laboratory Medicine, Children's Mercy Hospitals, Kansas City, MO 64108, USA

Full list of author information is available at the end of the article




Figure 1 Clinical characteristics of CMH812. Several congenital abnormalities were found, including **(A)** microcephaly, facial dysmorphism (small and bird-like face, malar and mandibular hypoplasia, prominent nasal bridge and columella, downslanting palpebral fissures, small mouth and low set, posteriorly rotated ears) **(B)** short stature with shortened long bones but no evidence of dysplasia or craniosynostosis. No major anomalies were found on imaging of her abdomen, brain, brain vasculature or heart. Genitalia and pubertal development were normal. No malabsorption or pituitary or thyroid insufficiency was found. **(C-D)** Skin findings in CMH812, showing one hypo- and three hyperpigmented patches on the skin. Biopsy of the skin was performed but not diagnostic. Microscopic examination displayed focal parakeratosis and mild spongiosis.




Figure 2 Growth charts of CMH812. The patient is a female with severe intrauterine and postnatal growth retardation head circumference **(A)** [<3 rd centile]; weight **(B)** [<2 nd centile], length **(C)** [<2 nd centile], based on CDC growth chart for girls, age 0–20 months. Growth remained poor postnatally and she was briefly started on growth hormone, which was discontinued due to suspicion for a disorder of DNA repair/instability. At age 15 months she was ~ 5 SD below the mean for weight and length and ~ 3 –4 SD below the mean for head circumference.

Table 1 Comparison of clinical features of *POLE1* patients and other chromosome instability syndromes

Clinical features	TAR syndrome	Ataxia telangiectasia	RAD50 deficiency	Fanconi anemia	Seckel syndrome	NBS	RS/SC syndrome	Warsaw syndrome	Cernnos-XLF syndrome	Bloom syndrome	LIG4 syndrome2	FILS syndrome*	CMH812
MIM #	274000	208900	613078	227650	210600	251260	268300 /269000	613398	611291	210900	606593	615139	615139
Microcephaly	–	v	v	v	v	v	v	v	v	v	v	v	v
Malar hypoplasia	–	n.a	v	–	n.a	n.a	v	v	–	v	n.a	v	v
Sloping head	v	–	n.a	–	v	v	v	v	v	–	v	v	v
Palpebral fissures, upslanting	–	–	n.a	–	–	v	–	v	–	–	v	v	–
Palpebral fissures, down-slanting	–	–	v	–	v	–	v	v	v	–	–	v	v
Epicanthic folds	–	–	v	–	–	v	v	v	v	–	v	v	v
Micrognathia	v	–	n.a	–	v	v	v	v	v	–	–	v	–
External ear abnormalities	–	–	n.a	v	v	v	v	v	v	v	–	v	v
Long/Large nose	v	–	v	–	–	v	v	v	v	v	v	v	–
Long philtrum	–	–	n.a	–	–	v	v	v	v	–	–	v	v
Cleft palate/high arched palate	–	–	n.a	–	v	v	v	v	–	–	–	–	–
Polydactily	–	–	n.a	v	–	v	–	–	–	v	–	–	–
Clinodactily	–	–	n.a	–	v	v	v	v	–	v	v	–	v
Syndactily													
Growth retardation	–	v	v	v	v	v	v	v	v	v	v	v	v
Short stature	v	v	n.a	v	v	v	v	v	v	v	v	v	v
Bone disease or anomalies	v	–	n.a	v	v	–	v	n.a	v	–	–	v	v
Thumb apasia/radial ray anomalies	v	–	n.a	v	v	–	–/v	n.a	–	–	v	–	–
Skin abnormalities	v	v	v	v	v	v	v	v	v	v	v	v	v

Table 1 Comparison of clinical features of *POLE1* patients and other chromosome instability syndromes (Continued)

Mental retardation	v	v	v	v	v	v	v	v	v	–	v	–	–
Developmental delay	v	v	v	v	v	v	v	v	v	v	v	–	v
Malignancy	v	v	–	v	v	v	v	n.a	v	v	v	–	–
Recurrent infections	–	v	–	v	v	v	v	v	v	v	v	v	v
Genital abnormalities	–	–	–	v	v	v	v	n.a	–	–	v	–	–
Ataxia	–	v	v	–	v	n.a	–	n.a	–	n.a	n.a	–	–
Brain anomalies/ degeneration	CA, CH, aCC, S	CA, CH,T, S	ND	v	CA, CH, PG, S	ND	S	n.a	PG	n.a	n.a	n.a	–
Endocrine	–	AZ, D, DP, HH	–	HH	D, DP	POF	n.a	n.a	–	AZ, D RF	–	–	–
Immunologic Features													
Pancytopenia	–	n.a	n.a	v	v	–	n.a	n.a	v	–	v	–	v
Thrombocytopenia	v	n.a	–	v	–	–	n.a	n.a	–	–	–	–	v
CID	–	v	–	–	n.a	–	n.a	n.a	v	n.a	v	2/14	v
SCID	–	v	–	–	v	–	n.a	n.a	v	n.a	v	n.a	–
Neutropenia	–	n.a	n.a	v	n.a	v	n.a	n.a	v	N	v	n.a	–*
B cell lymphocytopenia	–	v	N	v	–	–	n.a	n.a	v	–	2/17	v	v
T cell lymphocytopenia	–	v	N	v	v	v	n.a	n.a	v	v	N	v	v
IgA	n.a	↓	N	N	↓	n.a	n.a	n.a	↓	↓	↓	N	↑
IgE	n.a	↓	N	n.a	↓	n.a	n.a	n.a	n.a	N	n.a	N	N
IgG	n.a	↓	N	↓	↓	n.a	n.a	n.a	↓	↓	n.a	↓	↓**
IgM	n.a	↓	N	↓	↓	n.a	n.a	n.a	↓	↓	↓	↓	↓
Anti-pseudomonae polysaccharide IgG	n.a	n.a	N	n.a	↓	n.a	n.a	n.a	v	n.a	n.a	v	v
Auto-immunity	n.a	v	–	n.a	↓	v	n.a	n.a	1/17	n.a	0/17	–	–
Sister chromatide	n.a	–	n.a	n.a	↑ or N	v	v	v	n.a	v	n.a	N	N
DNA breakage studies	n.a	v	v	v	v	v	v	v	v	v	n.a	–	N

Table 1 Comparison of clinical features of *POLE1* patients and other chromosome instability syndromes (Continued)

Radiosensitivity	n.a.	v	v	v	v	v	v	v	v	v	v	–	n.a.
Gene	RBM8A	ATM	RAD50	FANC	ATR	NBS1	ESCO2	DDX11	NHEJ1	BLM	LIG4	POLE1	POLE1
Mode of Inheritance	AR	AR	AR	AR	AR	AR	AR	AR	AR	AR	AR	AR	AR

Legend

n.a; not reported/applicable.

–; negative.

v; positive.

N; normal range.

↓; decreased.

↑; increased.

NBS: Nijmegen breakage syndrome.

TAR: Thrombocytopenia-absent radius syndrome.

RS: Roberts syndrome.

SC: SC phocomelia syndrome.

aCC; absence of corpus collosum.

AR; autosomal recessive.

AZ; azoospermia.

CA; cerebellar ataxia.

CH; cerebellar hypoplasia.

CID; combined immunodeficiency.

D; diabetes.

DP; delayed puberty.

HH; hypergonadotropic hypogonadism.

NBS; Nijmegen breakage syndrome.

ND; neurodegenerative.

OA; oculomotor apraxia.

PG; polygyria.

POF; primary ovarian failure.

RF; reduced fertility.

S; seizures.

SCID; severe combined immunodeficiency.

T; Tremor.

*consanguineous family reported.

£ Ataxia-telangiectasia-like disorder.

- * Transient pancytopenia associated with CMV infection.

↓** IgG2 and IgG 4 ↓. Total IgG N.

to include appearance of poikiloderma (Figure 1C, D). Erupted teeth were found to be small and dysplastic. She developed a feeding aversion necessitating a gastrostomy tube. Growth remained poor postnatally (Figure 2). Her motor milestones were delayed but social development was normal.

She suffered chronic rhinosinusitis and pulmonary infections with purulent otitis media. At age 20 months she was admitted to the hospital with pancytopenia, splenomegaly, hepatitis and acute CMV infection. Laboratory data showed mild bone marrow myelodysplasia, normal total B, T, and NK cells, low class switched and non switched memory B cells, and high memory T cells. She had high IgA, normal total IgG and low IgM, IgG2 and IgG4. There was no serologic response to pneumococcal vaccine (Table 1). Lymphocyte response to mitogens was normal but absent to pertussis and candida antigens [11]. Hepatitis and pancytopenia resolved following the CMV infection. Extensive molecular and cytogenetic testing was negative, including array-CGH, telomere length studies, chromosome breakage, and gene sequencing with deletion/duplication analysis for the following genes: *ATM*, *ATR*, *BLM*, *CENPJ*, *CEP152*, *CEP63*, *ERCC6*, *ERCC8*, *LIG4*, *LMNA*, *MRE11A*, *NBN*, *PCNT*, *RBBP8*, *RECQL4*, *SHOX*, *WRN*, and *ZMPSTE24*.

Trio-exome sequencing was performed on CMH812 and her healthy parents (CMH813 & CMH814) following informed consent, and with methods as previously published [19-22]. Variants were filtered to 1% minor allele frequency, then prioritized by the American College of Medical Genetics (ACMG) categorization [23,24], OMIM identity and phenotypic assessment. This individual was homozygous for a splice-site variant, c.4444 + 3A > G, in intron 34 of the *POLE1* [11]. The parents were both heterozygous carriers (Additional file 1: Table S1). Homozygosity mapper was used to identify intervals of homozygosity and identity by descent segments (Additional file 2: Figure S1) [25].

The c.4444 + 3A > G variant was previously reported in consanguineous French family with three generations of affected members [11]. FILS phenotype was variable but included macrocephaly, recurrent respiratory infections, livedo and telangiectasia, bone dysplasia, short stature, and decreased IgM and IgG. The phenotype was considered similar to that of Bloom syndrome but with normal sister chromatid exchange. Table 1 compares the clinical and cellular features of CMH812 to those of individuals with inherited chromosomal instability and breakage syndromes, as well as the first reported FILS family. Features closely matched those reported in FILS with exceptions of microcephaly and intrauterine growth restriction. Although the *POLE1* variant identified in the present case is the same as previously reported, CMH812

seems to have had more significantly impaired growth and immunity, raising the hypothesis that rare variant(s) in other *POLE* subunits or MMR genes may act as phenotypic modifiers. However, no rare variant were detected in MMR genes, *POLE1* interacting proteins or other DNA breakage/instability syndrome genes.

The c.4444 + 3A > G *POLE1* variant confers abnormal splicing whereby exon 34 is deleted [11] leading to significant decrease in the *POLE1* subunit [11]. T- lymphocytes from affected individuals showed a proliferation defect as well as impaired cell cycle progression. The primary function of polymerase ϵ 1 is to synthesize DNA at the leading strand during replication [26,27], however, it is also involved in other cellular processes, including cell cycle progression and DNA repair/recombination [26,27]. Exonucleolytic proofreading and the MMR pathway act to maintain high-fidelity DNA replication and to protect against mutagenesis [28]. Somatic and germline heterozygous missense variants in *POLE1* have been associated with an increased cancer risk [28-32]. Functional studies in yeast showed that heterozygosity for a pathogenic allele can cause complete MMR deficiency, and that subsequent loss of heterozygosity is not required for the development of *POLE*-related tumors [28]. Taken together, these findings suggest that *POLE1* carriers are likely to be at increased risk for malignancy due to MMR deficiency.

Conclusions

In summary, we report a second family with *POLE1*-related disease. The clinical and immunologic features of our patient are reminiscent of *LIG4* syndrome, possibly representing the more severe end of an ill-defined clinical spectrum. For this reason, *POLE1* deficiency may be a more apt description of this disorder. This report illustrates the cost-effectiveness of trio-exome sequencing as a powerful diagnostic method considering that this family underwent an extensive diagnostic odyssey, with no molecular basis identified prior exome.

Consent statement

The project was approved by the research ethics committee of the Children's Mercy Hospitals. Written informed consent was obtained from the patient's legal guardians for publication of this case report. A copy of the written consent is available for review by the Editor-in-Chief of this journal.

Additional files

Additional file 1: Table S1. Trio-Exome sequencing data.

Additional file 2: Figure S1. Homozygosity Mapping Analysis. Homozygosity mapper was used to identify intervals of homozygosity. The homozygosity scores are plotted against the physical position. Red

and black bars represent the excess (red) or the shortage (black) of homozygosity (A) Eighteen loci of > 1 Mb of homozygosity were identified on seven chromosomes; loci ranged between 1.06 and 6.37 Mb in length. The length of identical by descent segments in genomes of CMH812 parents suggests that they are not closely related. However, they shared several relatively small identical segments of genome which is explained by the shared ancestry broken into pieces by the recombination events and Mendelian laws of inheritance. *POLE1* lies in a 1.1 Mb region of homozygosity on chromosome 12 (chr12:132635257–133702440), which includes 20 genes (B). It is reasonable to hypothesize that the *POLE1* variant, if not the result of a combination of recent origin and chance, owes its origin to a founder who lived several hundred years ago. Consistent with the rarity of the c.4444 + 3A > G variant and the small physical distances between each haplotype marker in relationship to *POLE1* gene (1.1 Mb); the size of the shared haplotype may have been broken into smaller segments due to genetic drift. This assumption could explain that the same variant (c.4444 + 3A > G) has been found in two apparently unrelated FILS families.

Abbreviations

ACMG: American College of Medical Genetics; aCGH: Array of comparative genomic hybridization; DNA: Deoxyribonucleic acid; FILS: Facial dysmorphism immunodeficiency livedo and short stature; IUGR: Intrauterine growth restriction; LIG4: DNA ligase IV; Mb: Megabase pair; MMR: DNA mismatch repair; NK: Natural Killer; OMIM: Online Mendelian Inheritance in Man; NICU: Neonatal intensive-care unit; PCR: Polymerase chain reaction; *POLE1*: DNA polymerase E1; TAR: Thrombocytopenia absent radii syndrome; TORCH: Toxoplasmosis, other infections, rubella, cytomegalovirus (CMV), and herpes simplex virus (HSV). The “other infections” usually include syphilis, hepatitis B, coxsackie virus, Epstein-Barr virus, varicella-zoster virus, and human parvovirus; SD: Standard deviation; SNP: Single nucleotide polymorphism; XLF: XRCC4-like factor.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

The authors have made the following declarations about their contributions: Conceived and designed the experiments: CJS, EF, IT, GT. Performed the experiments: CJS, EF, IT, NK, NM, LZ. Contributed reagents/materials/analysis tools: NM, EF, DZ, GT. Wrote the paper: IT, CJS, NPS, JJ. Reviewed the manuscript: DZ, EF, KC, MS, HW, NM, NR, SFK. Contributed to the recruitment and clinical investigations of the patient for the study: DZ, HW, KC, JJ, MS, NK, NPS, NS. All authors read and approved the final manuscript.

Acknowledgments

We are grateful to the patient and her family. We thank the other staff of the Center for Pediatric Genomic Medicine, Children's Mercy Kansas City, and Dr. Geneviève de Saint Basile and her team for discussions. This work was supported by the Marion Merrell Dow Foundation, Children's Mercy - Kansas City, Patton Trust, W.T. Kemper Foundation, Pat & Gil Clements Foundation, Claire Giannini Foundation, and Black & Veatch.

Bioinformatics

The accession numbers for the reference sequences are listed as follows: *POLE1* NM_006231.2; HGMD 290829429, ClinVarRCV000034317.25 1000 Genomes Project, <http://www.1000genomes.org/> NHBLI Exome Sequencing Project (ESP) Exome Variant Server, <http://evs.gs.washington.edu/EVS/> Online Mendelian Inheritance in Man (OMIM), <http://www.omim.org> SNPdb, <http://www.ncbi.nlm.nih.gov/projects/SNP> UCSC Genome Browser, <http://genome.ucsc.edu/> PolyPhen-2, <http://www.genetics.bwh.harvard.edu/pph2/> ClinSeq, <http://genome.gov/20519355> RefSeq, <http://www.ncbi.nlm.nih.gov/proxy2.library.mcgill.ca/RefSeq> SIFT, sift.jcvi.org/

Author details

¹Division of Clinical Genetics, Children's Mercy Hospital, 2420 Pershing Road, Suite 421, Kansas City, MO 64108, USA. ²Center for Pediatric Genomic Medicine, Children's Mercy Hospital, Kansas City, MO 64108, USA.

³Department of Pathology and Laboratory Medicine, Children's Mercy Hospitals, Kansas City, MO 64108, USA. ⁴Pediatric Allergy, Asthma and Immunology Clinic, Children's Mercy Hospitals, Kansas City, MO 64108, USA. ⁵Dermatology Clinic, Children's Mercy Hospitals, Kansas City, MO 64108, USA. ⁶Department of Hematology and Oncology, Children's Mercy Hospitals, Kansas City, MO 64108, USA. ⁷Department of Pediatrics, Children's Mercy Hospitals, Kansas City, MO 64108, USA. ⁸University of Missouri, Kansas City School of Medicine, Kansas City, MO, USA.

Received: 10 February 2015 Accepted: 24 April 2015

Published online: 07 May 2015

References

- Rucci F, Notarangelo LD, Fazeli A, Patrizi L, Hickernell T, Paganini T, et al. Homozygous DNA ligase IV R278H mutation in mice leads to leaky SCID and represents a model for human LIG4 syndrome. *Proc Natl Acad Sci U S A*. 2010;107(7):3024–9.
- O'Driscoll M, Cerosaletti KM, Girard PM, Dai Y, Stumm M, Kysela B, et al. DNA ligase IV mutations identified in patients exhibiting developmental delay and immunodeficiency. *Mol Cell*. 2001;8(6):1175–85.
- Ogi T, Walker S, Stiff T, Hobson E, Limsirichaikul S, Carpenter G, et al. Identification of the first ATRIP-deficient patient and novel mutations in ATR define a clinical spectrum for ATR-ATRIP Seckel Syndrome. *PLoS Genet*. 2012;8(11), e1002945.
- O'Driscoll M, Ruiz-Perez VL, Woods CG, Jeggo PA, Goodship JA. A splicing mutation affecting expression of ataxia-telangiectasia and Rad3-related protein (ATR) results in Seckel syndrome. *Nat Genet*. 2003;33(4):497–501.
- Dubowitz V. Familial low birthweight dwarfism with an unusual facies and a skin eruption. *J Med Genet*. 1965;2(1):12–7.
- Curry CJ, O'Lague P, Tsai J, Hutchison HT, Jaspers NG, Wara D, et al. ATFresno: a phenotype linking ataxia-telangiectasia with the Nijmegen breakage syndrome. *Am J Hum Genet*. 1989;45(2):270–5.
- Cerosaletti KM, Lange E, Stringham HM, Weemaes CM, Smeets D, Solder B, et al. Fine localization of the Nijmegen breakage syndrome gene to 8q21: evidence for a common founder haplotype. *Am J Hum Genet*. 1998;63(1):125–34.
- Carney JP, Maser RS, Olivares H, Davis EM, Le Beau M, Yates 3rd JR, et al. The hMre11/hRad50 protein complex and Nijmegen breakage syndrome: linkage of double-strand break repair to the cellular DNA damage response. *Cell*. 1998;93(3):477–86.
- Ben-Omran TI, Cerosaletti K, Concannon P, Weitzman S, Nezarati MM. A patient with mutations in DNA Ligase IV: clinical features and overlap with Nijmegen breakage syndrome. *Am J Med Genet A*. 2005;137A(3):283–7.
- Gruhn B, Seidel J, Zintl F, Varon R, Tonnie H, Neitzel H, et al. Successful bone marrow transplantation in a patient with DNA ligase IV deficiency and bone marrow failure. *Orphanet J Rare Dis*. 2007;2:5.
- Pachlopnik Schmid J, Lemoine R, Nehme N, Cormier-Daire V, Revy P, Debeurme F, et al. Polymerase epsilon1 mutation in a human syndrome with facial dysmorphism, immunodeficiency, livedo, and short stature (“FILS syndrome”). *J Exp Med*. 2012;209(13):2323–30.
- Waltes R, Kalb R, Gatei M, Kijas AW, Stumm M, Soback A, et al. Human RAD50 deficiency in a Nijmegen breakage syndrome-like disorder. *Am J Med Genet A*. 2009;84(5):605–16.
- Stewart GS, Maser RS, Stankovic T, Bressan DA, Kaplan MI, Jaspers NG, et al. The DNA double-strand break repair gene hMRE11 is mutated in individuals with an ataxia-telangiectasia-like disorder. *Cell*. 1999;99(6):577–87.
- Buck D, Malivert L, de Chasseval R, Barraud A, Fondaneche MC, Sanal O, et al. Cernunnos, a novel nonhomologous end-joining factor, is mutated in human immunodeficiency with microcephaly. *Cell*. 2006;124(2):287–99.
- Wu Y, Sommers JA, Khan I, de Winter JP, Brosh Jr RM. Biochemical characterization of Warsaw breakage syndrome helicase. *J Biol Chem*. 2012;287(2):1007–21.
- van der Lelij P, Oostra AB, Rooimans MA, Joenje H, de Winter JP. Diagnostic overlap between fanconi anemia and the cohesinopathies: Roberts syndrome and Warsaw breakage syndrome. *Anemia*. 2010;2010:565268.
- van der Lelij P, Chrzanowska KH, Godthelp BC, Rooimans MA, Oostra AB, Stumm M, et al. Warsaw breakage syndrome, a cohesinopathy associated with mutations in the XPD helicase family member DDX11/ChIR1. *Am J Med Genet A*. 2010;86(2):262–6.

18. Capo-Chichi JM, Bharti SK, Sommers JA, Yammine T, Chouery E, Patry L, et al. Identification and biochemical characterization of a novel mutation in DDX11 causing Warsaw breakage syndrome. *Hum Mutat.* 2013;34(1):103–7.
19. Soden S, Saunders CJ, Willig LK, Farrow EG, Smith LD, Petrikin JE, et al. Effectiveness of acuity – guided exome and genome sequencing for etiologic diagnosis of neurodevelopmental disorders. *Sci Transl Med.* 2014. In press.
20. Saunders CJ, Miller NA, Soden SE, Dinwiddie DL, Noll A, Alnadi NA, et al. Rapid whole-genome sequencing for genetic disease diagnosis in neonatal intensive care units. *Sci Transl Med.* 2012;4(154):154ra135.
21. Bell CJ, Dinwiddie DL, Miller NA, Hateley SL, Ganusova EE, Mudge J, et al. Carrier testing for severe childhood recessive diseases by next-generation sequencing. *Sci Transl Med.* 2011;65):65ra64.
22. Soden SE, Saunders CJ, Willig LK, Farrow EG, Smith LD, Petrikin JE, et al. Effectiveness of exome and genome sequencing guided by acuity of illness for diagnosis of neurodevelopmental disorders. *Sci Transl Med.* 2014;6(265):265ra168.
23. Richards CS, Bale S, Bellissimo DB, Das S, Grody WW, Hegde MR, et al. ACMG recommendations for standards for interpretation and reporting of sequence variations: revisions 2007. *Genet Med.* 2008;10(4):294–300.
24. ACMG. Points to Consider in the Clinical Application of Genomic Sequencing. In: Genetics ACoM, editor. American College of Medical Genetics 14 edn. Bethesda, MD: American College of Medical Genetics; 2012. p. 759–61.
25. Seelow D, Schuelke M, Hildebrandt F, Nurnberg P. HomozygosityMapper—an interactive approach to homozygosity mapping. *Nucleic Acids Res.* 2009;37(Web Server issue):W593–599.
26. Huang D, Pospiech H, Kesti T, Syvaaja JE. Structural organization and splice variants of the POLE1 gene encoding the catalytic subunit of human DNA polymerase epsilon. *Biochem J.* 1999;339(Pt 3):657–65.
27. Huang D, Knuuti R, Palosaari H, Pospiech H, Syvaaja JE. cDNA and structural organization of the gene Pole1 for the mouse DNA polymerase epsilon catalytic subunit. *Biochim Biophys Acta.* 1999;1445(3):363–71.
28. Kane DP, Shcherbakova PV. A common cancer-associated DNA polymerase epsilon mutation causes an exceptionally strong mutator phenotype, indicating fidelity defects distinct from loss of proofreading. *Cancer Res.* 2014;74(7):1895–901.
29. Palles C, Cazier JB, Howarth KM, Domingo E, Jones AM, Broderick P, et al. Germline mutations affecting the proofreading domains of POLE and POLD1 predispose to colorectal adenomas and carcinomas. *Nat Genet.* 2013;45(2):136–44.
30. Briggs S, Tomlinson I. Germline and somatic polymerase epsilon and delta mutations define a new class of hypermutated colorectal and endometrial cancers. *J Pathol.* 2013;230(2):148–53.
31. Church DN, Briggs SE, Palles C, Domingo E, Kearsley SJ, Grimes JM, et al. DNA polymerase epsilon and delta exonuclease domain mutations in endometrial cancer. *Hum Mol Genet.* 2013;22(14):2820–8.
32. Zou Y, Liu FY, Liu H, Wang F, Li W, Huang MZ, et al. Frequent POLE1 p.S297F mutation in Chinese patients with ovarian endometrioid carcinoma. *Mutat Res.* 2014;761:49–52.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

