

Children's Mercy Kansas City

SHARE @ Children's Mercy

Manuscripts, Articles, Book Chapters and Other Papers

10-2018

Dietary Intake and Physical Activity Assessment: Current Tools, Techniques, and Technologies for Use in Adult Populations.

Holly L. McClung

Lauren T. Ptomey

Robin P. Shook
Children's Mercy Hospital

Anju Aggarwal

Anna M. Gorczyca

See next page for additional authors

Follow this and additional works at: <https://scholarlyexchange.childrensmercy.org/papers>



Part of the [Dietetics and Clinical Nutrition Commons](#), and the [Public Health Commons](#)

Recommended Citation

McClung, H. L., Ptomey, L. T., Shook, R. P., Aggarwal, A., Gorczyca, A. M., Sazonov, E. S., Becofsky, K., Weiss, R., Das, S. Dietary Intake and Physical Activity Assessment: Current Tools, Techniques, and Technologies for Use in Adult Populations. *American journal of preventive medicine* 55, 93-104 (2018).

This Article is brought to you for free and open access by SHARE @ Children's Mercy. It has been accepted for inclusion in Manuscripts, Articles, Book Chapters and Other Papers by an authorized administrator of SHARE @ Children's Mercy. For more information, please contact library@cmh.edu.

Creator(s)

Holly L. McClung, Lauren T. Ptomey, Robin P. Shook, Anju Aggarwal, Anna M. Gorczyca, Edward S. Sazonov, Katie Becofsky, Rick Weiss, and Sai Krupa Das

Dietary Intake and Physical Activity Assessment: Current Tools, Techniques, and Technologies for Use in Adult Populations



Holly L. McClung, MS,¹ Lauren T. Ptomey, PhD,² Robin P. Shook, PhD,³ Anju Aggarwal, PhD,⁴
Anna M. Gorczyca, PhD, MS,² Edward S. Sazonov, PhD,⁵ Katie Becofsky, PhD,⁶ Rick Weiss, MS,⁷
Sai Krupa Das, PhD⁸

Accurate assessment of dietary intake and physical activity is a vital component for quality research in public health, nutrition, and exercise science. However, accurate and consistent methodology for the assessment of these components remains a major challenge. Classic methods use self-report to capture dietary intake and physical activity in healthy adult populations. However, these tools, such as questionnaires or food and activity records and recalls, have been shown to underestimate energy intake and expenditure as compared with direct measures like doubly labeled water. This paper summarizes recent technological advancements, such as remote sensing devices, digital photography, and multisensor devices, which have the potential to improve the assessment of dietary intake and physical activity in free-living adults. This review will provide researchers with emerging evidence in support of these technologies, as well as a quick reference for selecting the “right-sized” assessment method based on study design, target population, outcome variables of interest, and economic and time considerations.

Theme information: This article is part of a theme issue entitled Innovative Tools for Assessing Diet and Physical Activity for Health Promotion, which is sponsored by the North American branch of the International Life Sciences Institute.

Am J Prev Med 2018;55(4):e93–e104. Published by Elsevier Inc. on behalf of American Journal of Preventive Medicine. This is an open access article under the CC BY-NC-ND license.

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

INTRODUCTION

Accurate assessment of dietary intake (DI) and physical activity (PA) is essential for quality research in the fields of public health, nutrition, and exercise science. However, consistent and accurate estimation of both remains one of the largest challenges in these fields. Several subjective and objective measures of DI and PA assessment exist, each with its own limitations and biases.

Capture of DI in healthy adult populations is intricate and multidimensional, thus making accurate quantification challenging. DI is traditionally assessed using self-report measures, including food frequency questionnaires (FFQs), diet records, and recalls.^{1–3} Such self-report measures have been shown to underestimate energy intake by approximately 11%–35% (more prevalent among

obese individuals) compared with direct measures like doubly labeled water.^{4–7} Reporting error that includes bias, also known as systematic error, misestimation, and

From the ¹Biophysics and Biomedical Modeling Division, U.S. Army Research Institute of Environmental Medicine, Natick, Massachusetts; ²Cardiovascular Research Institute, University of Kansas Medical Center, Kansas City, Kansas; ³Children’s Mercy Hospital, Kansas City, Missouri; ⁴Department of Epidemiology, University of Washington, Seattle, Washington; ⁵Department of Electrical and Computer Engineering, University of Alabama, Tuscaloosa, Alabama; ⁶Department of Kinesiology, University of Massachusetts, Amherst, Massachusetts; ⁷Viocare Inc., Princeton, New Jersey; and ⁸Jean Mayer USDA Human Nutrition Research Center on Aging, Tufts University, Boston, Massachusetts

Address correspondence to: Holly L. McClung, MS, Biophysics and Biomedical Modeling Division, U.S. Army Research Institute of Environmental Medicine, 10 General Greene Avenue, Building 42, Natick MA 01760. E-mail: holly.l.mcclung.civ@mail.mil

0749-3797/\$36.00

<https://doi.org/10.1016/j.amepre.2018.06.011>

random error, and error related to nutrient databases for foods being reported are a few of the current criticisms that have questioned the adequacy of self-report DI measures as the basis for scientific conclusions regarding the link between DI and health.^{8–11} From the findings in studies with doubly labeled water, researchers have suggested that self-report measures should not be used to estimate energy intakes, but that they are useful to estimate usual intakes of other nutrients and food groups and their densities, inform nutrition policy, and assess diet and disease associations.¹² Several recent reports suggest that investigators should work to improve and apply newer methods of DI assessment suitable for use in free-living individuals, such as biomarkers,^{4,13,14} remote sensing devices,^{15,16} or digital photography,¹⁷ rather than continue to rely solely on self-report methods.

PA is typically assessed using both self-report measures and devices. Self-report measures of PA include administration of questionnaires and completion of detailed diaries or logs. Device-based measures include motion sensors, such as accelerometers, pedometers, heart rate (HR) monitors, and multisensor devices.¹⁸ Because of the complex and multidimensional nature of PA, precise quantification can be difficult.¹⁹ Improvement and innovation are needed to provide low-cost, accurate measures of PA for use in both large and small samples of free-living healthy adults.

The use of technology for individualized DI and PA assessment has expanded rapidly over the past decade.^{20–24} Although technology has brought about some advances in diet and PA assessment methodology, many limitations and challenges remain. The purpose of this paper is to review the current science and challenges in the assessment of DI and PA for healthy adults and to identify current gaps and future needs.

DIETARY INTAKE ASSESSMENT

Methods of DI have been assessed using several objective and subjective tools, each with its inherent strengths and limitations. Selection of the right tool for use in research varies, depending on the study design, nutrients of interest, target population, and economic and time resources available. Some caution the adequacy of self-report DI measures as the basis for scientific conclusions regarding the link between DI and health outcomes.^{8–10} However, traditional DI assessment measures (FFQs, diet records, and recalls) remain the mainstay in the field based on their cost and familiarity, as well as lack of consensus among more objective methods capable of providing the complex output required. Although these measures may be criticized for not being precise, such data remain useful for population guidance in maintaining healthy

eating practices, comparison across populations, informing nutrition policy, and elucidating the associations between diet and disease.¹² Additional information on traditional DI methods and the controversy can be found in recent reviews by Farshchi et al.,²⁵ Dhurandhar and colleagues,⁹ Archer et al.,¹⁰ Shim and colleagues,²⁶ and Kirkpatrick et al.²⁷ Additionally, researchers are encouraged to utilize the Dietary Assessment Primer by National Cancer Institute (NCI) to help them determine the best way to assess diet for any study in which estimates of group intakes are required.²⁸

Current Dietary Intake Technology

Recent advances in technology have led to the development of several automated dietary assessment tools that have overcome some limitations of the traditional subjective tools, while striving to meet time and cost efficiency. Although modern DI methods are attractive, researchers should consider that these methods often do not differ in errors associated with underreporting and reactivity as compared with traditional methods. Current examples of modern methods include automated 24-hour recalls and food records,^{29,30} automated and graphic FFQs, photo-assisted dietary assessments (PADAs),^{31–38} and image-based dietary assessments (IBDAs).^{39–45} Table 1 summarizes the current and emerging DI assessment tools using technology.

The NCI introduced a modified version of the U.S. Department of Agriculture's Multiple-Pass 24-Hour Recall Method enabling 24-hour recalls to be self-administered by a respondent (ASA24) and used over multiple days as a food record.⁴⁶ Multiple versions (i.e., languages) have since been released and are detailed elsewhere.⁴⁷ The ASA24 improves on the limitations of traditional 24-hour recalls, including lack of reliance on trained interviewers, reduced time and economic burden to the researcher, and reduced respondent burden.⁴⁸ Because of the need for a high-speed Internet connection and familiarity with Internet or web-based tools, the use of the ASA24 may be limited in some populations.

In an effort to limit the issues with paper-based traditional FFQs,⁴⁹ a number of innovative web-based self-administered FFQs have been developed to automate the tool, such as the NCI Block questionnaire developed by Nutrition Quest,⁵⁰ NCI's Diet History Questionnaire (DHQ) III,⁵¹ and the Fred Hutchinson Cancer Research Center FFQ.⁵² All are web based and contain 100 or more questions on food items, purchasing, and preparation, with variations in layout design and analysis (e.g., food lists and databases) with NCI's DHQ III free for use by researchers. A novel alternative, VioScreen, offers a graphical FFQ option that addresses limitations of traditional FFQs by utilizing

Table 1. Summary of Current DI Assessment Tools Using Technology

Method/ tool	Outcome measure	Appropriate population	Attributes	Limitations	Validity	Research gaps	References
PADA	Researcher-defined nutrient output, such as energy intake, nutrient intake, food groups, and individual foods	Individuals and small groups	DLW validation; includes objective measure plus self-report	Participant burden to collect photos; researcher burden for post-analysis; self-report measure may not suitable for low-literacy populations	Estimate of energy intake –8.8% to 6.8% error compared to DLW	Automated post-analysis to include large food database	31–38
IBDA	Energy intake and volume estimates	Individuals and small groups; laboratory data collection	Low participant burden; no self-report	Data storage; high error rates or not validated to estimate energy/nutrient intake; ethical issues	Underestimated energy intake by ~23% compared to DLW; mean portion size difference compared to seed displacement is –5% ± 21.1%	Automatic analysis of data can estimate volume and requires food density to convert to nutrient intake; most food-nutrient databases lack density values	39–45
Automated 24-hour recall/ food record	Short-term DI, including energy intake, nutrient intake, food group, and individual foods; provides indicators of overall diet quality	Individuals, small groups, and large groups	Self-administered; eliminates the need for an interviewer and coding of intakes; captures short-term diet; accessible by individuals using assistive technologies, such as screen readers; uses images to assist respondents in reporting portion size	Restricted to populations with access to computers, high-speed Internet, and familiarity with web-based tools; not suitable for low-literacy populations	Underreporting of energy intake ~11% to 35% compared to DLW; 72% of items consumed were exact or close matches to reported; use to obtain food record data has not been evaluated	Accurately reports energy intake in normal-weight subjects; however, research is warranted to enhance its accuracy in overweight and obese individuals	8, 46–48
Automated FFQ	Frequency and portion size of foods and beverages consume over a long-term period; can also be used to assess usual DI or particular aspects of diet, including food groups and individual foods	Individuals, small groups, and large groups	Self-administered; low cost; low researcher burden; captures long-term diet (months); not affected by reactivity	Not suitable for low-literacy populations; restricted to populations with access to computers; limited application among ethnic populations due to its finite list of foods and beverages; poor measure of energy intake and some micronutrients with variable preparations; not useful for estimating a population's intake	Underestimated energy intake by 24% to 33% compared to DLW	Diverse food list/nutrient data for more universal use	49–52

(continued on next page)

Table 1. Summary of Current DI Assessment Tools Using Technology (*continued*)

Method/ tool	Outcome measure	Appropriate population	Attributes	Limitations	Validity	Research gaps	References
Graphic FFQ	Frequency and portion size of foods and beverages consumed over a long-term period; can also be used to assess usual DI or particular aspects of diet including food groups and individual foods	Individuals, small groups, and large groups	Improved DI and dietary pattern assessment through the use of improved portion size estimation via food images; self-administered; low cost; low researcher burden; uses branching logic to reduce completion time; captures long-term diet (months); not affected by reactivity	Not suitable for low-literacy populations; restricted to populations with access to computers; limited application among ethnic populations due to its finite list of foods and beverages; poor measure of energy intake and some micronutrients with variable preparations; not useful for estimating a population's intake	Compared to traditional FFQs, nutrient correlations are 0.90 for alcohol, 0.84 for saturated fat, 0.82 for fat, 0.79 for carbohydrate, and 0.67 for protein	Diverse food list/nutrient data for more universal use	53,54
Smart kitchen (e.g., plates, tables)	Researcher-defined nutrient output; frequency and portion size of foods and beverages consumed over a long-term period	Individuals and small groups; laboratory or home-based data collection	Reduced participant burden; streamlined researcher collection and analysis	Limited eating environments; strength of nutrient data is dependent on database used for coding	Validity based on quality of inputs, including weights, images, and sensor-based data; nutrient database selection	Use in real time; development of enhanced computer vision systems; validation studies	55,56
UPC or grocery store purchase data	Nutrients limited to food label; foods and beverages purchased over a long-term period	Large populations, as an adjunct to mobile apps	Ease of collection, time efficient, and minimal training of participants	Data are of food purchases and not consumed intake (DI assumed); large amount of data; individualized DI difficult to interpret; nutrients limited to food label (some missing nutrient data); privacy concerns	Association of foods purchased to food group mapping: 77%–100%; agreement between UPC scanned data home food inventory: ~95%	Real-time data use and feedback; accountability for waste; validation studies; database use transparency	57,58
Body-worn monitors	Time and duration of food intake; meal microstructure; estimates of mass and energy; food imagery	Individuals and small groups	Potential ease of data collection; no self-report in some methods; potential use in just-in-time interventions	Not well tested (yet); sensors may not detect certain foods; the nutritional value of ingested foods is not measured directly; stigma to wearing the device; personal privacy of bystanders	Up to 90.1% accurate at identifying when a person is consuming food	Large-scale validation across populations and environments	59–67

DI, dietary intake; DLW, doubly labeled water; FFQ, food frequency questionnaire; IBDA, image-based dietary assessment; PADA, photo-assisted dietary assessment; UPC, Universal Product Code.

branching questions (i.e., avoids lost data and limits respondent burden) and offers multiple photographs for each food item to accurately capture serving size (i.e., avoids need for respondents to calculate DI into a standard serving size).^{53,54}

PADAs, in which images of food selections and any food remaining after the meal are used to estimate DI, may provide an efficient, unobtrusive method for DI assessment in large groups of free-living individuals. PADAs have been utilized to assess DI in military recruits during basic training,^{31,32} young adults,^{33,34} individuals with disabilities,^{35,36} and overweight and obese women.³⁷ PADA methods include traditional methods (photo match to weighed food standard),³⁸ as well as technological advancements with remote food photography³⁷ and digital photography plus recall,³⁴ both validated to direct energy measures (e.g., doubly labeled water) under different environmental and population extremes. PADA is limited by a lack of full automation for nutrient analysis after photo capture and the quality of the nutrient database used in analysis.

IBDAs are a technique in which images of food selections and any food remaining after the meal are used to estimate DI. Unlike PADA, IBDA image capture is passive (e.g., automatic from the device) and relies on the captured images as the main source of information with input from the user only for verification.³⁹ Updated versions of IBDAs have combined with automated food identification and portion size estimation software, as well as user prompts, in an attempt to accurately assess DI. Examples include the Nutricam Dietary Assessment Method,⁴⁰ the eButton,⁴¹ and the Technology-Assisted Dietary Assessment system.^{42–44} Most advanced is the mobile food intake visual and voice recognizer,⁴⁵ which incorporates mobile phone food photography methods using image recognition with speech recognition and physical location (mobile phone accesses to a GPS). Although IBDA minimizes participant and researcher burden during data collection, the amount of data influx is vast and requires further work to streamline data cleaning and analysis for efficient researcher/user feedback.

Emerging Dietary Intake Technology

One of the largest areas of technological growth is in sensors for DI assessment.⁵⁵ A majority of technologies are geared toward the consumer and have inherent flaws (per the research community), whereas others are in their infancy and show potential with future improvements and testing.

In an effort to improve data accuracy and participant burden, some techniques and tools aim to identify foods and portions consumed through the

automation at the point of sale or food preparation (e.g., the kitchen). Ease of capturing Universal Product Codes⁵⁷ and Global Trade Item Numbers with handheld scanners or smartphones enables the correct item type to be properly linked with serving size and nutrition information at the time of consumption.⁵⁸ Alternatively, use of grocery store receipts (e.g., data capture) is an attractive option to minimize participant burden with direct feed into a food record for timely DI assessment (e.g., eliminate matching food type and brand consumed with specific database item). Traditional portion size estimation methods use standardization tools, cards, or even anatomical measures (e.g., the user's thumb) as a reference for improved accuracy of written DI assessments or PADAs.⁵⁵ Preliminary studies have looked at the use of smart kitchen equipment (e.g., plates, bowls, and tables) capable of recording food weight (with or without plates) before and after meal consumption.⁵⁶

Wearable sensors offer automated capture of food consumptions through hand-to-mouth gestures,^{59,60} modality of chewing (e.g., microphones to detect food crushing,⁶¹ electromyographic sensors to detect muscle activations,⁶² or strain and acceleration sensors to capture the chewing motion⁶³), or swallowing frequency.^{64–66} Chewing monitors have been shown to be reliable indicators of ingestion in community-dwelling individuals.¹⁶ Of interest, chew counts show good correlation to ingested food mass.⁶⁷ However, they may be prone to false detections (e.g., because of gum chewing) and may not detect all liquids, although consumption of certain liquids (e.g., sucking through a straw) creates jaw motion similar to chewing and thus may potentially be detected. Swallowing has been shown to be one of the most reliable indicators of DI, as any food requires swallowing to contribute to nutrition. Consumption of both solid and liquid foods manifests as an increase in swallowing frequency⁶⁴ over spontaneous non-nutritive swallowing. Swallowing sensors include microphones,⁶⁵ electrical sensors, or motion sensors.⁶⁶ The frequency of swallowing may be used to differentiate consumption of solids and liquids,⁶⁴ and the count of swallows per meal may serve as an indicator of the amount consumed.⁶⁷ In general, a significant strength of the sensor-based approaches is that in most (not all) of these, the food intake can be detected automatically, without self-report. However, the technology behind sensor devices is new and most have not been thoroughly tested and validated for use in community-dwelling individuals, and there is concern that wearing the device may cause some reactivity bias. Furthermore, sensor devices can only detect the total amount of food ingested and are unable to identify types of foods, portion sizes, nutrient composition, or energy intake.^{55,68}

Other sensor- and informatics-based research tools have been developed to determine food type and nutritional composition, for example, food classification-based acoustical sensing,⁶¹ use of miniaturized hand-held (near-infrared) spectrometers that can scan food items and determine characteristic food matrix properties,⁶⁹ or recent miniaturized tooth-mounted sensors capable of detecting nutrients and wirelessly communicating to a mobile device.⁷⁰ Such technologies are still under research and development at this time, and many require support of comprehensive nutrient databases to support the technology and methodology to assess portion size.

PHYSICAL ACTIVITY ASSESSMENT

Similarly to DI, the assessment of PA can be measured through self-report or device-based techniques.^{18,19} Researcher-selected PA methods and tools are influenced by cost, participant burden, sample size, collection time frame, type of information required (e.g., steps, counts, energy expenditure [EE]), data management, and measurement error.^{71,72} The following section provides a brief overview of PA assessment tools. Additional information can be found in recent reviews by Ainsworth and colleagues,¹⁸ Sylvia et al.,⁷³ Welk and colleagues,⁷⁴ and the Physical Activity Resource Center for Public Health (www.parcph.org/). PA assessment techniques are summarized in [Table 2](#).

Device-Based Physical Activity Assessment Tools

Research grade devices. Triaxial accelerometers, such as the ActiGraph wGT3X-BT, measure PA volume and intensity. They are commonly worn on the wrist or hip, with the hip location providing better accuracy.^{80–83} The major strength of accelerometers is their ability to collect large amounts of data and measure intensity level. Limitations include expense and the inability to provide contextual information. Furthermore, data collection protocols (e.g., hip versus wrist placement, waking-hour versus 24-hour registration period) and data analysis approaches (e.g., non–wear-time definition, cutpoints for intensity classification) vary, making it very difficult to compare across studies.⁸⁴ Researchers have traditionally used “activity counts” to classify PA as light, moderate, or vigorous intensity, but the field is shifting to activity characterization from raw acceleration signals.^{85,86} Lastly, researchers are working to improve the ability to analyze data from wrist-worn devices,^{83,87,88} which may improve compliance.⁸⁵ For an extensive discussion of considerations when using accelerometers, see the 2017 review by Migueles et al.⁸⁴

The activPAL is a particularly useful device for researchers interested in sedentary behavior.^{108,109} The

activPAL is affixed to the thigh, which makes it uniquely capable of assessing postures (e.g., sitting versus standing). The device also measures step cadence and number of steps, therefore allowing activity to be classified as sitting, standing, or stepping. There is promising evidence that the activPAL can also accurately classify PA intensities.^{110,111} For an extensive discussion of considerations when using the activPAL, see the review by Edwardson and colleagues.⁸⁹

HR monitors are used in laboratory settings to assess exercise activity, intensity of the activity, and EE of activity^{90,91} due to the direct and linear relationship between HR and oxygen consumption.⁹² Recently, HR monitoring has been combined with accelerometry to more accurately account for the predictive power of HR at rest and during light activity in EE estimates.⁹³

GPS units enable collection of altitude, longitude, latitude, speed, distance traveled, and elevation data.^{94,95} Commercial GPS units can be accurate up to 15–20 meters; however, the clarity of the device signal to the satellites is crucial, affecting sample rates and signal validity.^{96,97} Compared with accelerometers, GPS units significantly underestimate PA (i.e., EE).⁹⁸ There have been suggestions to use GPS in combination with HR monitors and accelerometry.^{97,99}

Multisensor devices utilize multiple physiological and mechanical sensors in combination to improve precision of PA and EE measurements. For example, the SenseWear armband is worn on the upper arm and incorporates triaxial accelerometry, heat flux, galvanic skin response, skin temperature, and near-body ambient temperature to accurately determine when the device is being worn (i.e., a major consideration with traditional accelerometers).¹⁰⁰ These measures, in combination with entered data, enable accurate estimation of EE, minutes of activity, and sleep.^{101–105} However, Jawbone Inc. acquired BodyMedia in 2013 and discontinued support of the SenseWear armband. Thus, this device is no longer available for purchase. Another device, the Intelligent Device for Energy Expenditure and Activity, incorporates five sensors (chest, right and left thighs, and right and left legs) connected to a digital recorder that allows identification of 32 different activities and body postures for estimation of PA level and accurate EE.^{106,107}

General-Use Devices

Pedometers are simple devices that measure steps. They are inexpensive and useful in assessing behavioral feedback and motivation.¹¹⁸ Pedometer accuracy has improved with transition into microelectromechanical-based systems¹¹² specifically with measurements more than 2 mph.^{112–115} Pedometer output can vary

Table 2. Summary of Device-Based PA Assessment Tools

Method/tool	Appropriate populations	Outcome measure	Attributes	Limitations	Validity	Research gaps	References
General-use wearables (Fitbit, Garmin, Apple)	Large population; behavior change within individuals	EE	Popular; ease of collection and data upload (wireless); large amounts of data collected	Not a valid measure of TEE; underestimates free-living EE; overestimates PAEE; algorithms change with updates; not designed for research; cost to obtain data	Correlation between consumer activity monitors and accelerometers for sleep count and step count, $r > 0.8$; for TDEE, $r = 0.74$ to 0.81 ; and MVPA, $r = 0.52$ to 0.91	More accuracy research is needed	75–79
Accelerometers	Large populations	Minutes of physical activity, intensity	Commonly used in research settings and by NHANES; ability to collect large amounts of data	Expense; inability to provide contextual information; data collection protocols (e.g., hip vs wrist placement, waking-hour vs 24-hour registration period) and data analysis approaches (e.g., non-wear-time definition, valid day criteria, cutpoints for intensity classification) vary, making it very difficult to compare across studies using accelerometry	Correlations between daily PAEE and activity counts for the hip-worn ActiGraph range from $r = 0.77$ to 0.90 ; compared to wearable cameras measuring PA, hip-worn accelerometers had 89.4% accuracy and wrist-worn accelerometers had 84.6% accuracy	Lack of consensus regarding data processing	80–88
GPS	Large populations; outdoors	Distance and speed	Ideal use outdoors (free-living walking and running) or field testing	Underestimates EE for field activities; not a standalone measure for EE; not appropriate for indoor activities; issues with battery life	Compared to accelerometers, GPS underestimates EE by 42% to 50%	Stronger association with EE measure (accelerometer use)	89–95
HR monitor	Supervised exercise	HR, activity intensity	Direct measure, high validity to clinical measures	Uncomfortable when worn for long periods of time; not a valid estimate of EE at rest; must have an HR-O ₂ consumption curve for each person to measure their intensity; TDEE is hard to predict because daily HR is not linear	During PA, EE error rates are <3% compared to whole-room calorimetry; however, when doing light or sedentary activity, they have poor predictive power in terms of EE	Improved estimates of TEE	96–99

(continued on next page)

Table 2. Summary of Device-Based PA Assessment Tools (continued)

Method/tool	Appropriate populations	Outcome measure	Attributes	Limitations	Validity	Research gaps	References
Multisensor	Populations with a wide range of activities	Minutes of PA, EE	Multiple mechanical and physiological sensors improve estimates	Cost, availability	SenseWear: EE and activity ICC of 0.81–0.85 compared to DLW; IDEEA: EE within 98.9%±9.0% compared to indirect calorimetry	Improved estimates of individual-level estimates	100–107
ActivPAL	Large populations	Sedentary time, steps/day	Can be worn 24 hours/day	Relatively small body of literature compared to other accelerometers	Correlations range from 0.78 to 0.99 against direct observation	Validation for EE	88, 108–111
Pedometers	Large populations	Steps/day	Affordable; best used to assess walking; steps/day is well understood by the lay population; newer versions store data	Inaccurate at slow speeds; inter-individual variability-based difference; in some models, must manually record steps; readings vary according to anatomical location (e.g., hip or ankle)	Varies by model	Estimations of EE and exercise intensity	71–74, 80, 112–117

DLW, doubly labeled water; EE, energy expenditure; HR, heart rate; ICC, intraclass correlation coefficient; IDEEA, Intelligent Device for Energy Expenditure and Activity; MVPA, moderate to vigorous physical activity; NHANES, National Health and Nutrition Examination Survey; PA, physical activity; PAEE, physical activity energy expenditure; TDEE, total daily energy expenditure; TEE, total energy expenditure.

according to location worn (e.g., the ankle is the most accurate placement)^{114,116} or among individuals (e.g., foot-strike variability).¹¹⁷

Recently, commercial off the shelf (COTS) activity trackers from Fitbit, Garmin, and Apple have exploded onto the consumer market.⁷⁵ These newer devices use advanced technologies allowing expansion of monitoring capabilities (e.g., accelerations, HR, EE, and sleep) and are able to transmit and store PA data to smartphones, computers, and cloud-based storage. New devices and algorithm updates are released frequently with expanded capability to detect posture changes and type of activity for more accurate and precise estimates of EE. These wearables provide health data that are instantly available to the consumer through a smartphone application. Such wearable devices employ multiple engagement strategies to make them more attractive and interactive for the individual.⁷⁶

Popularity in the marketplace has led to more research in the past 3–5 years to validate accuracy and reliability of EE for COTS wearables compared with more traditional measures, such as the ActiGraph.^{76–78,111} Compared with EE measured by doubly labeled water, COTS wearables underestimated EE in free-living, normal-weight men and women aged 21–50 years.⁷⁹

Other limitations to using COTS wearables for EE assessment in research include the lack of transparency of cutpoint data and algorithms used to calculate activity intensity and EE. Data management can also become overwhelmingly expensive, and many companies employ a third party to clean and organize the data. COTS wearables were not developed to be research grade; therefore, inclusion of third-party sites for data management makes it difficult for researchers to obtain required data necessary for analysis.

CONCLUSIONS

Accurate measurement of DI and PA is needed for both population- and intervention-based assessments. Although there are many limitations to the measurement of DI and PA, there is progress and promise for using technology to improve these measures. Managing the current knowledge base and facilitating a resource center for new technology integration are key to the future success of accurate DI and PA measures through device-assisted methods.

ACKNOWLEDGMENTS

The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the Army or the Department of Defense, U.S. Department of Agriculture, American College of Sports Medicine

(ACSM), or the North American Branch of the International Life Sciences Institute (ILSI North America). Any citations of commercial organizations and trade names in this report do not constitute an official endorsement or approval of the products or services of these organizations.

This review was inspired by discussions at a 2016 expert forum (Tech Summit: Innovative Tools for Assessing Diet and Physical Activity for Health Promotion), which was organized by ILSI North America with the help of scientists from the University of California, San Diego (UCSD), the U.S. Department of Agriculture Agricultural Research Service, ACSM, and NIH. In-kind support was received from UCSD and financial contributions were provided by ACSM and the ILSI North America Committee on Balancing Food and Activity for Health. Additional funding from the ILSI North America Committee on Balancing Food and Activity for Health was allocated for the preparation and publication of papers following the forum. ILSI North America is a public, non-profit foundation that provides a forum to advance understanding of scientific issues related to the nutritional quality and safety of the food supply by sponsoring research programs, educational seminars and workshops, and publications. ILSI North America receives support primarily from its industry membership. The opinions expressed herein are those of the authors and do not necessarily represent the views of the funding organization. Lauren T. Ptomey received an honorarium from ILSI North America. This article has been reviewed and approved for submission to the *American Journal of Preventive Medicine* by ILSI North America. ILSI North America had no role in the study design; collection, analysis, and interpretation of data; writing the report; and the decision to submit the report for publication.

HLM, LTP, AA, ES, RW, and SKD composed and drafted the dietary intake information and table. RPS, AMG, and KB prepared and drafted the physical activity information and table. HLM, LTP, and SKD edited the manuscript. All authors read and approved the final manuscript.

No financial disclosures were reported by the authors of this paper.

THEME NOTE

This article is part of a theme issue entitled *Innovative Tools for Assessing Diet and Physical Activity for Health Promotion*, which is sponsored by the North American branch of the International Life Sciences Institute.

REFERENCES

- Chen M, Sun Q, Giovannucci E, et al. Dairy consumption and risk of type 2 diabetes: 3 cohorts of U.S. adults and an updated meta-analysis. *BMC Med*. 2014;12:215. <https://doi.org/10.1186/s12916-014-0215-1>.
- O'Neil CE, Nicklas TA, Fulgoni 3rd VL. Nutrient intake, diet quality, and weight/adiposity parameters in breakfast patterns compared with no breakfast in adults: National Health and Nutrition Examination Survey 2001–2008. *J Acad Nutr Diet*. 2014;114(12 suppl):S27–S43. <https://doi.org/10.1016/j.jand.2014.08.021>.
- Ford ES, Dietz WH. Modeling dietary patterns to assess sodium recommendations for nutrient adequacy. *Am J Clin Nutr*. 2013;97(4):848–853. <https://doi.org/10.3945/ajcn.112.052662>.
- Subar AF, Kipnis V, Troiano RP, et al. Using intake biomarkers to evaluate the extent of dietary misreporting in a large sample of adults: the OPEN study. *Am J Epidemiol*. 2003;158(1):1–13. <https://doi.org/10.1093/aje/kwg092>.
- Trabulsi J, Schoeller DA. Evaluation of dietary assessment instruments against doubly labeled water, a biomarker of habitual energy intake. *Am J Physiol Endocrinol Metab*. 2001;281(5):E891–E899. <https://doi.org/10.1152/ajpendo.2001.281.5.E891>.
- Freedman LS, Commins JM, Moler JE, et al. Pooled results from 5 validation studies of dietary self-report instruments using recovery biomarkers for energy and protein intake. *Am J Epidemiol*. 2014;180(2):172–188. <https://doi.org/10.1093/aje/kwu116>.
- Park Y, Dodd KW, Kipnis V, et al. Comparison of self-reported dietary intakes from the Automated Self-Administered 24-h recall, 4-d food records, and food-frequency questionnaires against recovery biomarkers. *Am J Clin Nutr*. 2018;107(1):80–93. <https://doi.org/10.1093/ajcn/nqx002>.
- Schoeller DA, Thomas D, Archer E, et al. Self-report-based estimates of energy intake offer an inadequate basis for scientific conclusions. *Am J Clin Nutr*. 2013;97(6):1413–1415. <https://doi.org/10.3945/ajcn.113.062125>.
- Dhurandhar N, Schoeller D, Brown A, et al. Energy balance measurement: when something is not better than nothing. *Int J Obes*. 2015;39(7):1109–1113. <https://doi.org/10.1038/ijo.2014.199>.
- Archer E, Hand GA, Blair SN. Validity of U.S. nutritional surveillance: National Health and Nutrition Examination Survey caloric energy intake data, 1971–2010. *PLoS One*. 2013;8(10):e76632. <https://doi.org/10.1371/journal.pone.0076632>.
- Archer E. The use of implausible data without caveats is misleading. *Am J Clin Nutr*. 2017;106(3):949–950. <https://doi.org/10.3945/ajcn.116.150870>.
- Subar AF, Freedman LS, Tooze JA, et al. Addressing current criticism regarding the value of self-report dietary data. *J Nutr*. 2015;145(12):2639–2645. <https://doi.org/10.3945/jn.115.219634>.
- Schoeller DA, Bandini LG, Dietz WH. Inaccuracies in self-reported intake identified by comparison with the doubly labelled water method. *Can J Physiol Pharmacol*. 1990;68(7):941–949. <https://doi.org/10.1139/y90-143>.
- Prentice R, Huang Y, Neuhauser M, et al. Associations of biomarker-calibrated sodium and potassium intakes with cardiovascular disease risk among postmenopausal women. *Am J Epidemiol*. 2017;186(9):1035–1043. <https://doi.org/10.1093/aje/kwx238>.
- Dong Y, Scisco J, Wilson M, Muth E, Hoover A. Detecting periods of eating during free-living by tracking wrist motion. *IEEE J Biomed Health Inform*. 2014;18(4):1253–1260. <https://doi.org/10.1109/JBHI.2013.2282471>.
- Fontana JM, Farooq M, Sazonov E. Automatic ingestion monitor: a novel wearable device for monitoring of ingestive behavior. *IEEE Trans Biomed Eng*. 2014;61(6):1772–1779. <https://doi.org/10.1109/TBME.2014.2306773>.
- Martin C, Nicklas T, Gunturk B, Correa J, Allen H, Champagne C. Measuring food intake with digital photography. *J Hum Nutr Diet*. 2014;27(s1):72–81. <https://doi.org/10.1111/jhn.12014>.
- Ainsworth B, Cahalin L, Buman M, Ross R. The current state of physical activity assessment tools. *Prog Cardiovasc Dis*. 2015;57(4):387–395. <https://doi.org/10.1016/j.pcad.2014.10.005>.
- Bassett Jr DR. Validity and reliability issues in objective monitoring of physical activity. *Res Q Exerc Sport*. 2000;71(2 suppl):30–36. <https://doi.org/10.1080/02701367.2000.11082783>.
- Norman GJ, Zabinski MF, Adams MA, Rosenberg DE, Yaroch AL, Atienza AA. A review of eHealth interventions for physical activity and dietary behavior change. *Am J Prev Med*. 2007;33(4):336–345. <https://doi.org/10.1016/j.amepre.2007.05.007>.
- Atkinson NL, Gold RS. The promise and challenge of eHealth interventions. *Am J Health Behav*. 2002;26(6):494–503. <https://doi.org/10.5993/AJHB.26.6.10>.

22. Kroeze W, Werkman A, Brug J. A systematic review of randomized trials on the effectiveness of computer-tailored education on physical activity and dietary behaviors. *Ann Behav Med*. 2006;31(3):205–223. https://doi.org/10.1207/s15324796abm3103_2.
23. Brug J, Oenema A, Campbell M. Past, present, and future of computer-tailored nutrition education. *Am J Clin Nutr*. 2003;77(4 suppl):1028s–1034s. <https://doi.org/10.1093/ajcn/77.4.1028S>.
24. Eng TR. *The eHealth Landscape: A Terrain Map of Emerging Information and Communication Technologies in Health and Health Care*. Princeton, NJ: Robert Wood Johnson Foundation, 2001.
25. Farshchi H, Macdonald I, Madjid A, Taylor M. Benefits and limitations of traditional self-report instruments. In: Schoeller D, editor. *Advances in the Assessment of Dietary Intake*. Boca Raton, FL: Taylor and Francis, 2017:1–17. <https://doi.org/10.1201/9781315152288-2>.
26. Shim J, Oh K, Kim J. Dietary assessment methods in epidemiologic studies. *Epidemiol Health*. 2014;36:e2014009. <https://doi.org/10.4178/epih/e2014009>.
27. Kirkpatrick SI, Subar AF, Toozé JA. Statistical approaches to mitigate measurement error in dietary intake data collected using 24-hour recalls, and food records/diaries. In: Schoeller D, editor. *Advances in the Assessment of Dietary Intake*. Boca Raton, FL: Taylor and Francis, 2017:19–43. <https://doi.org/10.1201/9781315152288-3>.
28. National Cancer Institute. Dietary Assessment Primer. <https://dietassessmentprimer.cancer.gov/>. Accessed June 1, 2018.
29. Moshfegh AJ, Rhodes DG, Baer DJ, et al. The U.S. Department of Agriculture Automated Multiple-Pass Method reduces bias in the collection of energy intakes. *Am J Clin Nutr*. 2008;88(2):324–332. <https://doi.org/10.1093/ajcn/88.2.324>.
30. Schatzkin A, Subar AF, Moore S, et al. Observational epidemiologic studies of nutrition and cancer: the next generation (with better observation). *Cancer Epidemiol Biomarkers Prev*. 2009;18(4):1026–1032. <https://doi.org/10.1158/1055-9965.EPI-08-1129>.
31. Williamson DA, Allen HR, Martin PD, Alfonso AJ, Gerald B, Hunt A. Comparison of digital photography to weighed and visual estimation of portion sizes. *J Am Diet Assoc*. 2003;103(9):1139–1145. [https://doi.org/10.1016/S0002-8223\(03\)00974-X](https://doi.org/10.1016/S0002-8223(03)00974-X).
32. Crombie AP, Funderburk LK, Smith TJ, et al. Effects of modified foodservice practices in military dining facilities on ad libitum nutritional intake of U.S. Army soldiers. *J Acad Nutr Diet*. 2013;113(7):920–927. <https://doi.org/10.1016/j.jand.2013.01.005>.
33. Wang DH, Kogashiwa M, Kira S. Development of a new instrument for evaluating individuals' dietary intakes. *J Am Diet Assoc*. 2006;106(10):1588–1593. <https://doi.org/10.1016/j.jada.2006.07.004>.
34. Ptomey LT, Willis EA, Honas JJ, et al. Validity of energy intake estimated by digital photography plus recall in overweight and obese young adults. *J Acad Nutr Diet*. 2015;115(9):1392–1399. <https://doi.org/10.1016/j.jand.2015.05.006>.
35. Elinder L, Brunosson A, Bergström H, Hagström M, Patterson E. Validation of personal digital photography to assess dietary quality among people with intellectual disabilities. *J Intellect Disabil Res*. 2012;56(2):221–226. <https://doi.org/10.1111/j.1365-2788.2011.01459.x>.
36. Ptomey LT, Herrmann SD, Lee J, Sullivan DK, Rondon MF, Donnelly JE. Photo-assisted recall increases estimates of energy and macronutrient intake in adults with intellectual and developmental disabilities. *J Acad Nutr Diet*. 2013;113(12):1704–1709. <https://doi.org/10.1016/j.jand.2013.07.029>.
37. Martin CK, Correa JB, Han H, et al. Validity of the Remote Food Photography Method (RFPM) for estimating energy and nutrient intake in near real-time. *Obesity (Silver Spring)*. 2012;20(4):891–899. <https://doi.org/10.1038/oby.2011.344>.
38. McClung H, Champagne C, Allen H, et al. Digital food photography technology improves efficiency and feasibility of dietary intake assessments in large populations eating ad libitum in collective dining facilities. *Appetite*. 2017;116:389–394. <https://doi.org/10.1016/j.appet.2017.05.025>.
39. Boushey C, Spoden M, Zhu F, Delp E, Kerr D. New mobile methods for dietary assessment: review of image-assisted and image-based dietary assessment methods. *Proc Nutr Soc*. 2017;76(3):283–294. <https://doi.org/10.1017/S0029665116002913>.
40. Rollo ME, Ash S, Lyons-Wall P, Russell AW. Evaluation of a mobile phone image-based dietary assessment method in adults with type 2 diabetes. *Nutrients*. 2015;7(6):4897–4910. <https://doi.org/10.3390/nu7064897>.
41. Jia W, Chen H-C, Yue Y, et al. Accuracy of food portion size estimation from digital pictures acquired by a chest-worn camera. *Public Health Nutr*. 2014;17(8):1671–1681. <https://doi.org/10.1017/S1368980013003236>.
42. Six BL, Schap TE, Zhu FM, et al. Evidence-based development of a mobile telephone food record. *J Am Diet Assoc*. 2010;110(1):74–79. <https://doi.org/10.1016/j.jada.2009.10.010>.
43. Zhu F, Bosch M, Woo I, et al. The use of mobile devices in aiding dietary assessment and evaluation. *IEEE J Sel Top Signal Process*. 2010;4(4):756–766. <https://doi.org/10.1109/JSTSP.2010.2051471>.
44. Zhu F, Bosch M, Khanna N, Boushey CJ, Delp EJ. Multiple hypotheses image segmentation and classification with application to dietary assessment. *IEEE J Biomed Health Inform*. 2015;19(1):377–388. <https://doi.org/10.1109/JBHI.2014.2304925>.
45. Weiss R, Stumbo PJ, Divakaran A. Automatic food documentation and volume computation using digital imaging and electronic transmission. *J Am Diet Assoc*. 2010;110(1):42–44. <https://doi.org/10.1016/j.jada.2009.10.011>.
46. National Cancer Institute. Automated Self-Administered 24-Hour (ASA24[®]) Dietary Assessment Tool. <https://epi.grants.cancer.gov/asa24/>. Accessed June 21, 2018.
47. National Cancer Institute. Comparison Among ASA24[®] Versions. <https://epi.grants.cancer.gov/asa24/comparison.html>. Accessed November 15, 2017.
48. Agricultural Research Service. AMPM-USDA Automated Multiple-Pass Method. www.ars.usda.gov/northeast-area/beltsville-md-bhnrc/beltsville-human-nutrition-research-center/food-surveys-research-group/docs/ampm-usda-automated-multiple-pass-method/. Updated 2016. Accessed November 15, 2017.
49. Subar AF, Thompson FE, Smith AF, et al. Improving food frequency questionnaires: a qualitative approach using cognitive interviewing. *J Am Diet Assoc*. 1995;95(7):781–788. [https://doi.org/10.1016/S0002-8223\(95\)00217-0](https://doi.org/10.1016/S0002-8223(95)00217-0).
50. NutritionQuest. <http://nutritionquest.com/>. Accessed November 15, 2017.
51. National Cancer Institute. Diet History Questionnaire (DHQ) III. <https://epi.grants.cancer.gov/dhq3/>. Accessed June 11, 2018.
52. Fred Hutchinson Cancer Research Center. Food Frequency Questionnaires (FFQ). <http://sharedresources.fredhutch.org/services/food-frequency-questionnaires-ffq>. Accessed November 15, 2017.
53. Kristal AR, Kolar AS, Fisher JL, et al. Evaluation of web-based, self-administered, graphical food frequency questionnaire. *J Acad Nutr Diet*. 2014;114(4):613–621. <https://doi.org/10.1016/j.jand.2013.11.017>.
54. VioScreen[™]. www.viocare.com/vioscreen.html. Accessed November 15, 2017.
55. McClung HL, Kehayias JJ, Zientara GP, Hoyt R. Direct and indirect measures of dietary intake: use of sensors and modern technologies. In: Schoeller D, ed. *Advances in the Assessment of Dietary Intake*. Boca Raton, FL: Taylor and Francis; 2017:164–183. <https://doi.org/10.1201/9781315152288-10>.
56. Chang J, Lui S, Chu H. The diet-aware dining table: observing dietary behaviors over a tabletop surface. In: Fishkin KP, Schiele B, Nixon P, Quigley A, eds. *Pervasive Computing. Pervasive 2006. Lecture Notes in Computer Science*. Berlin, Heidelberg: Springer, 2006:3968. https://doi.org/10.1007/11748625_23.
57. Stevens J, Bryant M, Wang L, Borja J, Bentley ME. Exhaustive measurement of food items in the home using a universal product code

- scanner. *Public Health Nutr.* 2011;14(2):314–318. <https://doi.org/10.1017/S1368980010001837>.
58. French SA, Shimotsu ST, Wall M, Gerlach AF. Capturing the spectrum of household food and beverage purchasing behavior: a review. *J Am Diet Assoc.* 2008;108(12):2051–2058. <https://doi.org/10.1016/j.jada.2008.09.001>.
 59. Dong Y, Hoover A, Scisco J, Muth E. A new method for measuring meal intake in humans via automated wrist motion tracking. *Appl Psychophysiol Biofeedback.* 2012;37(3):205–215. <https://doi.org/10.1007/s10484-012-9194-1>.
 60. Scisco JL, Muth ER, Hoover AW. Examining the utility of a bite-count-based measure of eating activity in free-living human beings. *J Acad Nutr Diet.* 2014;114(3):464–469. <https://doi.org/10.1016/j.jand.2013.09.017>.
 61. Passler S, Wolff M, Fischer WJ. Food intake monitoring: an acoustical approach to automated food intake activity detection and classification of consumed food. *Physiol Meas.* 2012;33(6):1073–1093. <https://doi.org/10.1088/0967-3334/33/6/1073>.
 62. Zhang R, Bernhart S, Amft O. Diet eyeglasses: recognising food chewing using EMG and smart eyeglasses. 2016 IEEE 13th International Conference on Wearable and Implantable Body Sensor Networks (BSN). <https://doi.org/10.1109/BSN.2016.7516224>.
 63. Sazonov ES, Fontana JM. A Sensor system for automatic detection of food intake through non-invasive monitoring of chewing. *IEEE Sens J.* 2012;12(5):1340–1348. <https://doi.org/10.1109/JSEN.2011.2172411>.
 64. Sazonov ES, Schuckers SA, Lopez-Meyer P, et al. Toward objective monitoring of ingestive behavior in free-living population. *Obesity (Silver Spring).* 2009;17(10):1971–1975. <https://doi.org/10.1038/oby.2009.153>.
 65. Sazonov E, Schuckers S, Lopez-Meyer P, et al. Non-invasive monitoring of chewing and swallowing for objective quantification of ingestive behavior. *Physiol Meas.* 2008;29(5):525–541. <https://doi.org/10.1088/0967-3334/29/5/001>.
 66. Kalantarian H, Alshurafa N, Le T, Sarrafzadeh M. Monitoring eating habits using a piezoelectric sensor-based necklace. *Comput Biol Med.* 2015;58:46–55. <https://doi.org/10.1016/j.compbiomed.2015.01.005>.
 67. Fontana JM, Higgins JA, Schuckers SC, et al. Energy intake estimation from counts of chews and swallows. *Appetite.* 2015;85:14–21. <https://doi.org/10.1016/j.appet.2014.11.003>.
 68. Stumbo PJ. New technology in dietary assessment: a review of digital methods in improving food record accuracy. *Proc Nutr Soc.* 2013;72(1):70–76. <https://doi.org/10.1017/S0029665112002911>.
 69. Lieberman HR, Caruso CM, Niro PJ, et al. A double-blind, placebo-controlled test of 2 d of calorie deprivation: effects on cognition, activity, sleep, and interstitial glucose concentrations. *Am J Clin Nutr.* 2008;88(3):667–676. <https://doi.org/10.1093/ajcn/88.3.667>.
 70. Tseng P, Napier B, Garbarin IL, Kaplan D, Omenetto F. Functional, RF-trilayer sensors for tooth-mounted, wireless monitoring of the oral cavity and food consumption. *Adv Mater.* 2018;30(18):1703257. <https://doi.org/10.1002/adma.201703257>.
 71. Schutz Y, Weinsier RL, Hunter GR. Assessment of free living physical activity in humans: an overview of currently available and proposed new measures. *Obes Res.* 2001;9(6):368–379. <https://doi.org/10.1038/oby.2001.48>.
 72. Hills AP, Mokhtar N, Byrne NM. Assessment of physical activity and energy expenditure: an overview of objective measures. *Front Nutr.* 2014;1:5. <https://doi.org/10.3389/fnut.2014.00005>.
 73. Sylvia LG, Bernstein EE, Hubbard JL, Keating L, Anderson EJ. A practical guide to measuring physical activity. *J Acad Nutr Diet.* 2014;114(2):199–208. <https://doi.org/10.1016/j.jand.2013.09.018>.
 74. Welk GJ, Morrow J, Saint-Maurice P. Measures Registry User Guide: Individual Physical Activity. www.nccor.org/wp-content/uploads/sites/2/2017/NCCOR_MR_User_Guide_Individual_PA-FINAL.pdf. Accessed June 30, 2018. Published 2017.
 75. Morabito V. Wearable technologies. In: *The Future of Digital Business Innovation*. New York, NY: Springer; 2016:23–42. https://doi.org/10.1007/978-3-319-26874-3_2.
 76. Case MA, Burwick HA, Volpp KG, Patel MS. Accuracy of smart-phone applications and wearable devices for tracking physical activity data. *JAMA.* 2015;313(6):625–626. <https://doi.org/10.1001/jama.2014.17841>.
 77. Nelson MB, Kaminsky LA, Dickin DC, Montoye A. Validity of consumer-based physical activity monitors for specific activity types. *Med Sci Sports Exerc.* 2016;48(8):1619–1628. <https://doi.org/10.1249/MSS.0000000000000933>.
 78. Ferguson T, Rowlands AV, Olds T, Maher C. The validity of consumer-level, activity monitors in healthy adults worn in free-living conditions: a cross-sectional study. *Int J Behav Nutr Phys Act.* 2015;12(1):42. <https://doi.org/10.1186/s12966-015-0201-9>.
 79. Murakami H, Kawakami R, Nakae S, et al. Accuracy of wearable devices for estimating total energy expenditure: comparison with metabolic chamber and doubly labeled water method. *JAMA Intern Med.* 2016;176(5):702–703. <https://doi.org/10.1001/jamainternmed.2016.0152>.
 80. Tudor-Locke C, Barreira TV, Schuna Jr JM. Comparison of step outputs for waist and wrist accelerometer attachment sites. *Med Sci Sports Exerc.* 2015;47(4):839–842. <https://doi.org/10.1249/MSS.0000000000000476>.
 81. Ellis K, Kerr J, Godbole S, Staudenmayer J, Lanckriet G. Hip and wrist accelerometer algorithms for free-living behavior classification. *Med Sci Sports Exerc.* 2016;48(5):933–940. <https://doi.org/10.1249/MSS.0000000000000840>.
 82. Ellis K, Kerr J, Godbole S, Lanckriet G, Wing D, Marshall S. A random forest classifier for the prediction of energy expenditure and type of physical activity from wrist and hip accelerometers. *Physiol Meas.* 2014;35(11):2191–2203. <https://doi.org/10.1088/0967-3334/35/11/2191>.
 83. Zhang S, Rowlands AV, Murray P, Hurst TL. Physical activity classification using the GENE wrist-worn accelerometer. *Med Sci Sports Exerc.* 2012;44(4):742–748. <https://doi.org/10.1249/MSS.0b013e31823bf95c>.
 84. Migueles JH, Cadenas-Sanchez C, Ekelund U, et al. Accelerometer data collection and processing criteria to assess physical activity and other outcomes: a systematic review and practical considerations. *Sports Med.* 2017;47(9):1821–1845. <https://doi.org/10.1007/s40279-017-0716-0>.
 85. Troiano RP, McClain JJ, Brychta RJ, Chen KY. Evolution of accelerometer methods for physical activity research. *Br J Sports Med.* 2014;48(13):1019–1023. <https://doi.org/10.1136/bjsports-2014-093546>.
 86. Liu S, Gao RX, Freedson PS. Computational methods for estimating energy expenditure in human physical activities. *Med Sci Sports Exerc.* 2012;44(11):2138–2146. <https://doi.org/10.1249/MSS.0b013e31825e825a>.
 87. Staudenmayer J, He S, Hickey A, Sasaki J, Freedson P. Methods to estimate aspects of physical activity and sedentary behavior from high-frequency wrist accelerometer measurements. *J Appl Physiol (1985).* 2015;119(4):396–403. <https://doi.org/10.1152/jappphysiol.00026.2015>.
 88. Hildebrand M, VAN Hees VT, Hansen BH, Ekelund U. Age group comparability of raw accelerometer output from wrist- and hip-worn monitors. *Med Sci Sports Exerc.* 2014;46(9):1816–1824. <https://doi.org/10.1249/MSS.0000000000000289>.
 89. Edwardson CL, Winkler EAH, Bodicoat DH, et al. Considerations when using the activPAL monitor in field-based research with adult populations. *J Sport Health Sci.* 2017;6(2):162–178. <https://doi.org/10.1016/j.jshs.2016.02.002>.
 90. Church TS, Blair SN, Cocroham S, et al. Effects of aerobic and resistance training on hemoglobin A1c levels in patients with type 2

- diabetes: a randomized controlled trial. *JAMA*. 2010;304(20):2253–2262. <https://doi.org/10.1001/jama.2010.1710>.
91. Church TS, Earnest CP, Skinner JS, Blair SN. Effects of different doses of physical activity on cardiorespiratory fitness among sedentary, overweight or obese postmenopausal women with elevated blood pressure: a randomized controlled trial. *JAMA*. 2007;297(19):2081–2091. <https://doi.org/10.1001/jama.297.19.2081>.
 92. Freedson PS, Miller K. Objective monitoring of physical activity using motion sensors and heart rate. *Res Q Exerc Sport*. 2000;71(suppl 2):21–29. <https://doi.org/10.1080/02701367.2000.11082782>.
 93. Dong L, Block G, Mandel S. Activities contributing to total energy expenditure in the United States: results from the NHAPS Study. *Int J Behav Nutr Phys Act*. 2004;1(1):4. <https://doi.org/10.1186/1479-5868-1-4>.
 94. Terrier P, Schutz Y. How useful is satellite positioning system (GPS) to track gait parameters? A review. *J Neuroeng Rehabil*. 2005;2(1):28. <https://doi.org/10.1186/1743-0003-2-28>.
 95. Witte T, Wilson A. Accuracy of WAAS-enabled GPS for the determination of position and speed over ground. *J Biomech*. 2005;38(8):1717–1722. <https://doi.org/10.1016/j.jbiomech.2004.07.028>.
 96. Wieters KM, Kim J, Lee C. Assessment of wearable global positioning system units for physical activity research. *J Phys Act Health*. 2012;9(7):913–923. <https://doi.org/10.1123/jpah.9.7.913>.
 97. Stopher P, FitzGerald C, Zhang J. Search for a global positioning system device to measure person travel. *Trans Res Part C Emerg Technol*. 2008;16(3):350–369. <https://doi.org/10.1016/j.trc.2007.10.002>.
 98. Hongu N, Orr BJ, Roe DJ, Reed RG, Going SB. Global positioning system watches for estimating energy expenditure. *J Strength Cond Res*. 2013;27(11):3216–3220. <https://doi.org/10.1519/JSC.0b013e31828bae0f>.
 99. Rodriguez DA, Brown AL, Troped PJ. Portable global positioning units to complement accelerometry-based physical activity monitors. *Med Sci Sports Exerc*. 2005;37(11):S572. <https://doi.org/10.1249/01.mss.0000185297.72328.ce>.
 100. Herrmann SD, Barreira TV, Kang M, Ainsworth BE. Impact of accelerometer wear time on physical activity data: a NHANES semisimulation data approach. *Br J Sports Med*. 2014;48(3):278–282. <https://doi.org/10.1136/bjsports-2012-091410>.
 101. Jakicic JM, Marcus M, Gallagher K, et al. Evaluation of the SenseWear Pro Armband™ to assess energy expenditure during exercise. *Med Sci Sports Exerc*. 2004;36(5):897–904. <https://doi.org/10.1249/01.MSS.0000126805.32659.43>.
 102. Johannsen DL, Calabro MA, Stewart J, Franke W, Rood JC, Welk GJ. Accuracy of armband monitors for measuring daily energy expenditure in healthy adults. *Med Sci Sports Exerc*. 2010;42(11):2134–2140. <https://doi.org/10.1249/MSS.0b013e3181e0b3ff>.
 103. St-Onge M, Mignault D, Allison DB, Rabasa-Lhoret R. Evaluation of a portable device to measure daily energy expenditure in free-living adults. *Am J Clin Nutr*. 2007;85(3):742–749. <https://doi.org/10.1093/ajcn/85.3.742>.
 104. Welk GJ, McClain JJ, Eisenmann JC, Wickel EE. Field validation of the MTI Actigraph and BodyMedia armband monitor using the IDEEA monitor. *Obesity*. 2007;15(4):918–928. <https://doi.org/10.1038/oby.2007.624>.
 105. Correa JB, Apolzan JW, Shepard DN, Heil DP, Rood JC, Martin CK. Evaluation of the ability of three physical activity monitors to predict weight change and estimate energy expenditure. *Appl Physiol Nutr Metab*. 2016;41(7):758–766. <https://doi.org/10.1139/apnm-2015-0461>.
 106. Zhang K, Pi-Sunyer FX, Boozer CN. Improving energy expenditure estimation for physical activity. *Med Sci Sports Exerc*. 2004;36(5):883–889. <https://doi.org/10.1249/01.MSS.0000126585.40962.22>.
 107. Zhang K, Werner P, Sun M, Pi-Sunyer FX, Boozer CN. Measurement of human daily physical activity. *Obes Res*. 2003;11(1):33–40. <https://doi.org/10.1038/oby.2003.7>.
 108. Kozey-Keadle S, Libertine A, Lyden K, Staudenmayer J, Freedson PS. Validation of wearable monitors for assessing sedentary behavior. *Med Sci Sports Exerc*. 2011;43(8):1561–1567. <https://doi.org/10.1249/MSS.0b013e31820ce174>.
 109. Bassett Jr DR, John D, Conger SA, Rider BC, Passmore RM, Clark JM. Detection of lying down, sitting, standing, and stepping using two activPAL monitors. *Med Sci Sports Exerc*. 2014;46(10):2025–2029. <https://doi.org/10.1249/MSS.0000000000000326>.
 110. Dowd KP, Harrington DM, Donnelly AE. Criterion and concurrent validity of the activPAL professional physical activity monitor in adolescent females. *PLoS One*. 2012;7(10):e47633. <https://doi.org/10.1371/journal.pone.0047633>.
 111. Lyden K, Keadle SK, Staudenmayer J, Freedson PS. The activPAL™ accurately classifies activity intensity categories in healthy adults. *Med Sci Sports Exerc*. 2017;49(5):1022–1028. <https://doi.org/10.1249/MSS.0000000000001177>.
 112. Berlin JE, Storti KL, Brach JS. Using activity monitors to measure physical activity in free-living conditions. *Phys Ther*. 2006;86(8):1137–1145.
 113. Crouter SE, Schneider PL, Karabulut M, Bassett DR. Validity of ten electronic pedometers for measuring steps, distance, and kcals. *Med Sci Sports Exerc*. 2003;35(5):S283. <https://doi.org/10.1097/00005768-200305001-01571>.
 114. Abel MG, Peritore N, Shapiro R, Mullineaux DR, Rodriguez K, Hannon JC. A comprehensive evaluation of motion sensor step-counting error. *Appl Physiol Nutr Metab*. 2011;36(1):166–170. <https://doi.org/10.1139/H10-095>.
 115. Le Masurier GC, Lee SM, Tudor-Locke C. Motion sensor accuracy under controlled and free-living conditions. *Med Sci Sports Exerc*. 2004;36(5):905–910. <https://doi.org/10.1249/01.MSS.0000126777.50188.73>.
 116. Crouter SE, Schneider PL, Bassett Jr DR. Spring-levered versus piezoelectric pedometer accuracy in overweight and obese adults. *Med Sci Sports Exerc*. 2005;37(10):1673–1679. <https://doi.org/10.1249/01.mss.0000181677.36658.a8>.
 117. Bassey E, Dallosso H, Fentem P, Irving J, Patrick J. Validation of a simple mechanical accelerometer (pedometer) for the estimation of walking activity. *Eur J Appl Physiol*. 1987;56(3):323–330. <https://doi.org/10.1007/BF00690900>.
 118. Normand MP. Increasing physical activity through self monitoring, goal setting, and feedback. *Behav Interv*. 2008;23(4):227–236. <https://doi.org/10.1002/bin.267>.