


# Antibiotics

For best viewing, please access on your desktop computer.



Cephalosporin or type I mediated penicillin allergy\*:

Ciprofloxacin replaces cephalosporin or piperacillin/Tazobactam

\*Type I mediated hypersensitivity includes hives, angioedema and/or anaphylaxis

\*\*Patient receiving the following medications:

- \* Oral or SQ methotrexate >5 mg
- \* Prednisone 2 mg/kg/day or ≥ 20 mg daily (> 2 weeks)
- \* Cyclophosphamide
- \* Rituximab
- \* Mycophenolate mofetil
- \* Azathioprine
- \* Anakinra
- \* Infliximab
- \* Etanercept
- \* Adalimumab

If patient experiences Toxin Mediated Syndrome: Consider addition of Clindamycin

## Consider IM Ceftriaxone if delay in IV/IO Access

*These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly these guidelines should guide care with the understanding that departures from them may be required at times.*

[Children's Mercy Kansas City](#) / [Health Care Providers](#) / [Evidence Based Practice](#) / [Clinical Practice Guidelines and Care Process Models](#) / [Sepsis ED Inpatient Algorithm](#) / Antibiotics

2401 Gillham Road, Kansas City, MO 64108

© Copyright 2022 The Children's Mercy Hospital

Language Assistance Available

[Notice of Nondiscrimination](#) [Español](#) [繁體中文](#) [Tiếng Việt](#) [Serbo-Croatian](#) [Deutsch](#) [한국어](#) [Français](#) [Laotian](#) [العربية](#) [Tagalog](#) [Burmese](#) [Persian \(Farsi\)](#)  
[Deutsch](#) [Oromo](#) [Português](#) [Amharic](#) [日本語](#) [Русский](#) [Hmong](#) [Swahili](#)