

Indications For Hematology Consultation

- Initiation of therapy for patients
 - Age < 30 days
 - Baseline INR \geq 1.2 prior to initiation of warfarin
 - Impaired renal function
- Maintenance of anticoagulation therapy with
 - Delay in reaching therapeutic anticoagulation
 - Progression of thrombus
 - Concern for heparin induced thrombocytopenia (HIT)
 - Hemorrhage and need for antidote
- Surgery or invasive procedure in patients with:
 - Mechanical/prosthetic mitral valves
 - Atrial fibrillation
 - Recent/recurrent thromboembolism
- Use of LMW heparin other than enoxaparin

References:

David, M., et al. Heparin and LMWH in Children. Thrombosis Interest Group of Canada. January 2007. <http://www.tigc.org/eguidelines/heparinchild07.htm>. Accessed 11/15/08.

Lexi-Drugs Online/Pediatric Lexi-Drugs Online, Enoxaparin, <http://online.lexi.com/crlsql/servlet/crlonline>, Copyright © 1978-2008 Lexi-Comp, Inc, Hudson, OH 44236

Monagle P, Chan AKC, Goldenberg NA, Ichord RN, Journeycake JM, Nowak-Gottl U, Vesely SK. Antithrombotic therapy in neonates and children: antithrombotic therapy and prevention of thrombosis, 9th ed: American College of Chest Physicians evidence-based clinical practice guidelines. Chest 2012; 141: e737S –e801S.

Roach ES. Golomb MR. Adams R. Biller J. Daniels S. Deveber G. Ferriero D. Jones BV.

Kirkham FJ. Scott RM. Smith ER. American Heart Association Stroke Council. Council on Cardiovascular Disease in the Young. Management of stroke in infants and children: a scientific statement from a Special Writing Group of the American Heart Association Stroke Council and the Council on Cardiovascular Disease in the Young. [Journal Article] Stroke. 39(9):2644-91, 2008 Sep.

These guidelines do not establish a standard of care to be followed in every case. It is recognized that each case is different and those individuals involved in providing health care are expected to use their judgment in determining what is in the best interests of the patient based on the circumstances existing at the time. It is impossible to anticipate all possible situations that may exist and to prepare guidelines for each. Accordingly these guidelines should guide care with the understanding that departures from them may be required at times.

[Children's Mercy Kansas City](#) / [Health Care Providers](#) / [Evidence Based Practice](#) / [Clinical Practice Guidelines and Care Process Models](#) / [Anticoagulation Therapy: Standard Heparin Algorithm](#) / Indications For Hematology Consultation

2401 Gillham Road, Kansas City, MO 64108

© Copyright 2022 The Children's Mercy Hospital

Language Assistance Available

[Notice of Nondiscrimination](#) [Español](#) [繁體中文](#) [Tiếng Việt](#) [Serbo-Croatian](#) [Deutsch](#) [한국어](#) [Français](#) [Laotian](#) [العربية](#) [Tagalog](#) [Burmese](#)
[Persian \(Farsi\)](#) [Deutsch](#) [Oromo](#) [Português](#) [Amharic](#) [日本語](#) [Русский](#) [Hmong](#) [Swahili](#)